

PII: S0040-4039(97)00117-2

## Allenyl(vinyl)methane Photochemistry. Photochemistry of 4,4,7-Trimethyl-5-phenyl-2,5,6-octatrienate Derivatives

Takashi Tsuno,\* Hidetaka Hoshino, and Kunio Sugiyama

Department of Industrial Chemistry, College of Industrial Technology, Nihon University, Narashino-shi, Chiba 275, Japan

Abstract: The direct photolysis of the diester in the title compounds gave a cross conjugated triene, [2+2] cycloadduct, pyran, and cyclopropene. The direct photolysis of the monoester afforded a cross conjugated triene, [2+2] cycloadduct, cyclopropene, and its Z-form. The sensitized photolysis of these esters afforded the trienes and [2+2] cycloadducts, which were derived from the triplet excited state of the different chromophores. © 1997 Elsevier Science Ltd. All rights reserved.

Previously, we reported the photochemistry of alkylidenemalonates and alkylidene Meldrum's acids possessing an allenyl group at the  $\gamma$ -position which is either the 1,2-propadienyl, 1,2-butadieneyl, or 3-methyl-1,2-butadienyl group, and these esters underwent the photoreaction as described in Scheme 1.<sup>1.2</sup> The photoreactions of these esters occur from the excited state of the vinyl chromophore or from the CT excited state between both chromophores and presented the possibility of new multiplicity control on the photochemistry of divinymethanes.<sup>3</sup> We then noted the photochemistry of allenyl(vinyl)methane 1a and 1b in which the allenyl group is attached to a phenyl group. It is expected that these compounds 1a and 1b will undergo the reaction not only via the above photoexcitation but also the allene excitation. We now wish to report the photochemistry of 1a and 1b.



The diester  $1a^4$  was obtained by the condensation of 2,2,5-trimethyl-3-phenyl-3,4-hexadienal<sup>5</sup> with dimethyl malonate using Lehnert's method.<sup>6</sup> The monoester 1b was also prepared by the reaction of the above aldehyde with methyl diethylphosphonoacetate in the presence of sodium hydride. Analogous to our previous reported allenyl(vinyl)methanes,<sup>1,2</sup> the esters 1a and 1b in several solvents showed a CT band in the range of 250 to 310 nm. The direct photolysis of 1a (0.005 mol dm<sup>-3</sup>) in hexane (400 cm<sup>3</sup>) using a 6-W low-pressure mercury lamp with a quartz jacket (254 nm irradiation) under argon gave a cross conjugated triene 2a, intramolecular [2+2] cycloadduct 3a, pyran 4, and cyclopropene 5a in 4%, 15%, 18%, and 4% isolated yields, respectively.<sup>7</sup> In a similar manner, the direct photolysis of 1b gave 2b, 3b, 5b, and 6 in 1%, 25%, 15%, and 2% isolated yields, respectively. In the case of the CT excitation (> 280 nm irradiation) of 1a and 1b, these photoproducts were not obtained. The acetone-sensitized photolysis of 1a gave 2a and 3a in 12% and 3% yields, respectively, as intramolecular photoproducts. Moreover, the acetone-sensitized photolysis of 1b afforded 2b (7%) and 3b (14%), respectively. During both acetone sensitizations, polymeric materials were significantly produced. The structures of the photoproducts were established from spectroscopic data and HRMS. Further, 3a was established by X-ray crystallographic analysis (Fig. 1).<sup>8</sup>





Fig. 1. ORTEP drawing of the [2+2] cycloadduct **3a**.

Table 1 shows quantum yields for the isomerization of 1a and 1b under the direct or sensitized irradiation. In the direct photolyses, the isomerization to 5a or 5b has a very low quantum efficiency and their formations showed an induction period by GLC during the direct photolyses of 1a and 1b. Moreover, the isolated [2+2] cycloadducts 3a and 3b were irradiated to afford 5a and 5b. Therefore, 5a and 5b are apparently photoproducts of 3a and 3b. Photochemical *E-Z* isomerization of 1b was expected, but scarcely occurred during the direct photolysis alone; the maximum yield of 6 was 3% by GLC. In the case of the direct photolyses in the presence of 2-methyl-1,3-butadiene as a quencher, the

quantum yields for the formation of the [2+2] cycloadducts, pyran, and Z-form are within experimental errors in comparison with the non-additive conditions, although those of 2a and 2b decrease in ca. 35%  $(1a\rightarrow 2a)$  and 38%  $(1b\rightarrow 2b)$ , respectively. Acetone and acetophenone are good sensitizers for the isomerization to the trienes and [2+2] cycloadducts. Benzophenone has sensitizing effects on their isomerization, although the qua. Im

Substrate	λ	Additive $(E_T/kJ \text{ mol}^{-1})^{b}$		Quantum Yields				
1	nm	-	$\Phi_{1 \rightarrow 2}$	$\Phi_{1 \rightarrow 3}$	$\Phi_{i \rightarrow 4}$	$\Phi_{1 \rightarrow 5}$	Φ <sub>1→6</sub>	
1a	254	none <sup>c)</sup>	0.013	0.043	0.030	< 0.0001		
		2-Methyl-1,3-butadiene (251) <sup>c,d</sup>	0.0085	0.041	0.028	< 0.0001		
	313	Acetone (377-334) <sup>c)</sup>	0.25	0.027				
	365	Acetophenone (310) <sup>c)</sup>	0.37	0.016				
		Benzophenone (287) <sup>c)</sup>	0.29	0.0018				
		2-Benzoylpyridine (279) <sup>c)</sup>	0.16	<0.0001				
		Anthraquinone (259) <sup>e)</sup>	0.082	<0.0001				
		4-Phenylacetophenone (254) <sup>c)</sup>	0.012	<0.0001				
		1-Acetonaphthone (236) <sup>c)</sup>	0.0031	<0.0001				
1b	254	none <sup>c)</sup>	0.0065	0.14		0.0096	0.014	
		2-Methyl-1,3-butadiene (251) <sup>c,f)</sup>	0.0040	0.15		0.0095	0.012	
	313	Acetone (377-334) <sup>c)</sup>	0.026	0.054				
	365	Acetophenone (310) <sup>c)</sup>	0.045	0.016				
		Benzophenone (287) <sup>c)</sup>	0.040	0.0022				
		2-Benzoylpyridine (279) <sup>c)</sup>	0.021	<0.0001				
		Anthraquinone (259) <sup>e)</sup>	0.0066	<0.0001				
		4-Phenylacetophenone (254) <sup>c)</sup>	0.0055	<0.0001				
		1-Acetonaphthone (236) <sup>c)</sup>	0.0028	<0.0001				

Table 1. Quantum Yields<sup>a)</sup> for the Isomerization of 1a and 1b

a) Total quantum yields for the formation of the photoproducts corresponded with the quantum yield for the disappearance of 1a or 1b below the conversion of 20%. b) Sensitizers absorb > 95% of incident light. c) In hexane. d) Quencher concn.: 0.03 mol dm<sup>-3</sup>. e) In benzene. f) Quencher concn.: 0.07 mol dm<sup>-3</sup>. yields of the formation of the [2+2] cycloadducts 3a and 3b are smaller values than those obtained with use of acetone or acetophenone. Furthermore, 2-benzoylpyridine, anthraquinone, 4-phenylacetophenone, and 1-acetonaphthone having triplet energy less than benzophenone show sensitizing effects for the formation of the triene only and the later two sensitizers have slight quantum efficiency.

Because these isomerizations did not occur by the CT excitation of 1a and 1b, it is considered that the photoproducts, except 5a and 5b, were derived from the excited state of either the allenyl or  $\alpha,\beta$ -unsaturated ester chromophore. Sensitizing effects suggest that the trienes and [2+2] cycloadducts are derived from the triplet excited state of a different chromophore in the substrates. Based on our previous report.<sup>1</sup> the triplet energy required for the formation of the [2+2] cycloadducts is estimated to be ca. 300 kJ mol<sup>-1</sup>; hence, the triplet excited state of the vinvl chromophore in the substrates affords the [2+2] cycloadducts. On the other hand, the triplet energy required for the formation of the trienes 2a or 2b is estimated to be ca. 260 kJ mol<sup>-1</sup>. This triplet energy is nearly value to the triplet energy of the phenyl-substituted allene.<sup>9</sup> Hence, acetone or acetophenone having a higher triplet energy competitively affects both chromophores in 1a or 1b, although the allenvi chromophore is excited effectively by sensitizers in the range of 260 to 290 kJ mol<sup>-1</sup>. By "the rule of five", <sup>10,11</sup> it is considered that the vinyl chromophore excited state in 1a and 1b initially bonds between the 2- and 6-positions to yield a cyclopenta-1,3-diyl radical 7 from which ring closure give the [2+2] cycloadduct (Scheme 2).<sup>12</sup> The formation of 3b during the direct irradiation predominant rather than during the sensitized irradiation, and, further, by the quenching effect it seems that the 12+21 cycloaddition during the direct irradiation involves the singlet pathway For the formation of the trienes, it is considered that the allenvl chromophore excited state bonds between the 3and 5-positions to afford 8, which affords a spirobiradical 9 and subsequent cyclopropane ring cleavage give the trienes 2a and 2b (Scheme 2). This mechanism was supported by PM3 MO calculation:<sup>13</sup> the optimal structure of biradical 9 (X=H) corresponded to the structure of 2b. Furthermore, it is considered that 4 was derived via the photochemical oxa-Diels-Alder reaction by the singlet excited state of the carbonyl at the cis-position of 1a 2a,14



Generally, the photoreaction via the triplet excited state of bichromophoric compounds takes place from the lower triplet of either chromophore by an intramolecular energy transfer.<sup>15</sup> However, it is interesting that, despite the short length between the allenyl and vinyl chromophores in **1a** and **1b**, the triene and [2+2] cycloadducts are derived from the triplet excited state of different chromophores.

## ACKNOWLEDGMENT

The authors are indebted to Professor Michikazu Yoshioka and Professor Sigekazu Kumakura of Faculty of Science, Saitama University, for their helpful discussions and for providing us with the X-ray diffractometer.

## **REFERENCES AND NOTES**

- 1. Sugiyama, K.; Tsuno, T. Chem. Express 1992, 7, 929.
- 2. a) Tsuno, T.; Sugiyama, K. Bull. Chem. Soc. Jpn. 1995, 68, 3175; b) idem Chem. Lett. 1991, 503.
- 3. For reports for the multiplicity control of acyclic divinylmethanes, see Ref. 2a.
- 4. 1a: IR (neat) 1965, 1734, 1638 cm<sup>-1</sup>; <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  1.31 (6H, s), 1.81 (6H, s), 3.68 (3H, s), 3.77 (3H, s), 7.20 (1H, s), 7.25-7.28 (5H, m); <sup>13</sup>C NMR (CDCl<sub>3</sub>)  $\delta$  20.4, 27.5, 40.5, 52.5, 54.6, 98.9, 110.8, 125.0, 126.4, 127.9, 128.3, 136.9, 154.7, 166.9, 201.5; MS *m/z* (rel intensity) 328 (M<sup>+</sup>, 6), 143 (100). HRMS Found: *m/z* 328.1650. Calcd for C<sub>20</sub>H<sub>24</sub>O<sub>4</sub>: 328.1675.
- 5. The allenic aldehyde was prepared from the reaction of 3-methyl-1-phenyl-1-butyne-3-ol with isobutyraldehyde in the presence of TsOH in benzene.
- 6. Lehnert. W. Tetrahedron 1973, 29, 635.
- 7 2a: IR (neat) 1737, 1597 cm<sup>-1</sup>; <sup>1</sup>H NMR (CDCl<sub>2</sub>) δ 1.66 (3H, s), 1.69 (6H, s), 1.85 (3H, s), 2.02 (3H, s), 3.51 (3H, s), 3.75 (3H, s), 7.24 (5H, s), 7.85 (1H, s); <sup>13</sup>C NMR (CDCl,) § 21.2, 22.0, 23.1, 23.3, 51.9. 52.5. 122.6. 126.0, 127.3, 129.3, 129.8, 131.9, 133.2, 135.0, 140.5, 147.6, 165.9, 166.5. HRMS Found: m/z 328.1725. Calcd for C<sub>20</sub>H<sub>24</sub>Q; 328.1675. 3a: mp 102-105 °C (hexane); IR (neat) 1733, 1602 cm<sup>-1</sup>; <sup>1</sup>H NMR (CDCl<sub>3</sub>) & 0.93 (3H, s), 1.27 (3H, s), 1.57 (3H, s), 1.73 (3H, s), 2.29 (1H, s), 3.77 (3H, s), 3.84 (3H, s), 7.22-7.50 (5H, m); <sup>13</sup>C NMR (CDCL, ) δ 18.0, 19.1, 20.1, 23.7, 29.2, 36.4, 48.1, 52.1, 52.6, 57.4, 126.7, 127.9, 129.0, 130.2, 131.2, 137.7, 170.9, 171.3. HRMS Found: 328.1731. Calcd for C<sub>20</sub>H<sub>24</sub>O<sub>4</sub>: 328.1675. 4: IR (neat) 1720, 1682 cm<sup>-1</sup>; <sup>1</sup>H NMR (CDCl<sub>3</sub>) & 1.04 (3H, s), 1.27 (3H, s), 1.68 (3H, s), 1,73 (3H, s, CH<sub>3</sub>), 1.78 (1H, s), 3.70 (3H, s), 3.98 (3H, s), 7.21-7.36 (5H, m); <sup>13</sup>C NMR (CDCl,) & 17.5, 19.9, 24.4, 29.4, 29.7, 32.1, 50.9, 55.6, 79.4, 118.2, 126.1, 128.3, 130.2, 14, 4, 144.6, 163.5, 167.8. HRMS Found: m/z 328.1693. Calcd for C<sub>20</sub>H<sub>24</sub>O<sub>4</sub>: 328.1675. 5a: IR (neat) 1824, 1741 cm<sup>-1</sup>; <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 1.32 (6H, s), 1.54 (3H, d, J=0.8 Hz), 1.82 (3H, d, J=1.1 Hz), 3.78 (6H, s), 6.08 (1H, septet, J=1.1 Hz), 7.26-7.49 (5H, m); <sup>13</sup>C NMR (CDCl<sub>3</sub>) & 19.0, 23.1, 24.7, 26.5, 53.0, 58.3, 120.3, 123.4, 125.1, 128.3, 128.5, 129.1, 129.6, 138.6, 169.3. HRMS Found m/z 328, 1670. Calcd for C<sub>20</sub>H<sub>24</sub>O<sub>4</sub>: 328.1675.
- 8. Crystal data for **3a**:  $C_{20}H_{24}O_4$ , Mw = 328.41, Monoclinic, space group P21/c (#14); a = 7.639(5), b = 20.832(9), c = 22.658(9) Å,  $\beta = 104.57(4)^\circ$ , V = 1795.6 (16) Å<sup>3</sup>, Z = 4, Dc = 1.20 g cm<sup>3</sup>, Cu-K $\alpha$  ( $\lambda = 0.71073$  Å<sup>-1</sup>), R = 0.0482, Rw = 0.0804 for 5033 unique reflections.
- 9. The triplet of 3-phenyl-1,2-butadiene is quenched by 1,3-pentadiene (E<sub>r</sub> =247 kJ mol<sup>-1</sup>),<sup>9a</sup> while the triplet of xanthene-9-thione (274 kJ mol<sup>-1</sup>) is quenched by phenylallene. Hence, it is considered that the triplet energy of the monophenyl-substituted allene lies in a range of 250 to 270 kJ mol<sup>-1</sup>.<sup>9b</sup> a) Fujita, K.; Matsui, K.; Shono, T. J. Am. Chem. Soc. 1975, 97, 6256; b) Kamphuis, J.; Bos, H. J. T.; Visser, R. J.; Huizer, B. H.; Varma, C. A. G. O. J. Chem. Soc., Perkin Trans. 2 1986, 1867.
- 10. Recently, "the rule of five" for the intramolecular [2+2] cycloaddition of cycloalkenones with term nal alkenes has been proved by a biradical trapping experiment using hydrogen selenide. Maradyn, D. J.; Weedon, A, C. J. Am. Chem. Soc. 1995, 117, 5359.
- 11. Crimmins, M. T. Chem. Rev. 1988, 88, 1453; Brian, T.; McMurry, H.; McKenna, B. J. Chem. Soc., Perkin Trans. 1, 1991, 811; Srinivasan, R.; Carlough, K. H. J. Am. Chem. Soc. 1967, 89, 4932.
- 12. We confirmed that the biradical intermediate for the intramolecular [2+2] cycloaddition of allenyl(vinyl)methane is a cyclopenta-1,3-diyl radical by the photochemistry of methyl 4,4,7-trimetyl-2,5,6-octatrienate in the presence of hydrogen selenide. Tsuno, T.; Sugiyama, K. unpublished result.
- 13. MO calculation was performed using UHF PM3 method of Mac Spartan Plus of Wavefunction, Inc.
- 14. Sugiyama. K.; Yoshida, M.; Tsuno. T. Heterocycles 1994, 38, 1721.
- 15. For review of photochemistry of bichromophoric molecules, see: De Schryver, F. C.; Bones, N.; Put, J. Advances in Photochemistry; Pitt, Jr. J. N.; Hammond, G. S.; Gollnick, K. Eds.; John Wiley and Sons, Inc.: New York, 1977, Vol. 10, pp. 359-465.

(Received in Japan 29 October 1996; revised 12 December 1996; accepted 16 January 1997)