

S0040-4039(96)00386-3

Trimethylsilylcyanation of Aldehydes and Ketones Catalyzed by Diorganotin Dichlorides

James K. Whitesell* and Richard Apodaca

Department of Chemistry and Biochemistry University of Texas at Austin Austin, TX 78712, USA

Abstract: Dibutyltin dichloride and diphenyltin dichloride functioned as effective catalysts for the trimethylsilylcyanation of aldehydes and ketones. Isolated yields ranged from 71% to 97%. Copyright © 1996 Elsevier Science Ltd

The development of catalysts for the addition reaction of cyanotrialkylsilanes with carbonyl compounds is an area of active study^{1, 2} due to the synthetic versatility of *O*-silylated cyanohydrins.³ Although diorganotin dihalides offer several attractive features⁴ which could be exploited in catalytic systems (for example: mild Lewis acidity; ease of synthesis; generally monomeric solution state structures; strong carbon-metal bonds relative to many organometallics; and relative stability to air), the use of these organometallic compounds as catalysts in preparative organic reactions is surprisingly uncommon.⁵ Recently, Scholl and Fu reported that tributyltin cyanide was an efficient catalysts for the trialkylsilylcyanation of aldehydes and activated ketones.^{1a} As part of an ongoing program to develop practical, enantioselective, catalytic reactions based on diorganotin dihalides, we have found dibutyltin dichloride and diphenyltin dichloride to function as effective catalysts for the addition of trimethylsilyl cyanide (TMSCN) to aldehydes and unactivated ketones.

$$R' + 1.3 \text{ Me}_{3}\text{SiCN} \xrightarrow{R_{2}\text{SnCl}_{2}} Me_{3}\text{SiO} \xrightarrow{CN} (1)$$

Treatment of a variety of aldehydes and ketones with TMSCN and a catalytic amount of diorganotin dihalide neat at 0 °C followed by warming to room temperature⁶ furnished the expected *O*-silylated cyanohydrins in good yield (Eq. 1, Table 1).⁷ Both sterically hindered and conjugated aldehydes were suitable substrates (entries 2-6, and 8), as were dialkyl and aryl alkyl ketones (entries 10-12). Diphenyltin dichloride was a more reactive catalyst than dibutyltin dichloride (entries 4, 5 and 11, 12). All uncatalyzed reactions exhibited less than 4% conversion under otherwise identical conditions except for pivaldehyde, which showed 20% conversion.⁸ In the case of benzaldehyde, freshly-distilled dibutyltin dichloride afforded results (entry 3) comparable to those obtained with dibutyltin dichloride which was used as received and handled in an air atmosphere (entry 2),

demonstrating that the catalyst is air-tolerant and suggesting that catalytic activity is not due to an inorganic impurity in the commercial organotin.

	R ₂ SnCl ₂ catalyst				
Entry	Substrate	R=	Mol%	Time (h)	Yield(%) ^a
1 ^b	<i>с</i> -С ₆ Н ₁₁ СНО	Bu	10	0.5	83 ^c
2	PhCHO	Bu	10	0.5	90
3 ^d	PhCHO	Bu	10	0.5	97
4	PhCHO	Bu	2	0.5	75 ^e
5	PhCHO	Ph	2	0.5	90
6 ^{b,f}	<i>t</i> -BuCHO	Bu	10	5.5	71
7 ^b	PhCHMeCHO	Bu	10	1.5	78 ^g
8	(E)-PhCH=CHCHO	Bu	10	0.5	86 ^h
9b	<i>п</i> -С ₇ Н ₁₅ СНО	Bu	10	0.2	85
10	<i>п</i> -С ₆ Н ₁₃ СОМе	Bu	10	22	93
11	PhCOMe	Bu	10	45	93 ^e
12	PhCOMe	Ph	4 ⁱ	1.8	95

 Table 1. Yields and reaction times for the diorganotin dihalide-catalyzed trimethylsilylcyanation of aldehydes and ketones.

a) Isolated yields of spectroscopically pure products are reported unless otherwise noted. All reactions were run to 100% conversion as determined by ¹H NMR of the crude product unless otherwise noted. All products were identified by comparison of their ¹H and ¹³C NMR spectra with samples prepared with ZnI₂ catalyst.⁹ b) The aldehyde was treated as described in reference 8. c) Traces of an unidentified impurity were present in the isolated product. d) The catalyst was distilled before use. e) Because of incomplete conversion, yield was estimated from the ¹H NMR spectrum of the crude reaction mixture. f) The uncatalyzed reaction proceeded to 20% conversion. g) A 1:1 ratio of diasteriomers was obtained. h) Exclusive 1,2-addition was observed. i) This loading represents the solubility limit of the catalyst in the reaction medium.

Although a detailed understanding of this reaction will require further studies, previous reports regarding organotin chemistry suggest that a tin cyanide may be a catalytically active species:¹⁰ (1) dimethyltin dicyanide has been prepared from the reaction of dimethyltin dichloride and TMSCN;^{11, 12} (2) tributyltin cyanide adds to aldehydes much more rapidly than does TMSCN;¹³ and (3) organotin alkoxides react with trialkylsilyl halides to afford the organotin halide and silylated alcohol.^{1a,14} A plausible catalytic cycle which takes these observations into account is depicted in Figure 1.



Figure 1. Possible catalytic cycle for the trimethylsilylcyanation of aldehydes and ketones catalyzed by diorganotin dichlorides.

Given our preliminary mechanistic considerations, the advantageous features of diorganotin dihalides, and the fact that a number of nonracemic chiral diorganotin dihalides have already been reported,^{5a, 5f, 15} we believe that the development of an enantioselective version of this new catalytic system may be feasible. Efforts directed toward this end are in progress.

Acknowledgments: Financial support of this research by the National Institutes of Health (Grant GM-31750) and by the Robert A. Welch Foundation (Grant F-626) are gratefully acknowledged. R.A. wishes to thank Texaco and the University of Texas Department of Chemistry and Biochemistry for graduate fellowships.

REFERENCES AND NOTES

- For recent reports of achiral homogeneous catalysts, see: (a) tributyltin cyanide: Scholl, M.; Fu, G.C. J. Org. Chem. 1994, 59, 7178. (b) tetracyanoethylene: Miura, T.; Masaki, Y. J. Chem. Soc., Perkin Trans. 1 1994, 1659. (c) [HC(py)₃W(NO)₂](SbF₆)₂: Faller, J.W.; Gunderson, L.-L. Tetrahedron Lett. 1993, 34, 2275. (d) LiClO₄: Reetz, M.T.; Fox, D.N.A. Tetrahedron Lett. 1993, 34, 1119. (e) Eu(fod)₃: Gu, J.-H.; Okamoto, M.; Terada, M.; Mikami, K.; Nakai, T. Chem. Lett. 1992, 1169. (f) Nitrogen bases: Kobayashi, S.; Tsuchiya, Y.; Mukaiyama, T. Chem. Lett. 1991, 537.
- Several enantioselective catalysts for the addition of TMSCN to aldehydes have been reported. For recent examples, see: (a) Bolm, C.; Müller, P. *Tetrahedron Lett.* 1995, *36*, 1625. (b) Hayashi, M.; Inoue, T.; Miyamato, Y.; Oguni, N. *Tetrahedron* 1994, *50*, 4385. For reviews on catalytic enantioselective cyanohydrin syntheses, see reference 3c and: (c) North, M. Synlett 1993, 807.

- For recent examples see: (a) Brown, R.F.C.; Donohue, A.C.; Jackson, W.R.; McCarthy, T.D. *Tetrahedron* 1994, 48, 13739. (b) Cainelli, G.; Panunzio, M.; Contento, M.; Giacomini, D.; Mezzina, E.; Giovagnoli, D. *Tetrahedron* 1993, 49, 3809. For a recent review of stereoselective reactions of cyanohydrins and cyanohydrin derivatives, see: (c) Effenberger, F. Angew. Chem. Int. Ed. Engl. 1994, 33, 1555. For a review of O-protected cyanohydrins as acyl anion equivalents, see: (d) Allbright, J.D. *Tetrahedron* 1983, 20, 3207.
- For reviews on organotin chemistry, see: (a) Omae, I. Organotin Chemistry; Elsevier: Amsterdam, 1989.
 (b) Davies, A.G.; Smith, P.J. Comprehensive Organometallic Chemistry; Wilkinson, G., Stone, G.A., Abel, E.W., Eds.; Pergamon Press: New York, 1982; Vol. 2, p 519.
- For examples, see: (a) Vedejs, E.; Erdman, D.E.; Powell, D.R. J. Org. Chem. 1993, 58, 2840.
 (b) Shibata, I.; Yoshimura, N.; Baba, A.; Matsuda, H. Tetrahedron Lett. 1992, 33, 7149.
 (b) Yano, K.; Baba, A.; Matsuda, H. Chem. Lett. 1991, 1181.
 (c) Kobayashi, S.; Uchiro, H.; Fujishita, Y.; Shiina, I.; Mukaiyama, T. J. Am. Chem. Soc. 1991, 113, 4247.
 (d) Yano, K.; Amishiro, N.; Baba, A.; Matsuda, H. Bull. Chem. Soc. Jpn. 1991, 64, 2661.
 (e) Lowenthal, R.E.; Abiko, A.; Masamune, S. Tetrahedron Lett. 1990, 31, 6005.
 (f) Krishnamurti, R. Ph.D. Dissertation, State University of New York at Albany, 1989.
 (g) Bacaloglu, R.; Cotarca, L.; Marcu, N.; Tölgyi, S. J. Prakt. Chem. 1988, 330, 541.
- 6. Reactions with aldehydes were generally mildly exothermic. Initial cooling was employed to prevent significant temperature increase.
- 7. General experimental procedure: Trimethylsilyl cyanide (CAUTION: *Toxic!*, 3.20 mmol, 1.3 eq) was added to the untreated organotin catalyst at room temperature under nitrogen. The homogeneous mixture was frozen in an ice bath after which the untreated aldehyde or ketone (2.46 mmol, 1 eq) was added and the ice bath was removed. After the indicated time, the resulting oil was pipetted onto a short plug of Florisil[®] and eluted with Skelly B. The catalyst-free product was obtained by evaporation of the solvent at water aspirator pressure.
- 8. We believe that trace amounts of carboxylic acids resulting from air-oxidation of the starting aldehydes act as catalysts in the absence of organotins. We found that washing the aldehyde sequentially with equal volumes of saturated NaHCO₃, water, 5% HCl, and water followed by vacuum distillation (see: Perrin, D.D., Armarego, L.F., Perrin, D.R.; *Purification of Laboratory Chemicals*; Pergamon Press: New York, 1966; p 350) minimized the uncatalyzed reaction. In the case of octanal, simply distilling the aldehyde had little effect. We observed substantial acceleration of the addition reaction of untreated octanal with TMSCN in the presence of 10 mol% acetic acid. There was no significant acceleration in the presence of 10 mol% ethanol. We also found that the reaction of untreated octanal with TMSCN alone could be inhibited by adding 4 mol% Me₃SiCl. This amount of additive had a negligable effect on the rate of the Bu₂SnCl₂-catalyzed reaction. The role Me₃SiCl in these reactions is not yet understood.
- 9. Evans, D.A.; Truesdale, L.K.; Carroll, G.L. J. Chem. Soc., Chem. Commun. 1973, 55.
- 10. For related discussions, see reference 1a and: Saito, S.; Yamashita, S.; Nishikawa, T.; Yokoyama, Y.; Inaba, M.; Moriwake, T. *Tetrahedron Lett.* **1989**, *30*, 4153. Activation of the carbonyl compound by the diorganotin dichloride in a Lewis acid sense is also possible.
- 11. Konnert, J.; Britton, D.; Chow, Y.M. Acta Cryst. 1972, B28, 180.
- 12. We have found by ¹H NMR spectroscopy that an equimolar mixture of Bu₂SnCl₂ and Me₃SiCN in CDCl₃ at room temperature contains approximately 5% Me₃SiCl.
- 13. Herranz, R.; Castro-Pichel, J.; García-Lopéz, T. Synthesis 1989, 703.
- 14. (a) Leigh, D.A.; Martin, R.P.; Smart, J.P.; Truscello, A.M. J. Chem. Soc., Chem. Commun. 1994, 1373. (b) Reginato, G.; Ricci, A.; Roelens, S.; Scapecci, S. J. Org. Chem. 1990, 55, 5132.
- (a) Krishnamurti, R.; Kuivila, H.G. J. Org. Chem. 1986, 51, 4947. (b) Schumann, H.; Wassermann, B.C.; Hahn, F.E. Organometallics 1992, 11, 2803.

(Received in USA 12 December 1995; revised 13 February 1996; accepted 15 February 1996)