

Nucleosides, Nucleotides and Nucleic Acids

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/lncn20>

SYNTHESIS OF 6-FORMYLURIDINE 5'-MONOPHOSPHATE: 6-FORMYL-UMP

Michael P. Groziak^a

^a Pharmaceutical Discovery Division, SRI International, Menlo Park, California, 94025-3493, U.S.A.

Published online: 07 Feb 2007.

To cite this article: Michael P. Groziak (2001) SYNTHESIS OF 6-FORMYLURIDINE 5'-MONOPHOSPHATE: 6-FORMYL-UMP, Nucleosides, Nucleotides and Nucleic Acids, 20:4-7, 1003-1006, DOI: [10.1081/NCN-100002478](https://doi.org/10.1081/NCN-100002478)

To link to this article: <http://dx.doi.org/10.1081/NCN-100002478>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

SYNTHESIS OF 6-FORMYLURIDINE 5'-MONOPHOSPHATE: 6-FORMYL-UMP

Michael P. Groziak


Pharmaceutical Discovery Division, SRI International,
Menlo Park, California 94025-3493

ABSTRACT

A synthesis of 6-formyluridine 5'-monophosphate (6-formylUMP) in 6 steps starting from uridine is described. This approach should be applicable to the preparation of other O5'-phosphorylated 6-formylUrds such as 6-formylUDP and 6-formylUTP.

Previous investigations in my laboratory have demonstrated that the nucleosides 6-formyluridine (6-formylUrd), 6-formyl-2-deoxyuridine, and 6-formyl-1-(β -D-arabinofuranosyl)uracil all possess the unusual carbohydrate-like property of existing to some degree in transglycosidic cyclic hemiacetal form, even in aqueous solution (1–6). This is a consequence of both the proximity of the formyl group to certain carbohydrate hydroxyl groups and to the strong electron-withdrawing force that the 1-alkyluracil-6-yl moiety exerts on the carboxaldehyde functionality. For example, 6-formylUrd exists in water as a 2:1 mixture of the aldehyde hydrate and the (7*R*)-O5',7-intramolecular cyclic hemiacetal. In Me₂SO, it exists as a 1:10 mixture of the aldehyde and the same hemiacetal diastereomer. Finally, it exists as a 1:1 mixture of diastereomeric ethyl hemiacetals in EtOH, but 6-formylUrd crystallizes out of this solvent exclusively as the (7*R*)-O5',7-cyclic hemiacetal.

The task of preparing a 5'-nucleotide derivative of 6-formylUrd such as 6-formylUMP presents a unique obstacle to overcome: Any attempt to directly O5'-phosphorylate an O5'-unprotected version of 6-formylUrd would surely fail because of the overwhelming predominance of the transglycosidic hemiacetal form that exists in anhydrous aprotic solvents. Thus, a less straightforward route—one involving a masked aldehyde—is required for accessing 6-formylUMP or indeed




any other O5'-phosphoryl derivatives that might be desired for investigations with nucleotide-utilizing enzymes.

RESULTS AND DISCUSSION

A 6-formylUrd derivative in which the 2'-OH, 3'-OH, and 6-CHO groups are all protected with acid-labile protecting groups might well be very useful in an efficient and successful synthetic approach to 6-formylUMP. Using such a derivative, a direct phosphorylation with, for instance, $POCl_3$ would be assuredly regiospecific for the 5'-OH group. Furthermore, the phosphorylation could be followed directly by an *in situ* removal of all of the protecting groups in the last step of the synthesis, taking advantage of the HCl that would be generated during an aqueous workup procedure.

With these points in mind, an efficient 6-step synthesis of 6-formylUMP starting from uridine was developed (Scheme 1). For its greater ease of removal compared to that of an isopropylidene group, one of Reese's 2',3'-O-alkoxymethylidene groups (7) was employed for the simultaneous protection of the 2'- and 3'-OH groups in uridine. Its 5'-OH group was then protected with a TBDMS group, giving nucleoside **1**. A Miyasaka dianion (a uridine-3,6-diyldilithium species)



Scheme 1.

(8) was then generated from **1**, and this was quenched with HCO_2Me to give the carbohydrate-protected 6-formylUrd **2** in an excellent yield. In our earliest work with the 6-formylUrds, we found that their formyl group would undergo acetalation only under forcing conditions, (1) and so for the present synthesis of 6-formylUMP it instead was masked as a 1,3-diphenylimidazolidin-2-yl (DPI) moiety as was found advantageous in our most recently completed investigation (6).

This conversion of the 6-formyl group of **2** to the DPI one of **3** was the only problematic step encountered. It was conducted without the aid of an acid catalyst to avoid labilizing the acid-sensitive methoxymethylidene group. Perhaps due to a steric hindrance provided by the carbohydrate fragment or a competing intramolecular 5'-O-desilylation via silyl transfer to the hemiaminal intermediate, this conversion (not optimized) proceeded in only a 27% yield. The 5'-O-TBDMS group in **3** was removed by exposure to TBAF, and the subsequent O5'-phosphorylation of **4** with POCl_3 proceeded smoothly, as did the anticipated *in situ* removal of all the protecting groups effected simply by the introduction of water after the phosphorylation was complete. The aqueous solution of 6-formylUMP (**5**) that this procedure generated was promptly treated with NH_4OH to convert **5** to its ammonium salt



form. Extraction with Et₂O to remove the (PhNHCH₂)₂ and excess PO(OEt)₃ was followed by lyophilization, giving the target 6-formylUMP **5** as an anhydrous powder ready for desalting (9).

CONCLUSIONS

With a careful consideration for the unusual carbohydrate-like properties of 6-formylUrd, a general route to its O5'-phosphorylated derivatives like 6-formylUMP has now been developed. It should prove to be useful for the preparation of 6-formylUDP, 6-formylUTP, and other similar 5'-phosphorylated 6-formylUrd derivatives.

ACKNOWLEDGMENTS

Some of the intermediates were generated in connection with a separate investigation funded by a Shannon Award from NIH/NIGMS (grant GM56878).

REFERENCES

1. Groziak, M. P.; Koohang, A. *J. Org. Chem.* **1992**, *57*, 940–944.
2. Groziak, M. P.; Koohang, A.; Stevens, W. C.; Robinson, P. D. *J. Org. Chem.* **1993**, *58*, 4054–4060.
3. Groziak, M. P.; Lin, R.; Robinson, P. D. *Acta Crystallogr.* **1995**, *C51*, 1204–1207.
4. Groziak, M. P.; Lin, R.; Stevens, W. C.; Wotring, L. L.; Townsend, L. B.; Balzarini, J.; Witvrouw, M.; De Clercq, E. *Nucleosides Nucleotides* **1996**, *15*, 1041–1057.
5. Groziak, M. P.; Lin, R. *ARKIVOC* **2000**, *1(1)*, 33–45.
6. Groziak, M. P.; Lin, R. *Tetrahedron*, submitted.
7. Griffin, B. E.; Jarman, M.; Reese, C. B.; Sulton, J. E. *Tetrahedron* **1967**, *23*, 2301–2313.
8. Tanaka, H.; Hayakawa, H.; Miyasaka, T. *Tetrahedron* **1982**, *38*, 2635–2642.
9. Preliminary characterization data obtained on un-desalted **5**: ¹H NMR (D₂O) δ 6.07 (s, 1, H5), 5.97 (s, 1, CH(OD)₂), 5.92 (d, *J* = 4Hz, 1, H1'), 4.40 (m, 1, H3'), 4.01 (m, 1, H4'), 3.87 (m, 2, 5'-CH₂). ¹³C NMR (D₂O) δ 166 (C4), 156 and 152 (C2/C6), 101 (C5), 92 (C1'), 86 (CH(OD)₂), 82 (C4'), 72 (C2'), 69 (C3'), 64 (C5'). Low resolution DCI (direct current ionization) mass spectrum, *m/e*: 213 (phosphoribosyl-H+H⁺).



Request Permission or Order Reprints Instantly!

Interested in copying and sharing this article? In most cases, U.S. Copyright Law requires that you get permission from the article's rightsholder before using copyrighted content.

All information and materials found in this article, including but not limited to text, trademarks, patents, logos, graphics and images (the "Materials"), are the copyrighted works and other forms of intellectual property of Marcel Dekker, Inc., or its licensors. All rights not expressly granted are reserved.

Get permission to lawfully reproduce and distribute the Materials or order reprints quickly and painlessly. Simply click on the "Request Permission/Reprints Here" link below and follow the instructions. Visit the [U.S. Copyright Office](#) for information on Fair Use limitations of U.S. copyright law. Please refer to The Association of American Publishers' (AAP) website for guidelines on [Fair Use in the Classroom](#).

The Materials are for your personal use only and cannot be reformatted, reposted, resold or distributed by electronic means or otherwise without permission from Marcel Dekker, Inc. Marcel Dekker, Inc. grants you the limited right to display the Materials only on your personal computer or personal wireless device, and to copy and download single copies of such Materials provided that any copyright, trademark or other notice appearing on such Materials is also retained by, displayed, copied or downloaded as part of the Materials and is not removed or obscured, and provided you do not edit, modify, alter or enhance the Materials. Please refer to our [Website User Agreement](#) for more details.

[Order now!](#)

Reprints of this article can also be ordered at

<http://www.dekker.com/servlet/product/DOI/101081NCN100002478>