



Diels–Alder approach to the synthesis of azulene-substituted benzene derivatives. Synthesis and redox behavior of 1,2-di(6-azulenyl)tetraphenylbenzenes

Shunji Ito,^{a,*} Haruki Inabe,^a Testuo Okujima,^a Noboru Morita,^a Masataka Watanabe^b and Kimiaki Imafuku^c

^aDepartment of Chemistry, Graduate School of Science, Tohoku University, Sendai 980-8578, Japan

^bInstitute for Chemical Reaction Science, Tohoku University, Sendai 980-8577, Japan

^cDepartment of Chemistry, Faculty of Science, Kumamoto University, Kumamoto 860-8555, Japan

Received 11 July 2000; revised 18 August 2000; accepted 25 August 2000

Abstract


1,2-Di(6-azulenyl)tetraphenylbenzenes and (6-azulenyl)pentaphenylbenzenes were synthesized by Diels–Alder reaction of di(6-azulenyl)acetylenes and 6-(phenylethynyl)azulenes with tetraphenylcyclopentadienone. Mono(6-azulenyl)benzenes exhibited a reduction wave upon cyclic voltammetry (CV). In contrast to the benzene derivatives, di(6-azulenyl)benzenes showed a two-step reduction wave at similar potential region upon CV, which revealed the formation of dianions stabilized by 6-azulenyl substituents under electrochemical reduction conditions. © 2000 Elsevier Science Ltd. All rights reserved.

Keywords: ethynylazulenes; Pd-cross-coupling reaction; Diels–Alder reaction; azulenylbenzenes.

Azulene (C₁₀H₈) is a theoretically interesting compound with many unusual properties.¹ Especially, the azulene system has a tendency to stabilize cations, as well as anions owing to its remarkable polarized properties, which are utilized to construct extremely stable methyl cations.² Recently, much attention has been focused on the aromatic compounds with extended π -electron systems because of their special optical and electrical properties.³ However, to date, molecules with potentially useful electronic properties constructed by non-benzenoid aromatic compounds are fairly scarce. We have recently developed a simple route to unknown 1,3,5-tri(1-azulenyl)benzenes by tetrachlorosilane-mediated cyclotrimerization of 1-acethylazulenes.⁴ Very recently, Hafner et al. reported the ethynylation of azulenes in a five-membered ring utilizing Pd-catalyzed cross-coupling reaction of bromo- and iodoazulenes with trimethylsilylacetylene⁵ and the cyclooligomerization of the 1-ethynylazulenes catalyzed by CpCo(CO)₂ to afford 1-azulenylbenzene derivatives and (cyclobutadiene)cobalt complexes in low yields.⁶


* Corresponding author. Tel/fax: +81-22-217-7714; e-mail: ito@funorg.chem.tohoku.ac.jp

Herein we report an efficient preparation of azulene-substituted benzene derivatives by Diels–Alder reaction of di(6-azulenyl)acetylenes (**3a** and **3b**) and 6-(phenylethynyl)azulenes (**4a** and **4b**) with tetraphenylcyclopentadienone (**5**) to afford novel 6-azulenylbenzene derivatives **1a**, **1b**, **2a**, and **2b**, which are expected to have a different character from that of 1-azulenylbenzene derivatives.⁷ According to the nature of the azulene systems, **1a** and **1b** would exhibit the formation of stabilized closed shell dianions **6** by electrochemical reduction (Scheme 1). Although there are many reports concerning carbocations stabilized by 1-azulenyl substituents, little is known about carbanions stabilized by 6-azulenyl substituents because of the synthetic difficulty of the functionalization of azulene in a seven-membered ring.¹ Thus, we also report here the redox behavior of these novel 6-azulenylbenzene derivatives (**1a**, **1b**, **2a**, and **2b**) and 6-azulenylacetylenes (**3a**, **3b**, **4a**, and **4b**) examined by cyclic voltammetry (CV).




Scheme 1. Redox system for the reduction of **1a**, **1b** and that for a general structure of violenes **7**

Pd-catalyzed cross-coupling reaction of 6-bromoazulenes (**8a** and **8b**)⁸ with trimethylsilylacetylene under Sonogashira–Hagihara conditions⁹ at room temperature afforded 6-(trimethylsilylethynyl)azulenes (**9a** and **9b**) in 86 and 84% yields, respectively. It is noteworthy that the ethynylation of azulenes in the seven-membered ring readily proceeded using bromides at room temperature. Treatment of **9a** and **9b** with potassium fluoride in DMF furnished 6-ethynylazulenes (**10a** and **10b**)¹⁰ in 78 and 93% yields, respectively. Cross-coupling reaction of **10a** with **8a** using $\text{PdCl}_2(\text{PPh}_3)_2$ as a catalyst afforded an inseparable mixture (1:2.2) of di(6-azulenyl)acetylene (**3a**) and di(6-azulenyl)diacetylene (**11**), which was obtained from the oxidative coupling of **10a** by the Pd-catalyst.^{9,11} However, we found that the reaction of **10a** with **8a** utilizing $\text{Pd}(\text{PPh}_3)_4$ as a catalyst exclusively gave the desired **3a** in 98% yield. Similarly, **3b** was obtained by the reaction of **10b** with **8b** under similar conditions in 96% yield. Diels–Alder reaction of **3a** and **3b** with a large excess of **5** in diphenyl ether at 160°C under an Ar atmosphere for 1–2 days and subsequent chromatographic purification of the reaction mixture on silica gel afforded **1a** and **1b** in 13 and 91% yields, respectively (Scheme 2).¹²



Scheme 2. Synthesis of **1a** and **1b**. (a) TMSA, $[\text{PdCl}_2(\text{PPh}_3)_2]$, PPh_3 , CuI , Et_3N , toluene, rt, 2.5 h (86%) for R = H, 1.5 h (84%) for R = COOEt; (b) KF , H_2O , DMF, rt, 3 h (78%) for R = H, 3 h (93%) for R = COOEt; (c) **8a** and **8b**, $[\text{Pd}(\text{PPh}_3)_4]$, PPh_3 , CuI , Et_3N , toluene, rt, 3 h (98%) for R = H, 1.5 h (96%) for R = COOEt; (d) **5**, diphenyl ether, 160°C , 2 days (13%) for R = H, 1 day (91%) for R = COOEt

Synthesis of **2a** and **2b** as outlined in Scheme 3 also commenced with **8a** and **8b**, respectively. 6-(Phenylethynyl)azulenes (**4a** and **4b**) were prepared by the Sonogashira reaction of **8a** and **8b** with phenylacetylene in 90 and 87% yields, respectively. Diels–Alder reaction of **4a** and **4b** with a large excess of **5** in diphenyl ether at 140 and 160°C , respectively, under an Ar atmosphere for 5 days afforded **2a** and **2b** in 32 and 93% yields, respectively. The spectral features of **1a**, **1b**, **2a**, and **2b** are in agreement with the structure of these products. **1a**, **1b**, **2a**, and **2b** are remarkably stable, showing no decomposition even after several weeks at room temperature.



Scheme 3. Synthesis of **2a** and **2b**. (a) Phenylacetylene, $[\text{PdCl}_2(\text{PPh}_3)_2]$, PPh_3 , CuI , Et_3N , toluene, rt, 5 h (90%) for R = H, 9 h (87%) for R = COOEt; (b) **5**, diphenyl ether, 160°C , 1 day (32%) for R = H, 140°C , 5 days (93%) for R = COOEt

1,2-Di(6-azulenyl)benzene **1b** exhibited the voltammogram, as shown in Fig. 1(a). The voltammogram was characterized by a barely separated two-step reduction wave at $E_{1/2}$ -1.57 and -1.72 V. **1a** also showed the transfer of two electrons at -2.00 and -2.20 V upon CV. Thus, the redox system of **1a** and **1b** is depicted to be like a general structure of violenes **7**, as shown





Figure 1. Cyclic voltammograms of (a) **1b** and (b) **3b** in THF containing $n\text{-Bu}_4\text{N}^+\text{BF}_4^-$ (0.1 M) as a supporting electrolyte; V versus Ag/Ag^+ , Pt electrode, scan rate 100 mV s^{-1} , and $\text{Fc}/\text{Fc}^+ = 0.10 \text{ V}$

in Scheme 1.¹³ The less negative reduction potentials of **1b** compared with those of **1a** are attributable to the stabilization of anions by the ethoxycarbonyl groups substituted in the five-membered ring. The first reduction potentials and even the second ones of **1a** and **1b** are almost comparable with those of **2a** (-2.06 V) and **2b** (-1.63 V), respectively. Therefore, the two 6-azulenyl substituents on benzene in a 1,2 relationship increased its electron-accepting properties because of the closed-shell dianionic structure **6**. Unfortunately, the reduction waves of **1a** and **1b** showed poor reversibility, which exhibited the low stabilities of the expected radical anion and dianion **6** under the electrochemical reduction conditions. Thus, the measurement of the UV-vis spectroscopy of **1b** under electrochemical reduction conditions did not afford any evidence of the formation of the radical anion and/or dianion **7** due to the instability of these species.

1,2-Di(6-azulenyl)acetylenes **3a** and **3b** are also an example of the two-electron redox system stabilized by two 6-azulenyl substituents, as illustrated in Scheme 4. In contrast to the poor reversibility of **1a** and **1b**, **3a** and **3b** exhibited a reversible two-step reduction wave (**3a**, $E_{1/2}$ -1.41 and -1.61 V ; **3b**, $E_{1/2}$ -0.96 and -1.15 V , respectively) upon CV (Fig. 1(b)). These reduction potentials of **3a** and **3b** are less negative than those of **4a** (-1.66 V) and **4b** (-1.29 V), respectively. Thus, the two 6-azulenyl substituents of **3a** and **3b** also increase the electron-accepting properties.



Scheme 4. Redox system for the reduction of **3a** and **3b**

We have demonstrated the Pd-catalyzed ethynylation of azulenes in a seven-membered ring using bromoazulenes and the efficient preparation of novel 6-azulenylbenzene derivatives **1** and **2**. Although the radical anion and dianion of **1** were not detected by UV-vis spectroscopy due to the instability of these species, **1** was clarified to represent the presumed two-step redox system upon CV. A cyclooligomerization approach to the synthesis of novel 6-azulenylbenzene derivatives using (6-azulenyl)acetylenes by transition metal-catalyzed reaction will be published elsewhere.

References

1. Zeller, K.-P. *Houben-Weyl; Methoden der Organischen Chemie*; 4th Ed.; Azulene. Georg Thieme: Stuttgart, 1985; Vol. V, Part 2C, pp. 127–418.
2. Ito, S.; Kikuchi, S.; Morita, N.; Asao, T. *J. Org. Chem.* **1999**, *64*, 5815–5821 and references cited therein.
3. Berresheim, A. J.; Müller, M.; Müllen, K. *Chem. Rev.* **1999**, *99*, 1747–1785; Martin, R. E.; Diederich, F. *Angew. Chem. Int. Ed.* **1999**, *38*, 1350–1377.
4. Sato, K.; Yamashiro, S.; Imafuku, K.; Ito, S.; Morita, N.; Fujimori, K. *J. Chem. Res. (S)* **2000**, 334–335.
5. Fabian, K. H. H.; Elwahy, A. H. M.; Hafner, K. *Tetrahedron Lett.* **2000**, *41*, 2855–2858.
6. Elwahy, A. H. M.; Hafner, K. *Tetrahedron Lett.* **2000**, *41*, 2859–2862.
7. A part of this study was presented at CSJ Local Meeting of Tohoku District, Ishinomaki, October 3, 1999, Abstract No. 2PB65 and 78th CSJ National Meeting, Funabashi, March 29, 2000, Abstract No. 2G4 30.
8. McDonald, R. N.; Richmond, J. M.; Curtis, J. R.; Petty, H. E.; Hoskins, T. L. *J. Org. Chem.* **1976**, *41*, 1811–1821.
9. Sonogashira, K. *Comprehensive Organic Synthesis*; Trost, B. M., Fleming, I., Eds.; Pergamon Press: Oxford, 1991; Vol. 3, Chapter 2.4, pp. 521–529; Takahashi, S.; Kuroyama, Y.; Sonogashira, K.; Hagihara, N. *Synthesis* **1980**, 627–630; Sonogashira, K.; Tohda, Y.; Hagihara, N. *Tetrahedron Lett.* **1975**, 4467–4470.
10. Morita, T.; Fujita, T.; Takase, K. *Bull. Chem. Soc. Jpn.* **1980**, *53*, 1647–1651; Fujita, T.; Morita, T.; Takase, K. *Tetrahedron Lett.* **1974**, *30*, 2585–2586.
11. Rossi, R.; Carpita, A.; Bigelli, C. *Tetrahedron Lett.* **1985**, *26*, 523–526; Takahashi, A.; Endo, T.; Nozoe, S. *Chem. Pharm. Bull.* **1992**, *40*, 3181–3184.
12. Ogliaruso, M. A.; Romanelli, M. G.; Becker, E. I. *Chem. Rev.* **1965**, *65*, 261–367.
13. Deuchert, K.; Hünig, S. *Angew. Chem., Int. Ed. Engl.* **1978**, *17*, 875–958; Hünig, S.; Ort, B. *Liebigs Ann. Chem.* **1984**, 1959–1971; Hünig, S.; Ort, B. *Liebigs Ann. Chem.* **1984**, 1936–1951.