This article was downloaded by: [Australian National University] On: 01 January 2015, At: 08:45 Publisher: Taylor & Francis Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information: <u>http://www.tandfonline.com/loi/gpss20</u>

Studies on 1,10-Phenanthroline Complexes of Some Hydroxyaryltellurium Trihalides

K. K. Verma^a, Seema^a & Shashi Chhabra^a ^a Department of Chemistry, M.D. University Rohtak, Rohtak, India Published online: 03 Nov 2008.

To cite this article: K. K. Verma , Seema & Shashi Chhabra (2008) Studies on 1,10-Phenanthroline Complexes of Some HydroxyaryItellurium Trihalides, Phosphorus, Sulfur, and Silicon and the Related Elements, 183:12, 3037-3045, DOI: <u>10.1080/10426500802059760</u>

To link to this article: http://dx.doi.org/10.1080/10426500802059760

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at http://www.tandfonline.com/page/terms-and-conditions



Studies on 1,10-Phenanthroline Complexes of Some Hydroxyaryltellurium Trihalides

K. K. Verma, Seema, and Shashi Chhabra Department of Chemistry, M.D. University Rohtak, Rohtak, India

The synthesis and structural features ofnewly some synthesized 1,10-phenanthroline ofp-hydroxyphenyland3-methyl-4complexes trihalides (chlorides, bromides, hydroxyphenyltellurium and iodides) are reported. The resulting complexes have been subjected to elemental analyses, conductance and cryoscopic measurements, infra-red and proton magnetic resonance spectral studies. Solution studies reveal the weak to 1:1 electrolyte type behavior of these complexes in solution. Spectral studies indicate the linkage of phenanthroline to the tellurium atom through the nitrogen atoms. Central tellurium atom in these complexes is hexa-coordinated in an octahedral way.

Keywords p-Hydroxyphenyltellurium trihalides; 3-methyl-4-hydoxyphenyltellurium trihalides; phenanthroline complexes

INTRODUCTION

Aryltellurium trihalides are known to behave as Lewis acids and form molecular complexes with several nitrogen, oxygen, and sulfur donor bases.^{1–14} Tellurium in these complexes is penta- or hexa-coordinated, but the reports on hexa-coordinated complexes are few.^{4,9–13} A small number of such compounds are known^{6.7} to posses antimicrobial activity, as well. The hydroxyaryltellurium trihalides have two active sites to interact with the nitrogen donor bases: the acidic phenolic proton; and the acceptor tellurium atom. Earlier, we reported^{10–13} the complexes of various hydroxyaryltellurium trihalides with pyridine, 2,2'-bipyridyl, piperidine, and thiourea. The present paper describes the synthesis and characterization of 1,10-phenanthroline complexes of hydroxyaryltellurium trihalides derived from phenol and *o*-cresol.

Received 2 September 2007; accepted 5 November 2007.

The authors are thankful to Maharshi Dayanand University, Rohtak (India) for necessary facilities.

Address correspondence to K. K. Verma, Department of Chemistry, Maharshi Dayanand University, Rohtak 124001, India. E-mail: vermakk123@rediffmail.com

RESULTS AND DISCUSSION

The formation of *p*-hydroxyphenyltellurium trichloride and 3-methyl-4-hydroxyphenyl-tellurium trichloride by reaction of tellurium tetrachloride with phenol^{15,16} and *o*-cresol^{15,17} involves the electrophilic substitution of the aromatic ring, by a TeCl_3^+ group at a position *para* to the hydroxyl group:

$$R-H + TeCl_4 \rightarrow RTeCl_3 + HCl.$$
(1)

Hydroxyaryltellurium trichlorides undergo halogen-exchange $^{15-17}$ when treated with KBr or KI:

$$RTeCl_3 + 3KX \rightarrow RTeX_3 + 3KCl(X = Br, I).$$
 (2)

Reactions of these hydroxyaryltellurium trihalides with 1,10- phenanthroline in 1:1 molar ratio yield $RTeX_3$. Phen type complexes:

$$\operatorname{RTeX}_3 + \operatorname{Phen} \to \operatorname{RTeX}_3$$
. Phen. (3)

These complexes are colored crystalline solids, fairly stable in dry air, and soluble in polar organic solvents.

Conductance and Cryoscopic Studies

The molar conductance data of ca. 10^{-3} M solutions in nitrobenzene, acetone and acetonitrile (Table I) reflect the weak to 1:1 electrolytic behavior of these complexes in solution, presumably due to ionization into [RTeX₂.Phen]⁺ and X⁻ ions. Complexes of 3-methyl-4hydroxyphenyltellurium trihalides appear to be more dissociated than those of *p*-hydroxy phenyltellurium trihalides. Also, the complexes of triiodides appear to be most dissociated and those of trichlorides least dissociated. The lower values of molar conductance than those of 1:1 electrolytes,¹⁸ in some cases, may be due to incomplete dissociation or ion-pair formation.

It has been noted¹⁸ that the concentration ranges selected for such measurements are often arbitrarily chosen and that the method of determination requires the assumption of the molecular weight, which may be erroneous. The complexes that are predicted as 1:1 electrolytes by $\Lambda_{\rm M}$ data and have sufficient solubility were, therefore, investigated in acetonitrile over a wider range (1 – 5 mM), and the results subjected to the Onsager Law. Values of equivalent conductance at infinite dilution, Λ_0 and (A + ω B Λ_0) are recorded in Table I. A comparison of calculated and experimental values of (A + ω B Λ_0) shows^{17,19} the compounds to behave as 1:1 electrolytes. A higher ratio than expected reflects the incomplete dissociation or ion-pair formation. Molecular Downloaded by [Australian National University] at 08:45 01 January 2015

TABLE I Molar Conductance, Cryoscopic, and (A + ω BA_o) Data of Phenanthroline Complexes of Hydroxyaryltellurium Trihalides

| RComplexNitro- benzene p -HydroxyphenylRTeCl3.Phen14.3066.41RTeBr3.Phen24.20116.26RTeBr3.Phen19.8078.90 | at ca. 10^{-3} M, ohm ⁻¹ ci | $m^2 mol^{-1}$ | rormula weight | | | | | |
|---------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------|----------------|-------------------|-------------|----------------------|-------|-----------------------------------|---------|
| RComplexNitro- benzene p -HydroxyphenylRTeCl3.Phen14.3066.41RTeBr3.Phen24.20116.26RTeBr3.Phen24.20116.26 | | 3 | conc. range, A | verage Mol. | Λ_0 | (A) | $+\omega B \Lambda_0 \sin \theta$ | Ratio |
| RComplexbenzeneAcetone p -HydroxyphenylRTeCl3.Phen14.3066.41RTeBr3.Phen24.20116.26RTel3.Phen19.8078.90 | Nitro- | Aceto- | ò | wt. | , , | Found | aceton- | (found/ |
| $p-{\rm Hydroxyphenyl} \ {\rm RTeCl_3.Phen} \ 14.30 \ 66.41 \\ {\rm RTeBr_3.Phen} \ 24.20 \ 116.26 \\ {\rm RTel_3.Phen} \ 19.80 \ 78.90 \\ \end{array}$ | benzene Acetone n | itrile | m moles/L | Found | (ohm ⁻¹) | Calcd | itrile | calcd.) |
| $\begin{tabular}{lllllllllllllllllllllllllllllllllll$ | n 14.30 66.41 9 | 93.45 | 507.4 | 339.6 | | I | I | |
| RTeBr ₃ .Phen 24.20 116.26 RTel ₃ .Phen 19.80 78.90 | | Ŭ | 2.45 - 3.57) | | | | | |
| RTel ₃ .Phen 19.80 78.90 | n 24.20 116.26 E | 58.62 | 640.6 | 359.3 | Ι | | I | |
| RTel ₃ .Phen 19.80 78.90 | | Ŭ | 1.75 - 3.45) | | | | | |
| | 19.80 78.90 1 | .17.80 | 781.6 | 397.2 | 132 | 525.0 | 329.6 | 1.592 |
| | | Ŭ | 1.95 - 3.01) | | | | | |
| 3-Methyl-4- RTeCl ₃ .Phen 22.00 66.74 | n 22.00 66.74 1 | 12.81 | 521.4 | 295.5 | 136 | 640.0 | 332.6 | 1.924 |
| hydroxyphenyl | | Ŭ | 2.32 - 4.68) | | | | | |
| $RTeBr_{3}$. Phen 25.30 73.50 | 1 25.30 73.50 (| 62.31 | 654.6 | 328.3 | I | | | |
| | | Ŭ | 2.47 - 5.75) | | | | | |
| $RTeI_{3}$. Phen 27.50 96.30 | 27.50 96.30 1 | 27.23 | 795.6 | 413.7 | 141 | 422.2 | 336.2 | 1.255 |
| | |) | 1.09 - 3.98) | | | | | |

Values of $\Lambda_{\rm M}$ reported ¹⁸ for 1: 1 electrolyte, nitrobenzene = 20-30, acetone = 100-140 and acetonitrile = 120-160.

| R | Complex | υ0-H | $\delta_{\rm (OH)}$ | $v_{\rm C}=_{\rm N}$ |
|--------------------------|--------------------------|---------------------|---------------------|----------------------|
| <i>p</i> -Hydroxyphenyl | RTeCl ₃ .Phen | 3400 mb | 1380 w | 1630 w |
| | 0 | | 1368 w | 1614 m |
| | RTeBr ₃ .Phen | $3429 \mathrm{~sb}$ | 1385 w | 1618 m |
| | RTeI ₃ .Phen | 3401 mb | 1364 w | 1628 m |
| 3-Methyl-4-hydroxyphenyl | RTeCl ₃ .Phen | 3306 mb | 1390 w | 1625 w |
| | | | 1370 w | 1616 m |
| | RTeBr ₃ .Phen | 3435 mb | 1395 w | 1630 w |
| | | | | 1613 m |
| | $RTeI_3$.Phen | 3430 mb | 1393 w | 1626 w |

TABLE II Important IR Spectral Data (cm⁻¹) of PhenanthrolineComplexes of Hydroxyaryltellurium Trihalides

w = weak; m = medium; s = strong; and b = broad.

weights in nitrobenzene (Table I) are much lower than the formula weights, supporting the results of conductance studies.

Infra-Red Spectra

The IR spectra of phenanthroline adducts are quite complex and therefore, an attempt has been made to identify the donor sites of the ligand by comparing the spectra of the complexes with those of parent hydroxyaryltellurium trihalide and phenanthroline.

All the complexes (Table II) show a broad band at about 3400 cm⁻¹ (O-H stretching) and at about 1360-1390 cm⁻¹ (OH deformation), which reflects the non participation of hydroxyl group of hydroxvaryltellurium trihalides toward bonding with phenanthroline. Several researchers^{7-9,20-22} have discussed the IR spectra of phenanthroline and its complexes and have exploited the spectral data for identification of donor sites. In the spectra of phenanthroline complexes, strong bands are expected²⁰ in three frequency regions namely between 700 and 900 cm^{-1} , between 1125 and 1250 cm^{-1} , and between 1400 and 1650 $\rm cm^{-1}$. In the spectra of complexes of hydroxyaryltellurium trihalides, the satisfactory assignments of bands in earlier two regions are not possible due to the presence of organyl moiety of RTeX₃group. It is the third region that characteristic ring frequencies of phenanthroline occur. The intense bands at about 1500 cm^{-1} , 1590 cm^{-1} and 1423 cm^{-1} are reported ²⁰ to move towards higher frequencies on coordination, the 1423 cm⁻¹ band being least sensitive. This is the $v_{\rm C=N}$ band, which most researchers have reported $7^{-9,21,22}$ to shift towards higher frequency upon complexation. The complexes of hydroxyaryltellurium trihalides show medium to strong intensity bands around $1610-1630 \text{ cm}^{-1}$, which were not present in the parent hydroxyaryltellurium trihalides. This certainly suggests the linkage of 1,10-phenanthroline molecule to the tellurium atom of RTeX₃ through the two nitrogen atoms, thus enhancing the coordination number of tellurium to six in these complexes.

Proton Magnetic Resonance Spectra

¹H NMR spectral data for hydroxyaryltellurium trichlorides complexes are presented in Table III.

1,10- Phenanthroline shows²³ four sets of two equivalent protons each at δ 7.62 (a), 7.80 (b), 8.26 (c), and 9.21(d) ppm. Upon complexation with hydroxyaryltellurium trichlorides, these are shifted to downfield side, whereas the aryl protons of hydroxyaryltellurium moiety shift to upfield side. The OH proton of hydroxyaryl group remains intact in the complexes showing its non participation as predicted by IR studies, as well. Although a lot of mixing of signals of protons of phenanthroline and hydroxyaryl groups takes place, yet independent assignments of downfield protons (H_c and H_d) are possible. These results of ¹H NMR spectra suggest the linkage of phenanthroline to the tellurium atom through nitrogen atoms.

On the basis of above spectral evidences, tellurium appears to be hexacoordinated in these phenanthroline complexes, presumably in an octahedral environment. However, in the absence of X-ray studies it is very difficult to assign the exact geometry of the molecules and stereochemical role of the lone pair on tellurium.

EXPERIMENTAL

All preparations were carried out under an atmosphere of dry nitrogen and the solvents were dried and purified by standard methods 24,25 before use.

Preparation of Hydroxyaryltellurium Trihalides

The *p*-hydroxyphenyltellurium trichloride and 3-methyl-4-hydroxyphenyltellurium trichloride were prepared by reactions of tellurium tetrachloride with phenol and *o*-cresol, respectively.^{15–17} The corresponding tribromide and triiodide were prepared by halogen-exchange process using potassium bromide or potassium iodide.^{15–17}

| Complex | Chemical shift (δ ppm) | Assignment |
|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------|
| a O A | $\begin{array}{c} 6.84^{d} \ (2H) \\ 7.82 - 8.09^{cm} \ (6H) \\ 8.58^{d} \ (2H) \\ 9.20^{d} \ (2H) \\ 9.80^{b} \ (1H) \end{array}$ | H _B H _A , H _a , and H _b H _c H _d OH |
| <i>p</i> -HOC ₆ H ₄ TeCl ₃ . Phen $c \rightarrow b \rightarrow b \rightarrow c \rightarrow c$ $a \rightarrow d \rightarrow c \rightarrow d$ $TeCl_3 \rightarrow H_A \rightarrow H_A \rightarrow H_B \rightarrow CH_3$ | $\begin{array}{c} 2.31^{s}(3H)\\ 6.84,6.98^{b}\;(1H)\\ 7.70{-}8.21^{cm}\;(7H)\end{array}$ | —CH ₃ proton H _B H _A , H _C , H _a , H _b , and OH |
| $RTeCl_3.Phen$ (R = 3-methyl-4-hydroxyphenyl) | 8.58 ^d (2H) 9.20 ^d (2H) | ${ m H_c} { m H_d}$ |

TABLE III PMR Spectral Data of Phenanthroline Complexes ofHydroxyaryltellurium Trichlorides in DMSO- d_6

b = broad; d = doublet; cm = complex multiplet; and s = singlet.

Synthesis of 1,10-Phenanthroline Complexes

A saturated solution of 1,10-phenanthroline monohydrate (0.99 g, 5.0 mmol) in about 20 mL dichloromethane was added to a saturated solution of 5.0 mmol (1.64 g, 2.30 g, 3.01 g for *p*-hydroxyphenyltellurium trichloride, tribromide, triiodide; 1.71 g, 2.37 g, 3.08 g for 3-methyl-4-hydroxyphenyltellurium trichloride, tribromide, and triiodide, respectively) of hydroxyaryltellurium trihalide in about 20 mL dry methanol/

| TABLE IV Physical Characteristics and Elemental Analyses of Phenanthroline Complexes of Hydroxyaryltellurium Trihalides | |
|-------------------------------------------------------------------------------------------------------------------------|--|
| Elemental analyses found (calc | |

| | Complex | | M.P. °C | Yield. | Elemen | tal anal | yses four | od (calc | 1.) % |
|-------------------------|----------------------------|----------------------|-----------|--------|---------|----------|-----------|----------|--------|
| R | (empirical formula) | Color | (dec.) | % | Te | Х | С | Η | z |
| | RTeCl ₃ .Phen | Light pink | 188 - 190 | 80 | 24.66 | 21.23 | 41.98 | 2.38 | 5.22 |
| | $(C_{18}H_{13}Cl_3N_2OTe)$ | | | | (25.14) | (20.97) | (42.62) | (2.58) | (5.52) |
| | $ m RTeBr_3.Phen$ | Light green | 178 - 180 | 80 | 19.98 | 37.11 | 33.26 | 1.81 | 4.01 |
| | $(C_{18}H_{13}Br_3N_2OTe)$ | | | | (19.92) | (37.42) | (33.75) | (2.05) | (4.37) |
| <i>p</i> -Hydroxyphenyl | RTeI ₃ . Phen | Red | 152 - 155 | 72 | 16.10 | 48.36 | 27.83 | 1.36 | 1.89 |
| | $(C_{18}H_{13}I_3N_2OT_e)$ | | | | (16.33) | (48.71) | (27.66) | (1.68) | (2.04) |
| | $RTeCl_3$. Phen | Light pink | 98 - 100 | 90 | 24.11 | 20.36 | 43.28 | 2.67 | 5.11 |
| | $(C_{19}H_{15}Cl_3N_2OTe)$ | | | | (24.48) | (20.40) | (43.78) | (2.90) | (5.37) |
| 3-Methyl-4- | $ m RTeBr_3.Phen$ | Light orange | 200 - 203 | 85 | 19.11 | 36.40 | 34.21 | 2.44 | 4.66 |
| hydroxyphenyl | $(C_{19}H_{15}Br_3N_2OTe)$ | | | | (19.49) | (36.62) | (34.86) | (2.31) | (4.28) |
| | RTeI ₃ . Phen | Red | 138 - 141 | 75 | 15.80 | 47.36 | 28.54 | 1.67 | 1.87 |
| | $(C_{19}H_{15}I_3N_2OTe)$ | | | | (16.04) | (47.85) | (28.68) | (1.90) | (2.01) |

chloroform. The contents were refluxed for about 2 h. The solvent was distilled off and the solid thus obtain was extracted with benzene repeatedly and finally with chloroform and then dried over P_4O_{10} in a vacuum dessicator. The complexes were recrystallized from a mixture of methanol and chloroform (50:50).

The purity of these compounds was checked by T.L.C. using silica gel- G. The analytical data and physical properties of the complexes are presented in Table IV.

Analytical Methods and Physical Measurements

Carbon, hydrogen, and nitrogen analyses were obtained microanalytically from sophisticated Analytical Instrumentation Facility, Panjab University, Chandigarh. The tellurium content was estimated volumetrically.²⁶ Conductivity was measured at $25 \pm 2^{\circ}$ C with a conductivity bridge type 305 Systronics model and the molecular weights were determined by the cryoscopic method in nitrobenzene up to the saturation point. IR spectra were recorded in the region 4000–400 cm⁻¹ at University of Delhi on a SHIMAZDU FTIR - 8300 Spectrometer, using the KBr pellets technique. ¹H NMR spectra were recorded at SAIF, Panjab University, Chandigarh on BRUKER AVANCE II 400 NMR Spectrometer operating at 400.13 MHz in DMSO- d₆using tetramethylsilane as an internal reference

REFERENCES

- [1] K. J. Wynne and P. S. Pearson, Inorg. Chem., 10, 2735 (1971).
- [2] K. J. Wynne and P. S. Pearson, J. Chem. Soc., Chem. Commun., 556 (1970).
- [3] K. J. Wynne, A. J. Clark, and M. Berg, J. Chem. Sec. Dalton, 2370 (1972).
- [4] E. R. Clark, A. J. Collet, and D. G. Naik, J. Chem. Soc. Dalton, 1961 (1973).
- [5] M. C. Berg, Diss. Abstr. Int. B, 33 (7), 2982 (1972).
- [6] T. N. Srivastava, M. Singh, and H. B. Singh, Indian J. Chem., 21 (A), 307 (1982).
- [7] T. N. Srivastava, R. C. Srivastava, and M. Srivastava, Indian J. Chem., 21 (A), 539 (1982).
- [8] T. N. Srivastava, R. C. Srivastava, and V. K. Srivastava, J. Indian Chem. Soc., 60, 891(1983).
- [9] M. V. Garad, Polyhedron, 4, 1353(1985).
- [10] K. K. Verma and Reena, Synth. React. Inorg. Met. -Org. Chem., 29, 499 (1999).
- [11] K. K. Verma, R. Dahiya, and D. Soni, Synth. React. Inorg. Met. -Org. Chem., 29, 1033 (1999).
- [12] K. K. Verma and R. Dahiya, Synth. React. Inorg. Met.-Org. Chem., 29, 1299 (1999).
- [13] K. K. Verma and Reena, Phosphorus, Sulfur, and Silicon and the Related Elements, 148, 227 (1999).
- [14] Reena, Ph.D. Thesis, M. D. University Rohtak, Rohtak, India (1999).
- [15] K. Kumar, Ph.D. Thesis, IIT Delhi, New Delhi, India (1981).

- [16] B. L. Khandelwal, K. Kumar, and F. J. Berry, Inorg. Chim. Acta, 47, 135 (1981).
- [17] B. L. Khandelwal, K. Kumar, and K. Raina, Synth. React. Inorg. Met. –Org. Chem., 11, 65 (1981).
- [18] W. J. Geary, Coord. Chem. Revs., 7, 81 (1971).
- [19] K. K. Verma, S. Chand, and S. Garg, Synth. React. Inorg. Met. -Org. Chem., 24, 979, (1994).
- [20] A. A. Schilt and R. C. Taylor, J. Inorg. Nucl. Chem., 9, 211 (1959).
- [21] B. Sey Ferth and N. Kahlen, J. Organometal. Chem., 25, 809 (1960).
- [22] A. A. Osowole, J. A. O. Woods, and O. A. Odunola, Synth. React. Inorg. Met. -Org. Chem., 32, 783 (2002).
- [23] Shin-ichi, Hand Book of Proton NMR Spectra and Data, Asani Research Centre, Ed. (Academic Press, Amsterdam, 1985).
- [24] A. I. Vogel, A Text Book of Practical Organic Chemistry (Longman, London, 1991), 5th ed., pp. 131–234.
- [25] A. Weissberger, Ed., Techniques of Organic Chemistry (Interscience Publishers, Inc., New York, 1967), Vol. VII, 2nd ed.
- [26] A. I. Vogel, A Text Book of Quantitative Inorganic Analysis Including Elementary Instrumental Analysis (Longman, London, 1975) 3rd ed., p. 324.