# First efficient synthesis of (±)-erythro-roccellic acid †

# Sivaprakasam Mangaleswaran and Narshinha P. Argade\*

Division of Organic Chemistry (Synthesis), National Chemical Laboratory, Pune 411 008, India. E-mail: argade@dalton.ncl.res.in; Fax: + 91-20-5893153

Received (in Cambridge, UK) 4th May 2001, Accepted 19th June 2001 First published as an Advance Article on the web 9th July 2001



An easy, four-step route to  $(\pm)$ -*erythro*-2-dodecyl-3-methylbutanedioic acid (roccellic acid, 9) with 60% overall yield has been described. Wittig reaction of the citraconimide–TPP adduct with dodecanal furnished the mixture of (E)- and (Z)-isomers 3 and 4, which, on base-catalysed exocyclic to endocyclic double bond isomerisation, yielded maleimide 5. Catalytic *cis*-hydrogenation of 5 using Adam's platinum dioxide catalyst followed by acid-catalysed hydrolysis of formed succinimide 8 gave desired  $(\pm)$ -*erythro*-roccellic acid 9.

### Introduction

(+)-Roccellic acid [(2R,3S)-2-dodecyl-3-methylbutanedioic acid, 1] occurs in lichens<sup>1,2</sup> and it was first isolated in 1898. In the past century it has been isolated from the following lichen species: Roccella capensis,<sup>3</sup> R. fuciformis,<sup>4,5</sup> R. hypomecha,<sup>6</sup> R. gayana,<sup>7</sup> R. fucides,<sup>7,8</sup> R. condensata,<sup>9</sup> R. montagnei,<sup>10,11</sup> Dirinaria aegialita,<sup>12</sup> D. applanata,<sup>12</sup> D. confusa saxicola,<sup>12</sup> D. consimilis,<sup>12</sup> D. leopoldii,<sup>12</sup> Pyxine berteriana,<sup>12</sup> P. caesiopruinosa,<sup>12</sup> P. pungens,<sup>12</sup> Lobodirina cerebriformes,<sup>13</sup> L. mahuiana,<sup>14</sup> Acarospora chlorophana,<sup>15</sup> Lecanora riparia,<sup>16</sup> L. rupicola,<sup>17,18</sup> L. sordida,<sup>19,20</sup> Lepraria latebrarum,<sup>21</sup> L. aeruginosa,<sup>22</sup> Dirina lutosa,<sup>23</sup> Crocynea membranacea<sup>24,25</sup> and more recently from Haematomma nemetzii<sup>26</sup> and Tornabena scutellifera<sup>26</sup> with a major contribution from Siegfried Huneck's group. The structural assignment of roccellic acid 1 has been made on the basis of analytical and spectral data.<sup>17,19,24,26</sup> Its absolute configuration was established by Åkermark by degrading 1 to its two isomeric monomethyl esters.<sup>24,27</sup> Roccellic acid **1** possesses antituberculosis activity.<sup>28-30</sup> and concentration-dependent plant growth promoter  ${}^{31-33}$ /inhibitor  ${}^{33,34}$  activity. It is also used for (i) the synthesis of structural analogues of the antibiotic actinonin,35 (ii) the precipitation of human serum albumin<sup>36</sup> and (iii) the preparation of coloured metal complexes.<sup>37</sup> To date, two syntheses  $^{38,39}$  of unnatural (±)-*threo*roccellic acid (major) are known starting from diethyl malonate and during these studies a small amount (25 mg) of  $(\pm)$ -erythroroccellic acid has been obtained by an 8-step synthesis with 0.026% overall yield.<sup>38</sup> The provision of facile synthetic approaches to this bioactive natural product roccellic acid is a challenging task of current interest. Recently we have reported 40 the first coupling reaction of the citraconimide 2 and triphenylphosphine (TPP) adduct<sup>41</sup> with aliphatic aldehydes and used it for the synthesis of the recently isolated bioactive natural products chaetomellic acid A,<sup>40</sup> (±)-piliformic acid,<sup>42</sup> and tyromycin A.<sup>43</sup> We herein report yet another useful application of this coupling reaction to complete the first efficient synthesis of  $(\pm)$ -erythro-roccellic acid 9 by the cis-reduction of maleimide 5 to succinimide 8 employing a catalytic hydrogenation reaction with Adam's catalyst (Scheme 1).



<sup>†</sup> NCL Communication No. 6609.

1764 J. Chem. Soc., Perkin Trans. 1, 2001, 1764–1766



Scheme 1 Reagents, conditions and yields: (i) TPP, AcOH, dodecanal, reflux, 10 h (82%); (ii) TEA, THF, reflux, 48 h (98%); (iii) (a) KOH, THF, MeOH, H<sub>2</sub>O, reflux, 2 h; (b) H<sup>+</sup>/HCl (98%); (iv) Adam's catalyst, petroleum spirit, H<sub>2</sub>, rt, 10 h (9: 60%; 8: 95%); (v) CF<sub>3</sub>COOH, conc. HCl, reflux, 48 h (98%); (vi) CH<sub>2</sub>N<sub>2</sub>, Et<sub>2</sub>O, rt, 2 h, (98%).

#### **Results and discussion**

A mixture of equimolar amounts of citraconimide **2** and TPP on refluxing with 1.5 equivalents of dodecanal in glacial acetic acid for 10 h yielded a combination of geometric isomers **3** and

DOI: 10.1039/b104105n

4 in 82% yield by a Wittig reaction. The integration values for the vinylic protons in the <sup>1</sup>H NMR spectra of this mixture of geometric isomers (E)-3 and (Z)-4 revealed that they are formed in an 85:15 ratio. The mixture of exo-isomers 3 and 4 on refluxing with triethylamine-THF (1:1) for 48 h, underwent a smooth exocyclic to endocyclic carbon-carbon double bond isomerisation to yield dodecylmethylmaleimide 5 in 98% yield. On alkaline hydrolysis followed by acidification, the maleimide 5 gave the desired dodecylmethylmaleic anhydride 6 in 98% yield. We systematically studied the catalytic cis-hydrogenation reactions of maleimide 5 and disubstituted maleic anhydride 6 to obtain succinimide derivative 8 and succinic anhydride derivative 7 respectively. In our hands, the anhydride 6 and imide 5, when subjected to catalytic hydrogenation with palladium on charcoal as catalyst in methanol, remained completely unreacted for 12 h at 50 psi pressure of hydrogen. The anhydride 6 in petroleum spirit on treatment with Adam's platinum dioxide catalyst at 50 psi pressure of hydrogen for 10 h followed by filtration, concentration and recrystallisation with acetone gave the desired  $(\pm)$ -erythro-roccellic acid 9 in 60% yield. We were unable to isolate the intermediate anhydride 7 under the present set of reaction conditions. The maleimide 5 underwent a very smooth catalytic cis-hydrogenation at rt in petroleum spirit with Adam's platinum dioxide catalyst at 50 psi hydrogen pressure in 10 h to yield the desired cissuccinimide derivative 8 in 95% yield. The reaction of maleimide 5 to succinimide 8 was much cleaner and more efficient compared to the conversion of 6 to 9 via 7. The cis-succinimide derivative 8 on hydrolysis using refluxing mixture of acetic acid plus conc. hydrochloric acid (1:1) gave a mixture of erythroand threo-isomers of roccellic acid, in a 2 : 1 ratio (by <sup>1</sup>H NMR, from the integration values of the methine protons) in 48 h with 98% yield. The cis-succinimide derivative 8 on hydrolysis in refluxing mixture of trifluoroacetic acid and conc. hydrochloric acid (1:1) gave a mixture of the desired erythro- and undesired threo-isomers with a much better proportion of the erythroisomer (93 : 7, by <sup>1</sup>H NMR, from the integration values of the methine protons) in 48 h with 98% yield. Recrystallisation of the above mixture of erythro- and threo-isomers (93:7) from acetone gave the desired  $(\pm)$ -erythro-roccellic acid 9 in 80% yield. The formed acid 9 was further characterised as its dimethyl ester 10. The analytical and spectral data obtained for (±)-ervthro-roccellic acid 9 and its corresponding dimethyl ester 10 were in complete agreement with reported data.<sup>2,17,38</sup> The asymmetric catalytic hydrogenation<sup>44-46</sup> of maleimide 5 will provide an elegant route to the desired naturally occurring bioactive (+)-(2R,3S)-roccellic acid 1.

In summary, we have demonstrated the first efficient fourstep synthesis of  $(\pm)$ -*erythro*-roccellic acid **9** with 60% overall yield, starting from citraconimide **2** by a catalytic *cis*hydrogenation pathway and the present strategy will be useful for obtaining several 2,3-dialkyl substituted succinic acid derivatives.

## Experimental

Melting points were taken on a Büchi melting point B-540 apparatus and are uncorrected. The <sup>1</sup>H NMR spectra were recorded in CDCl<sub>3</sub> with TMS as an internal standard on a Bruker AC 200 NMR spectrometer (200 MHz) and for roccellic acid **9** in pyridine- $d_5$  with TMS as an internal standard on a Bruker DRX 500 NMR spectrometer (500 MHz). The <sup>13</sup>C NMR spectra were recorded on Bruker AC 200 (50 MHz), Bruker MSL 300 (75 MHz) and Bruker DRX 500 NMR spectrometers (125 MHz). Mass spectra were recorded on a Finnigan Mat 1020 °C mass spectrometer at 70 eV. The FT-IR spectra were recorded on an FT-IR-8300 Shimadzu spectrometer and microanalyses were carried out on a Carlo-Erba instrument; column chromatographic separation was performed with ACME silica gel (60–120 mesh). Triphenylphos-

phine and dodecanal were obtained from Aldrich Chemical Co. Petroleum spirit refers to the fraction with distillation range 60-80 °C.

# (±)-(E/Z)-2-Dodecylidene-3-methyl-N-(p-tolyl)succinimides 3 and 4

A mixture of citraconimide 2 (3.01 g, 15 mmol), TPP (3.93 g, 15 mmol) and dodecanal (4.14 g, 22.5 mmol) in glacial acetic acid (50 mL) was refluxed with stirring for 10 h. Acetic acid was distilled off in vacuo at 50 °C and the residue was dissolved in ethyl acetate (60 mL). The organic layer was washed successively with water (30 mL), brine (20 mL) and dried over Na<sub>2</sub>SO<sub>4</sub>. Concentration of the organic layer in vacuo followed by silica gel column purification of the residue using petroleum spirit-ethyl acetate (9:1) gave a mixture of 3 and 4 (3:4)85:15 by <sup>1</sup>H NMR) (4.54 g, 82%); mp 49–52 °C; IR (Nujol)  $v_{\text{max}}$  1717, 1458 cm<sup>-1</sup>; <sup>1</sup>H NMR  $\delta_{\text{H}}$  0.88 (t, *J* 6 Hz, 3 H), 1.27 (br s, 16 H), 1.45–1.60 (m, 2 H), 1.48 (d, J 6 Hz, 0.45 H, Z-isomer), 1.52 (d, J 6 Hz, 2.55 H), 2.20-2.36 (m, 1.7 H), 2.38 (s, 3 H), 2.74-2.92 (m, 0.3 H, Z-isomer), 3.34-3.59 (m, 1 H), 6.23 (dt, J 8 and 3 Hz, 0.15 H, Z-isomer), 6.92 (dt, J 8 and 3 Hz, 0.85 H), 7.20 (d, J 8 Hz, 2 H), 7.28 (d, J 8 Hz, 2 H); MS m/z 369 (M<sup>+</sup>) 85%), 354 (4), 340 (4), 270 (5), 256 (8), 242 (100), 229 (28), 216 (47), 203 (80), 186 (8), 170 (4), 157 (4), 144 (4), 132 (12), 118 (18), 107 (46), 95 (62), 91 (29), 81 (55), 68 (82), 55 (9) (Calc. for C24H35NO2: C, 78.00; H, 9.55; N, 3.80. Found: C, 78.16; H, 9.39; N, 3.92%).

#### 2-Dodecyl-3-methyl-N-(p-tolyl)maleimide 5

A solution of the mixture of **3** and **4** (3.3 g) in THF (15 mL) and triethylamine (15 mL) was refluxed for 48 h with stirring. The reaction mixture was then allowed to reach rt, and was concentrated *in vacuo*; silica gel column chromatographic purification of the residue using petroleum spirit–ethyl acetate (9 : 1) gave pure **5** (3.23 g, 98%), mp 68–69 °C; IR (Nujol)  $v_{max}$  1710, 1690, 1650 cm<sup>-1</sup>; <sup>1</sup>H NMR  $\delta_{\rm H}$  0.89 (t, *J* 8 Hz, 3 H), 1.27 (br s, 18 H), 1.50–1.75 (m, 2 H), 2.05 (s, 3 H), 2.38 (s, 3 H), 2.46 (t, *J* 8 Hz, 2 H), 7.10–7.40 (m, 4 H); <sup>13</sup>C NMR (CDCl<sub>3</sub>, 50 MHz)  $\delta_{\rm C}$  8.7, 14.0, 21.0, 22.6, 23.8, 28.1, 29.3, 29.6 (6 × CH<sub>2</sub>), 31.9, 125.6, 129.5, 137.1, 137.2, 141.4, 159.7, 170.7, 171.1; MS *m*/*z* 369 (M<sup>+</sup>, 85%), 256 (6), 228 (10), 215 (100), 203 (8), 107 (48), 91 (29), 81 (41), 67 (20), 55 (10) (Calc. for C<sub>24</sub>H<sub>35</sub>NO<sub>2</sub>: C, 78.00; H, 9.55; N, 3.80. Found: C, 77.72; H, 9.78; N, 3.71%).

#### 2-Dodecyl-3-methylmaleic anhydride 6

To the solution of imide 5 (1.11 g, 3 mmol) in a THF-methanol mixture (1:2, 18 mL) was added 30% aqueous KOH solution (10 mL) and the reaction mixture was refluxed for 2 h with stirring. The reaction mixture was concentrated in vacuo and the residue was acidified with dilute HCl, extracted with diethyl ether  $(3 \times 50 \text{ mL})$  and the organic layer was washed with water (20 mL), brine (20 mL) and dried over Na<sub>2</sub>SO<sub>4</sub>. Concentration of the organic layer in vacuo followed by silica gel column chromatographic purification of the residue using petroleum spirit-ethyl acetate (9:1) furnished pure 6 as a thick oil (0.82 g, 98%); IR (Nujol) v<sub>max</sub> 1823, 1767, 1674, 1466 cm<sup>-1</sup>; <sup>1</sup>H NMR δ<sub>H</sub> 0.88 (t, J 6 Hz, 3 H), 1.26 (br s, 18 H), 1.45–1.70 (m, 2 H), 2.07 (s, 3 H), 2.46 (t, J 6 Hz, 2 H); <sup>13</sup>C NMR (CDCl<sub>3</sub>, 50 MHz)  $\delta_{\rm C}$  9.2, 13.9, 22.5, 24.3, 27.4, 29.3, 29.5 (6 × CH<sub>2</sub>), 31.8, 140.3, 144.6, 165.7, 166.1; MS m/z 281 (MH<sup>+</sup>, 10%), 262 (18), 252 (12), 235 (13), 207 (20), 196 (14), 178 (42), 168 (29), 150 (15), 139 (10), 126 (95), 109 (12), 98 (26), 81 (33), 67 (30) (Calc. for C<sub>17</sub>H<sub>28</sub>O<sub>3</sub>: C, 72.82; H, 10.06. Found: C, 73.05; H, 9.93%).

#### (±)-erythro-2-Dodecyl-3-methyl-N-(p-tolyl)succinimide 8

A mixture of maleimide 5 (1.1 g, 3 mmol) and Adam's catalyst (25 mg) in petroleum spirit (50 mL) was subjected to hydrogenation at 50 psi hydrogen pressure for 10 h at rt. The reaction

mixture was filtered through Celite and the filtrate was concentrated in vacuo. The silica gel column purification of the residue using petroleum spirit-ethyl acetate (9:1) gave succinimide 8 (1.05 g, 95%), mp 67–68 °C (from methanol); IR (Nujol)  $v_{max}$ 1774, 1709, 1516, 1470 cm<sup>-1</sup>; <sup>1</sup>H NMR (CDCl<sub>3</sub>, 200 MHz)  $\delta_{\rm H}$  0.91 (t, J 6 Hz, 3 H), 1.29 (br s, 20 H), 1.35 (d, J 6 Hz, 3 H), 1.60-1.95 (m, 2 H), 2.39 (s, 3 H), 2.85-3.20 (m, 2 H), 7.16 (d, J 8 Hz, 2 H), 7.28 (d, J 8 Hz, 2 H); <sup>13</sup>C NMR (CDCl<sub>3</sub>, 50 MHz)  $\delta_{\rm C}$  11.6, 13.9, 21.0, 22.6, 26.5, 27.6, 29.3, 29.5 (6 × CH<sub>2</sub>), 31.8, 38.6, 43.8, 126.1, 129.6, 138.2, 138.4, 178.4, 179.3; MS m/z 371 (M<sup>+</sup>, 24%), 265 (4), 216 (29), 203 (80), 175 (7), 134 (12), 107 (20), 91 (12), 69 (18) (Calc. for C<sub>24</sub>H<sub>37</sub>NO<sub>2</sub>: C, 77.58; H, 10.03; N, 3.77. Found: C, 77.76; H, 10.14; N, 3.53%). Similarly, on catalytic hydrogenation, anhydride 6 (0.56 g, 2 mmol) furnished (±)-*erythro*-roccellic acid **9** (0.36 g, 60%).

#### (±)-erythro-Roccellic acid 9

Succinimide 8 (0.74 g, 2 mmol) was dissolved in trifluoroacetic acid-conc. hydrochloric acid (1:1, 15 mL) and the reaction mixture was refluxed for 48 h. The reaction mixture was kept aside at rt for 12 h and the white precipitate of roccellic acid was filtered in vacuo to obtain 9 (0.59 g, 98%) (erythro: threo = 93:7, proved by <sup>1</sup>H NMR). The mixture of erythro- and threoroccellic acid (0.50 g, 93:7) was recrystallised from acetone (30 mL) to obtain pure (±)-erythro-roccellic acid 9 (0.40 g, 80%), mp 141–142 °C (from acetone); IR (Nujol)  $v_{\text{max}}$  1693, 1464, 1271, 1202 cm<sup>-1</sup>; <sup>1</sup>H NMR (pyridine- $d_5$ , 500 MHz)  $\delta_{\text{H}}$ 0.86 (t, J 5 Hz, 3 H), 1.20 (br s, 16 H), 1.30-1.45 (m, 2 H), 1.62 (d, J 10 Hz, 3 H), 1.53-1.75 (m, 2 H), 1.92-2.03 (m, 1 H), 2.10-2.23 (m, 1 H), 3.17-3.33 (m, 2 H); <sup>13</sup>C NMR (pyridine-d<sub>5</sub>, 125 MHz)  $\delta_{\rm C}$  14.2, 16.2, 22.8, 28.2, 29.5, 29.8 (6 × CH<sub>2</sub>), 31.7, 32.0, 43.4, 50.0, 177.0, 177.7; MS m/z 282 (M<sup>+</sup>, 1%), 254 (2), 227 (7), 209 (3), 170 (10), 156 (17), 132 (35), 97 (33), 83 (28), 69 (54) (Calc. for C<sub>17</sub>H<sub>32</sub>O<sub>4</sub>: C, 67.96; H, 10.73. Found: C, 67.85; H, 10.65%).

#### Dimethyl ester of (±)-erythro-roccellic acid 10

A solution of  $(\pm)$ -erythro-roccellic acid 9 (0.30 g, 1 mmol) in ether (10 mL) was treated with a solution of diazomethane in ether at 0 °C, until the complete consumption of the starting material (2 h). The excess of diazomethane was quenched with acetic acid and the reaction mixture was concentrated in vacuo. Silica gel column chromatographic purification of the residue using petroleum spirit and ethyl acetate mixture (9:1) gave pure 10 as a thick oil (0.32 g, 98%); IR (neat) v<sub>max</sub> 1740, 1464, 1435, 1195 cm<sup>-1</sup>; <sup>1</sup>H NMR (CDCl<sub>3</sub>, 200 MHz)  $\delta_{\rm H}$  0.87 (t, J 6 Hz, 3 H), 1.13 (d, J 6 Hz, 3 H), 1.23 (br s, 18 H), 1.30-1.50 (m, 2 H), 1.50–1.75 (m, 2 H), 2.58–2.78 (m, 2 H), 3.68 (s, 3 H), 3.69 (s, 3 H); <sup>13</sup>C NMR (CDCl<sub>3</sub>, 75 MHz)  $\delta_{\rm C}$  14.0, 15.1, 22.7, 27.5, 29.4, 29.6 (6 × CH<sub>2</sub>), 30.7, 31.9, 42.1, 48.6, 51.4, 51.7, 174.6, 175.3; MS m/z 328 (M<sup>+</sup>, 1%), 298 (24), 267 (7), 242 (100), 184 (16), 170 (18), 160 (88), 128 (66), 111 (17), 101 (32), 91 (68), 81 (29), 69 (68), 55 (92) (Calc. for C<sub>19</sub>H<sub>36</sub>O<sub>4</sub>: C, 69.47; H, 11.04. Found: C, 69.54; H, 11.23%).

# Acknowledgements

S. M. thanks C.S.I.R., New Delhi, for the award of a fellowship. We thank Dr K. N. Ganesh, Head of the Division of Organic Chemistry (Synthesis), for constant encouragement.

#### References

1 The Merck Index, 12th Edition, 1996, p. 1419.

- 2 S. Huneck, J. Schmidt and A. Porzel, Z. Naturforsch., B: Chem. Sci., 1994, 49, 561.
- 3 S. Huneck, J. Jakupovic and G. Follmann, Z. Naturforsch., B: Chem. *Sci.*, 1991, **46**, 969. 4 S. Huneck, *Phytochemistry*, 1972, **11**, 1489.
- 5 S. Huneck, A. Mathey and G. Trotet, Z. Naturforsch., B: Anorg. Chem. Org. Chem., 1967, 22, 1367.
- 6 S. Huneck and G. Follmann, Ber. Dtsch. Bot. Ges., 1968, 81, 125 (Chem. Abstr., 1969, 70, 35025j).
- 7 S. Huneck and G. Follmann, Z. Naturforsch., B: Anorg. Chem. Org. Chem., 1967, 22, 1369.
- 8 S. Huneck, G. Follmann, A. W. William and G. Trotet, Z. Naturforsch., B: Anorg. Chem. Org. Chem., 1967, 22, 671.
- 9 S. Huneck and G. Follmann, Z. Naturforsch., B: Anorg. Chem. Org. Chem., 1967, 22, 1185.
- 10 V. Subba Rao and T. R. Seshadri, Proc. Indian Acad. Sci., Sect. A, 1940, 12, 466.
- 11 V. Subba Rao and T. R. Seshadri, Proc. Indian Acad. Sci., Sect. A, 1941, 13, 199.
- 12 S. Huneck, M. A. Morales and K. Kalb, J. Hattori Bot. Lab., 1987, 62, 331 (Chem. Abstr., 1988, 108, 19191h).
- 13 W. Quilhot, J. A. Garbarino and V. Gambaro, J. Nat. Prod., 1983, 46. 593
- 14 W. Quilhot, J. Redon, E. Zuniga and S. Vidal, Phytochemistry, 1975, 14. 1865.
- 15 S. Huneck, Lichenologist, 1980, 12, 239.
- 16 S. Huneck, Phytochemistry, 1972, 11, 1493.
- 17 J. P. Devlin, C. P. Falshaw, W. D. Ollis and R. E. Wheeler, J. Chem. Soc. (C), 1971, 1318.
- 18 C. H. Fox and S. Huneck, Phytochemistry, 1969, 8, 130.
- 19 G. Kennedy, J. Breen, J. Keane and T. J. Nolan, Sci. Proc. R. Dublin Soc., 1937. 21, 557.
- 20 O. Hesse, J. Prakt. Chem., 1898, 58, 497.
- 21 K. Soviar, Acta Fac. Pharm. Univ. Comenianae, 1971, 20, 27 (Chem. Abstr., 1972, 76, 123969z).
- 22 K. Soviar, M. Bachrata, D. Georch and L. Krasnec, Farm. Obz., 1967, 36, 161 (Chem. Abstr., 1969, 70, 44812r).
- 23 S. Huneck and G. Follmann, Z. Naturforsch., B: Anorg. Chem. Org. Chem., 1964, 19, 658.
- 24 B. Åkermark, Acta Chem. Scand., 1962, 16, 599.
- 25 B. Åkermark, H. Erdtman and C. A. Wachtmeister, Acta Chem. Scand., 1959, 13, 1855.
- 26 S. Huneck, U. Himmelreich, J. Schmidt, J. Volker and U. Zeybek, Z. Naturforsch., B: Chem. Sci., 1994, 49, 1561.
- 27 B. Åkermark, Acta Chem. Scand., 1970, 24, 1456.
- V. C. Barry and P. A. McNally, *Nature*, 1945, **156**, 48.
  V. C. Barry, J. G. Belton, R. M. Kelly and D. Twomey, *Nature*, 1950, 166, 303.
- 30 F. M. Pereira, J. de sa and S. S. Bhatnagar, Indian J. Pharm., 1953, 15, 287
- 31 C. F. Garcia, G. A. Espinoza, S. G. Coltantes, V. V. Rios and P. W. Quilhot, J. Hattori Bot. Lab., 1982, 53, 443 (Chem. Abstr., 1982, 97, 212854i).
- 32 P. W. Quilhot, V. J. Thompson, B. S. Vidal and P. G. Campos, J. Hattori Bot. Lab., 1981, 49, 273 (Chem. Abstr., 1981, 95, 1773c).
- 33 S. Huneck and K. Schreiber, Phytochemistry, 1972, 11, 2429.
- 34 K. Schreiber, Environ. Qual. Saf. Suppl., 1975, 3, 483 (Chem. Abstr., 1976, 85, 105260t).
- 35 J. P. Devlin, W. D. Ollis and J. E. Thorpe, J. Chem. Soc., Perkin Trans. 1, 1975, 846.
- 36 W. McD. Armstrong, Proc. R. Ir. Acad., Sect. B, 1956, 58, 71.
- 37 I. K. Iskandar and J. K. Syers, J. Soil Sci., 1972, 23, 255.
- 38 B. Åkermark and N. G. Johansson, Ark. Kemi, 1967, 27, 1.
- 39 V. C. Barry and D. Twomey, Proc. R. Ir. Acad., Sect. B, 1947, 51, 137
- 40 S. B. Desai and N. P. Argade, J. Org. Chem., 1997, 62, 4862.
- 41 E. Hedaya and S. Theodoropulos, Tetrahedron, 1968, 24, 2241.
- 42 S. Mangaleswaran and N. P. Argade, J. Chem. Soc., Perkin Trans. 1, 2000, 3290.
- 43 S. Mangaleswaran and N. P. Argade, J. Org. Chem., 2001, 66, 5259.
- 44 H. Muramatsu, H. Kawano, Y. Ishii, M. Saburi and Y. Uchida, J. Chem. Soc., Chem. Commun., 1989, 769.
- 45 M. Kitamura, M. Tokunaga and R. Noyori, J. Org. Chem., 1992, 57, 4053
- 46 M. V. Berg, A. J. Minnaard, E. P. Schudde, J. V. Esch, A. H. M. Vries, J. G. Varies and B. L. Feringa, J. Am. Chem. Soc., 2000, 122, 11539.