

PROTOLYSES OF $(\text{CH}_3)_3\text{SnM}(\text{CH}_3)_3$ ($\text{M} = \text{Sn}, \text{Ge}, \text{Si}, \text{C}$)

MATTHEW J. CUTHBERTSON, DARRYL W. HAWKER and PETER R. WELLS *

Department of Chemistry, University of Queensland, Brisbane 4067 (Australia)

(Received November 8th, 1984)

Summary

Product and kinetic studies on the reactions of hydrogen chloride in methanol solution with the substrates $(\text{CH}_3)_3\text{SnM}(\text{CH}_3)_3$ ($\text{M} = \text{Sn}; \text{Ge}$ and Si) show that both $\text{Sn}-\text{M}$ and $\text{Sn}-\text{CH}_3$ cleavage reactions occur, at similar rates, and are followed by other reactions giving complex but explicable mixtures of products. Similar behaviour is observed for trifluoroacetolysis in carbon tetrachloride solution, and some intermediates are observable. Trifluoroacetolysis of $(\text{CH}_3)_3\text{SnC}(\text{CH}_3)_3$ results in exclusive $\text{Sn}-\text{CH}_3$ cleavage. The very slow apparent solvolysis in acetic acid solution is thought to involve reaction with dissolved oxygen.

Introduction

chlorotrimethylsilane, chlorotrimethylstannane (by ^1H NMR), and an unidentified white precipitate of m.p. 108°C . The latter corresponds to dichlorodimethylstannane, m.p. 107°C , which has a low solubility in hydrocarbon solvents. It was also reported that $(\text{CH}_3)_3\text{SiGe}(\text{CH}_3)_3$ shows no indications of any reaction with hydrogen chloride at 70°C after 14 h. We have found similarly, and in common with many other workers, that hexamethyldigermane and hexamethyldisilane are quite stable towards acid.

The reaction of hexamethyldistannane with hydrogen chloride in methanol solution has been studied by Tagliavini, Belluco and Pilloni [8] who used solvolysis of acetyl chloride to generate the acid *in situ*. They report that the primary reaction gives hydrogen and chlorotrimethylstannane only, but that this reaction is accompanied by the chlorotrimethylstannane-catalysed decomposition of hexamethyldistannane to tetramethylstannane and yellow polymeric "dimethyltin". There is almost complete suppression of this second reaction when three equivalents of acetyl chloride are used.

Reactions of hexamethyldistannane with halogenated acetic acids, but not with acetic acid itself, have been reported by Birchall and Johnson [9] to give good yields of the $(\text{CH}_3)_4\text{Sn}_2\text{X}_2$ species, including the trifluoroacetoxy derivative.

Experimental

Materials

Acetolysis reactions

Acetic acid, then the cyclohexane reference, were added to a pre-weighed NMR tube which was re-weighed at each stage then the substrate was added to produce a ca. 0.2 *M* solution. Reactions were so slow that the first ^1H NMR spectrum showed no reaction products. Reaction times of several days were required before significant product formation was observed.

Reactions with trifluoroacetic acid in carbon tetrachloride solution

Small aliquots (ca. 0.2 equivalents) of $\text{CF}_3\text{CO}_2\text{H}$ were added to a solution of the substrate of known concentration (ca. 0.2 *M*) in CCl_4 containing cyclohexane as a reference, contained in an NMR tube, and the ^1H NMR spectra were recorded. Reactions were rapid and complete before the spectra were obtained. This procedure was repeated until most of the substrate had been consumed. (In the case of trimethylgermyltrimethylstannane the reaction was terminated at ca. 50% when product began to precipitate out).

Analysis of reaction mixtures

The composition of the reaction mixture at any time was determined from corrected peak height measurements of the various identified components, as previously described [5]. For kinetic studies the progress of the reaction was best followed by observation of the substrate resonances either as two separate measures or as their average. In the competition experiments, the complexity




Fig. 1. ^1H NMR spectrum of reaction mixture: $(\text{CH}_3)_3\text{SnSi}(\text{CH}_3)_3$ (0.19 M) + HCl (0.50 M) after 1 day.

TABLE 1

ASSIGNMENT OF RESONANCES IN FIG. 1, 2 AND 3 ^a

Peak	Chemical shift (ppm)	$J(^{119}\text{Sn}-\text{H})$ (Hz)	Identity
A	1.45		cyclohexane
B	1.13	93	$(\text{CH}_3)_2\text{SnCl}_2$
C	0.60	68	$(\text{CH}_3)_3\text{SnCl}$
D	0		

Allowing for an overall stoichiometry of 2.5 with respect to HCl, i.e. half-way between the values of 2 and 3 for paths A and B, respectively, an overall rate constant may be estimated as ca. $3 \times 10^{-5} M^{-1} s^{-1}$. Two kinetic runs were studied at stoichiometric ratios close to this value and evaluated by plots of $([S] + \delta)^{-1}$ vs. time, where $2\delta = \frac{1}{2.5}[HCl]_0 - [S]_0$. Several estimates of the path A/path B ratio were also obtained during these studies. The following results were obtained:

$[S]_0$	$[HCl]_0$	Second order rate constants *	Path A/Path B
0.232 M	0.604 M	$\begin{cases} 2.0_5 \times 10^{-5} M^{-1} s^{-1} \text{ (a)} \\ 1.8_5 \times 10^{-5} M^{-1} s^{-1} \text{ (b)} \end{cases}$	57/43
0.266 M	0.639 M	$\begin{cases} 1.6 \times 10^{-5} M^{-1} s^{-1} \text{ (a)} \\ 1.4_5 \times 10^{-5} M^{-1} s^{-1} \text{ (b)} \end{cases}$	60/40

Reaction of HCl with $(CH_3)_3GeSn(CH_3)_3$




Fig. 2. ^1H NMR spectrum of reaction mixture: $(\text{CH}_3)_3\text{SnGe}(\text{CH}_3)_3$ (0.24 M) + HCl (0.29 M) after 50 h.

progressive analysis of this system is illustrated in Table 2, which shows that the concentration changes not employed in the analysis are correctly predicted, with the exception of the consumption of HCl . However hydrolysis of chlorotrimethylger-

TABLE 2

ANALYSIS OF REACTION PRODUCTS FROM $(\text{CH}_3)_3\text{SnGe}(\text{CH}_3)_3 + \text{HCl}$

	t 0	t 50 h	Change observed	a ^a	b	c</
--	-------	----------	--------------------	----------------	---	-----

mane to the extent of some 20% would provide the required additional reagent.

If trimethylgermane survives completely in this system, then the yield of this product compared with the total yield of Ge-products via paths A and A' indicates that path A is responsible for ca. 35% of the Sn-Ge cleavage process. This is, of course, a minimum value.

Kinetic studies were carried out with a large excess of HCl over substrate (S) and the data evaluated by plots of $\ln[S]$ vs. time with $[S]$ determined as the average from observation of the Sn-CH₃ and Ge-CH₃ signals to give the following results:

$[S]_0$	$[HCl]_0$	second order rate constant	Path A/Path B
0.054 <i>M</i>	1.52 <i>M</i>	$9.35 \times 10^{-6} M^{-1} s^{-1}$	34.5/65.5
0.061 <i>M</i>	1.52 <i>M</i>	$9.23 \times 10^{-6} M^{-1} s^{-1}$	34.5/65.6

Under the conditions of these kinetic studies only very small amounts of trimethylgermane are observed, presumably due

and an equivalent amount of dichlorodimethylstannane. This yields measures of the extent of reaction proceeding by path B and hence estimates of path A. As Table 3 shows, there is good agreement between the chlorotrimethylstannane calculated to be so produced and twice the hexamethyldistannane estimated to be consumed by this path.

From the extent of reaction at 30 and 60 min approximate first order rate constants can be calculated. Assuming an essentially constant $[HCl] \sim 0.58 M$ these give a second order rate constant of ca. $8 \times 10^{-5} M^{-1}s^{-1}$. The analyses for 24 and 48 h indicate ratios for path A/path B of 79/21 and 63/37, respectively. But kinetic studies under these conditions are difficult since the rate constants for $(CH_3)_6Sn_2 / (CH_3)_2SnCl_2$ and $/(CH_3)_3SnCl$ are 4.8×10^{-4} and $1.0 \times 10^{-4} M^{-1}s^{-1}$ respectively [5].

A kinetic run with $[S]_0 0.51 M$ and $[HCl]_0 1.52 M$ in ca. 12% aqueous methanol gave a second order rate constant of $9.0_5 \times 10^{-5} M^{-1}s^{-1}$ and the average of several estimates of the extent of reaction by path B was $18(\pm 1)\%$, (i.e. a ratio of 82/18). Competition between $(CH_3)_6Sn_2$ (III) and $(CH_3)_3SiSn(CH_3)_3$ (I) in ca. 1




Fig. 3. ^1H NMR spectrum of reaction mixture: $(\text{CH}_3)_6\text{Sn}_2$ (0.28 M) + $(\text{CH}_3)_3\text{SnCl}$ (0.56 M) after 70 h (a) before (b) 10 min after addition of HCl.

The quantity of dichlorodimethylstannane produced corresponds almost exactly to material present in the system that would have arisen in the preceding reaction from dimethylstannylene and which would ultimately have appeared as precipitated "dimethyltin polymer".

During 10 min there is a small extent of reaction involving conversion of hexamethyldistannane into trimethylchlorostannane, presumably due to the action of the acid, but without detectable production of tetramethylstannane.

Acetolysis reactions

In all cases acetolysis is extremely slow and proceeds at essentially the same rate for $(\text{CH}_3)_6\text{Sn}_2$ (ca. 34% reaction in 21 days), $(\text{CH}_3)_3\text{SnGe}(\text{CH}_3)_3$ (ca. 29% reaction in 20 days), and $(\text{CH}_3)_3\text{SnSi}(\text{CH}_3)_3$ (ca. 39% reaction in 21 days), at which stage products, i.e. diacetoxymethylstannane and tetramethylstannane, begin to appear arising from reactions of the substrates with acetoxymethylstann

TABLE 4

¹H NMR SPECTRAL DATA FOR ACETOLYSES AND TRIFLUOROACETOLYSES ^a

Species	Acetic acid solutions ^b	CCl ₄ solutions ^c
F (CH ₃) ₄ Sn	0.07 (53.5)	0.05 (52)
E (CH ₃) ₆ Sn ₂	0.22 (47.5 and 16)	0.20 (48 and 16)
(CH ₃) ₃ Sn C(CH ₃) ₃		0.01 (48)
		1.06 (64)
I (CH ₃) ₃ Sn	0.06 (45)	0.05 (45)
H Si(CH ₃) ₃	0.32 (28)	0.35 (27)
S (CH ₃) ₃ Sn	0.12 (48.5)	0.11 (47)
R Ge(CH ₃) ₃	0.32 (26)	0.31 (24.5)
C (CH ₃) ₃ SnX	0.53 ₃ (62)	0.69 (60)
B (CH ₃) ₂ SnX ₂	0.95 (?)	1.15 (?)


Fig. 4. ^1H NMR spectrum of $(\text{CH}_3)_6\text{Sn}_2 + \text{CF}_3\text{CO}_2\text{H}/\text{CCl}_4$ system No. 3.

TABLE 5

ANALYSES OF PRODUCTS OF SUCCESSIVE $\text{CF}_3\text{CO}_2\text{H}$ ADDITIONS (IN CCl_4)

	1	2	3	4	5
(a) $(\text{CH}_3)_6\text{Sn}_2 = \text{S}$					
$([\text{S}]_0 - [\text{S}])/[\text{S}]_0$	0.14	0.39	0.66	0.90	
$[(\text{CH}_3)_3\text{SnX}]/[\text{S}]_0$	0.17	0.31	0.33	0.53	
$[(\text{CH}_3$					




Fig. 5. ^1H NMR spectrum of $(\text{CH}_3)_3\text{SnGe}(\text{CH}_3)_3 + \text{CF}_3\text{CO}_2\text{H}/\text{CCl}_4$ system No. 5.




Fig. 6. ^1H NMR spectrum of $(\text{CH}_3)_3\text{SnSi}(\text{CH}_3)_3 + \text{CF}_3\text{CO}_2\text{H}/\text{CCl}_4$ system No. 3.

obtained, and that in the time between the additions of trifluoroacetic acid decompositions via dimethylstannylene and reactions between the substrate and $(\text{CH}_3)_3\text{SnOCOCF}_3$ may have occurred. In making allowances for these reactions, so that the extent of reaction by paths A and B may be estimated it is presumed that the only source of $(\text{CH}_3)_4\text{Sn}_2(\text{OCOCF}_3)_2$ when $(\text{CH}_3)_2\text{Sn}(\text{OCOCF}_3)_2$ is absent is the trifluoroacetolysis of $(\text{CH}_3)_5\text{Sn}_2\text{OCOCF}_3$, and that it arises otherwise solely from dimethylstannylene or its equivalent.

It is noteworthy that $(\text{CH}_3)_5\text{Sn}_2\text{OCOCF}_3$ survives sufficiently for it to be observed by ^1H NMR spectroscopy as a substantial component of the reaction mixture from $(\text{CH}_3)_6\text{Sn}_2$. We assign to this trifluoroacetate the two resonances labelled Y and Z in Fig. 4 at δ 0.42 and 0.74 in 3/2 ratio). It was completely decomposed, however, to $(\text{CH}_3)_3\text{SnOCOCF}_3$ and "polymeric dimethyltin" during attempts to recover it from this mixture, and did not survive long enough for satisfactory ^{13}C or ^{119}Sn NMR spectroscopy.

The major product of the trifluoroacetolysis of trimethylgermyltrimethylstannane is $(\text{CH}_3)_3\text{GeSn}(\text{$

case, and makes path A' redundant. Moreover, since trimethylgermane is only observed in systems where the acid concentration is low, it is most likely that path A represents the total Sn-Ge cleavage process in this case, even though we have chosen to analyse the data in terms of both paths A and A'. Probably it is the presence of a suitable leaving group which makes the $(\text{CH}_3)_3\text{SnM}(\text{CH}_3)_3$ substrates reactive, whereas $(\text{CH}_3)_3\text{GeSi}(\text{CH}_3)_3$, $(\text{CH}_3)_6\text{Ge}_2$ and $(\text{CH}_3)_6\text{Si}_2$ are inert. An additional path (B') in which methyl groups are removed from Ge or Si can be rejected on the grounds of insufficient reactivity.

Gaseous products are formed in these reactions, but were not observed in the NMR spectra. Hydrogen is anticipated to show up at δ 4.2 ppm but this spectral region is obscured by the solvent. Methane at δ 0.16 ppm might well be observable in the spectra, but Levi [13] reports that at 25°C one atmosphere pressure of methane over methanol only gives a mole fraction of 7.1×10^{-4} in solution, i.e. $1.5 \times 10^{-2} M$. This would give a signal indistinguishable from the background noise in the spectra shown in Fig. 1 and 2. However methane is clearly identified in the products of trifluoroacetylisis.

The first intermediate formed in path B, $(\text{CH}_3)_3\text{MSn}(\text{$

M = Sn, Ge and Si they do not differ from one another by more than a factor of 10. Attack via path A is faster in general than attack via path B, as judged by the statistically corrected rate constants given in Table 6. The rate constants for Sn-CH₃ cleavage seem to be essentially the same for all three substrates, but Sn-Ge cleavage appears to be significantly slower than Sn-Sn or Sn-Si cleavage.

Although comparative rate data have been obtained for the HCl reactions, our principal aim was simply to identify the possible complications arising in reactions where (CH₃)₃GeCl and (CH₃)₃SiCl were produced and would be solvolysed. These complications arise in an essentially water-free solution rather than the partially aqueous media of the present studies. It may be noted that the "anomalously" high path A/path B ratio for (CH₃)₃GeSn(CH₃)₃ in the least aqueous system accords well with our interpretation of this as a side reaction in other cases [6].

The reactions of these substrates with acetic acid are extremely slow and hence very sensitive to trace reagents and catalytic impurities. While the exclusive Sn-M cleavage observed and the absence of Sn-CH₃ cleavage for related substrates might seem to be appropriate to a highly discriminating reagent, the general similarity of the observed rates argues against this view. The kinetic studies suggest that there is an initial somewhat faster reaction before the first order acetolysis sets in. It is quite likely that the initial phase involves reaction with dissolved oxygen,

constitutes $80(\pm 3)\%$ of the overall reaction. In contrast, we estimate from its observable decomposition products that path B, yielding $(\text{CH}_3)_3\text{SiSn}(\text{CH}_3)_2\text{OCOCF}_3$, proceeds to an extent of only some 30%.

Acknowledgements

These investigations were supported by grants from the Australian Research Grants Scheme and by Commonwealth Postgraduate Awards (to M.J.C. and D.W.H.).

References

- 1 D.C. McWilliam and P.R. Wells, *J. Organomet. Chem.*, 85 (1975) 165.
- 2 D.C. McWilliam and P.R. Wells, *J. Organomet. Chem.*, 85 (1975) 335,347.
- 3 D.P. Arnold and P.R. Wells, *J. Organomet. Chem.*, 108 (1976) 345.
- 4 D.P. Arnold, M.J. Cuthbertson and P.R. Wells, *J. Organomet. Chem.*, 184 (1980) 39.
- 5 M.J. Cuthbertson and P.R. Wells, *J. Organomet. Chem.*, 216 (1981) 331,349.
- 6 M.J. Cuthbertson and P.R. Wells, *J. Organomet. Chem.*, 287 (1985) 25.
- 7 C.F. Shaw and A.L. Allred, *Inorg. Chem.*, 10 (1971) 1340.
- 8 G. Tagliavini, V. Belluco and G. Pill