# Total Synthesis of (+)-Irciniastatin A (a.k.a. Psymberin) and (-)-Irciniastatin B

Chihui An, Jon A. Jurica, Shawn P. Walsh, Adam T. Hoye, and Amos B. Smith, III\*

Department of Chemistry, Laboratory for Research on the Structure of Matter and Monell Chemical Senses Center, University of Pennsylvania, Philadelphia, Pennsylvania 19104, United States

**S** Supporting Information

ABSTRACT: A unified synthetic strategy to access (+)-irciniastatin A (a.k.a. psymberin) and (-)-irciniastatin B, two cytotoxic secondary metabolites, has been achieved. Highlights of the convergent strategy comprise a boron-mediated aldol union to set the C(15)-C(17) syn-syn triad, reagent control to set the four stereocenters of the tetrahydropyran core, and a



late-stage Curtius rearrangement to install the acid-sensitive stereogenic N,O-aminal. Having achieved the total synthesis of (+)-irciniastatin A, we devised an improved synthetic route to the tetrahydropyran core (13 steps) compared to the firstgeneration synthesis (22 steps). Construction of the structurally similar (-)-irciniastatin B was then achieved via modification of a late-stage (-)-irciniastatin A intermediate to implement a chemoselective deprotection/oxidation sequence to access the requisite oxidation state at C(11) of the tetrahydropyran core. Of particular significance, the unified strategy will permit late-stage diversification for analogue development, designed to explore the biological role of substitution at the C(11) position of these highly potent tumor cell growth inhibitory molecules.

## INTRODUCTION

In 2004 two new potent cytotoxins, (+)-irciniastatin A (1) and (-)-irciniastatin B (2), isolated from the Indo-Pacific marine sponge Ircinia ramose, were discovered by Pettit and co-workers (Figure 1).<sup>1</sup> In the same year, a closely related metabolite,



(+)-psymberin, was isolated independently by Crews and coworkers from marine sponge Psamminocinia.<sup>2</sup> Analysis of these reports suggests that irciniastatin A (1), irciniastatin B (2), and psymberin (1) possessed the same architectural features, including a highly substituted 2,6-trans-tetrahydropyran core, a dihydroisocoumarin, and an N,O-aminal. Crews postulated that both irciniastatin A (1) and psymberin might be identical,<sup>2</sup> but unfortunately the NMR spectra of the two congeners were taken in different solvents, and thus the exact stereochemical relationship at C(4) and C(8) could not be established. In 2005, De Brabander and colleagues resolved the structural ambiguity with the first total synthesis of psymberin by construction of all four C(4)-C(8) diastereomers of psymberin.

This effort not only yielded the absolute configuration of (+)-psymberin, but confirmed that both (+)-irciniastatin A (1)and (+)-psymberin possessed identical chemical structures. More recently, De Brabander reported a full account on their first- and second-generation synthesis of (+)-irciniastatin A (1)(a.k.a. psymberin), in conjunction with the synthesis of a series of novel analogues.<sup>4</sup>

The discovery that (+)-irciniastatin A (1) (a.k.a. psymberin) and (-)-irciniastatin B (2) display impressive therapeutic properties, including tumor cell growth inhibition at the nanomolar level (0.7 to 0.8 nM),<sup>1</sup> in conjunction with the observation by Crews et al. that (+)-irciniastatin A (1) displayed high differential cytotoxicity (>10 000-fold) in the NCI 60 human tumor cell panel, raised the intriguing possibility that the observed cytotoxicity might arise via a novel mode of action.<sup>2</sup> Interestingly, even though the chemical structures of (+)-irciniastatin A (1) and (-)-irciniastatin B (2) differ only in the oxidation level at C(11), the ketone congener (2) was reported to be nearly 10 times more active than the alcohol (1) against human pancreas (BXPC-3), breast (MCF-7), and central nervous system (SF268) cancer cell lines.<sup>1</sup> Subsequently, a group at Schering-Plough<sup>5</sup> reported that (+)-C(11)-deoxyanalogue possesses 3-10 times the cytotoxic activity compared to (1). Taken together, these results suggest that the C(11)hydroxyl group is not critical for potent cytotoxic activity.

In 2010 Usui and co-workers reported that the tumor growth inhibition activity of (+)-irciniastatin A (1) arises from activation of stress-activated protein kinases, such as JNK and p38, that in

Received: February 4, 2013

turn leads to apoptosis.<sup>6</sup> Subsequent to this report, De Brabander, in collaboration with Roth, disclosed a forward genetic screen of (+)-irciniastatin A (1) employing *Caenorhabditis elegans* that demonstrated 1 binds to the ribosome to induce cell death.<sup>7</sup> Also of interest, totally synthetic (+)-irciniastatin A (1), from the De Brabander group, did not reveal the high differential cytotoxicity<sup>7</sup> previously reported for natural (+)-irciniastatin A (1).<sup>2</sup>

Given the impressive biological activity, in conjunction with limited natural abundance of the irciniastatins, seven total syntheses<sup>3,8-13</sup> of (+)-irciniastatin A (1), including a report from our laboratory,9 have been disclosed since 2004. After DeBrabander's seminal total synthesis,<sup>3</sup> Floreancig's clever strategy to installing the N,O-aminal to construct (1) proved to be the shortest longest linear sequence to date (14 steps).<sup>13</sup> Somewhat more surprising, the first total synthesis of the substantially more active congener, (-)-irciniastatin B (2), was only recently achieved in our laboratory.<sup>14</sup> Although several SAR studies have shed some light on the mode of action of (+)-irciniastatin A (1), the biological properties associated with the structural differences of the alcohol (1) and the ketone (2)remain unknown. In particular, we were interested in what role the C(11) substituents in the irciniastatins would play in an SAR study. Toward this end, we report here a full account of the synthesis of both (+)-irciniastatin A (1) and (-)-irciniastatin B (2), including our first-generation synthesis of (+)-irciniastatin A (1), a revised second-generation synthesis of the 2,6-transtetrahydropyran core of (+)-irciniastatin A (1), and the subsequent development of a unified strategy, utilizing a latestage selective deprotection/oxidation sequence, which not only led to the first synthesis of (-)-irciniastatin B (2), but also the construction of both C(11) alcohol epimers.

# RESULTS AND DISCUSSION

First-Generation Synthesis of (+)-Irciniastatin A (1). Our strategy to construct (+)-irciniastatin A (1) began with disconnection at the amide linkage, leading to the acid side chain 3 and the Teoc-protected N,O-aminal 4 (Scheme 1). The acid-sensitive N,O-aminal moiety would be installed late in the synthesis, with complete retention of configuration via a Curtius rearrangement, a strategy first developed and successfully exploited in our 2002 synthesis of (+)-zampanolide, bearing a similar N,O-aminal group.<sup>15</sup> Disconnection at C(16)-C(17)next provided aldehyde 5 and 2,6-trans-tetrahydropyran 6, which we envisioned would be united via a substrate-controlled aldol reaction. Aryl aldehyde 5 in turn would derive by a [4 + 2] cycloaddition between known bis-silyl enol ether 7<sup>16</sup> and allene 8,<sup>17</sup> while 2,6-*trans*-tetrahydropyran **6** would arise via a 6-*exo*-tetcyclization of the C(13), hydroxyl with an epoxide that would be installed at the C(8)-C(9). Epoxide 9 in turn would be generated by iterative chemoselective functionalization of diene 10, a synthetic tactic that directs reactivity to the most electronrich olefin, distal to the electron withdrawing ester group. Synthetic strategies that selectively introduce functionality taking advantage of the difference in electron density of the olefinic linkage along a linear polyene, possessing a terminal electronwithdrawing group, have not, as of yet, been widely exploited.<sup>18-21</sup>

Synthesis of the requisite acid side chain (-)-3 began with methyl ether (+)-11,<sup>3</sup> which was constructed from commercially available (+)-isopropylidene glyceraldehyde in a stereocontrolled fashion in two steps; the overall yield was 57% yield (dr > 20:1 Scheme 2). Removal of the acetonide was next achieved by

Scheme 1. Retrosynthetic Analysis



Scheme 2. Synthesis of Acid Side Chain (-)-3



treatment of (+)-11 with aqueous hydrochloric acid. The primary alcohol was then protected chemoselectively as the pivalate ester (+)-12, followed by protection of the secondary alcohol as a SEM ether. Reduction with DIBAL-H then provided primary alcohol (+)-13, which was oxidized via a two-step Parikh-Doering<sup>22</sup>/Pinnick<sup>23</sup> oxidation sequence to provide the desired acid side chain (-)-3.

The requisite aryl aldehyde **5** was constructed via a Diels– Alder cycloaddition between 1,3-bis(trimethylsiloxy)-1,3-diene  $7^{16}$  and dimethyl-1,3-allene-dicarboxylate **8**,<sup>17</sup> followed by a fluoride-mediated aromatization to furnish known homopthalate **14**<sup>24</sup> in 83% yield (Scheme 3). Both phenols were then masked as SEM ethers, followed by chemoselective reduction to furnish aryl aldehyde **5** in an overall yield of 55% for the three-step sequence.

Access to 2,6-*trans*-tetrahydropyran 6, the core of (+)-irciniastatin A (1), began with commercially available *S*-epoxybutane (-)-15. Selection of (-)-15 was simply a matter of synthetic convenience, in order to operate with a single diastereomer, as the advanced alcohol would ultimately be oxidized to a ketone (vide infra). Treatment of epoxide (-)-15 with propynyllithium and BF<sub>3</sub>·OEt<sub>2</sub> provided the homopropargylic alcohol,<sup>25</sup> which was protected to furnish benzyl ether

#### Scheme 3. Synthesis of Dihydroisocoumarin Fragment 5



(-)-16 under acidic conditions. Next, following Negishi's protocol<sup>26</sup> for the in situ generation of the Schwartz reagent from zirconocene dichloride and DIBAL-H, iodination with *N*-iodosuccinimide furnished the external vinyl iodide (-)-17. Pleasingly, the regioselectivity was excellent (>20:1). Vinyl iodide (-)-17 was then united with methyl acrylate, exploiting a palladium-mediated Heck reaction<sup>27</sup> to complete diene (-)-18 in 78% yield (Scheme 4).



Installation of the first of the three epoxides to construct the linear precursor of tetrahydropyran 6 entailed chemoselective epoxidation of the most electron-rich olefin in (-)-18 with the Shi fructose-derived catalyst (-)-19<sup>28</sup> to furnish (+)-20; excellent diastereoselectivity (14:1) resulted (Scheme 5). Regioselective opening was then achieved with trimethylaluminum<sup>29</sup> to provide a single regioisomeric alcohol, which was treated with PMB-Cl to furnish ether (+)-21. Asymmetric epoxidation to introduce the second epoxide required DIBAL-H reduction of ester (+)-21 followed by a Sharpless asymmetric epoxidation.<sup>30</sup> Epoxy alcohol (+)-22 resulted as a single diastereomer in 92% yield for the two steps, which was then subjected to Parikh-Doering oxidation conditions;<sup>22</sup> the resulting aldehyde was then treated with Horner-Wadsworth-Emmons ylide 23 to provide ester (+)-24 in 89% yield (two steps). Palladium-catalyzed hydrogenolysis<sup>31</sup> employing formic acid as the hydrogen source next opened the epoxide in (+)-9, again with excellent chemoselectivity. Subsequent protection of the resulting alcohol as the TBS ether furnished (+)-24 in 92% yield for two steps. The third and final epoxide was installed by reduction of ester (+)-24 with DIBAL-H to reveal the allylic alcohol; Sharpless asymmetric epoxidation,<sup>30</sup> this time

utilizing (–)-diisopropyl tartrate, furnished epoxide (+)-25 with 20:1 *dr*. To complete the construction of the linear precursor for tetrahydropyran 6, alcohol (+)-25 was oxidized<sup>32</sup> directly to the carboxylic acid, followed by treatment with diazomethane to provide the corresponding methyl ester in 76% yield for the two steps. Oxidative removal of the PMB ether protecting group with DDQ completed the construction of (+)-26, the tetrahydropyran cyclization precursor. The overall yield for the 18-step synthetic sequence was 9.3% yield.

With (+)-26 in hand, we examined the acid-promoted cyclization to generate the tetrahydropyran core. Baldwin rules<sup>33</sup> suggest that both the desired 6-exo-tet and undesired 7endo-tet cyclization pathways could operate. However, the sixmembered ring transition state, in conjunction with the electronwithdrawing nature of the ester, destabilizing the partial cationic character at the  $\alpha$ -carbon under Lewis- or Brønsted- acidic conditions, suggested that the tetrahydropyran would predominate. Indeed, treatment with 20 mol % of camphorsulfonic acid (CSA) in methylene chloride achieved the desired 6-exo-tet cyclization pathway, which provided the 2,6-tetrahydropyran (+)-27 in excellent yield (92%), with no trace of the sevenmembered ring congener (Scheme 6). Alcohol (+)-27 was then methylated with dimethyl sulfate, followed by palladiumpromoted hydrogenolysis to remove the benzyl ether. Dess-Martin periodinane<sup>34</sup> oxidation completed construction of the requisite 2,6-trans-tetrahydropyran fragment (+)-6 in 88% yield for the three steps.

Having constructed the three fragments for the proposed synthesis of (+)-irciniastatin A (a.k.a. psymberin) (1), we turned to the union of tetrahydropyran (+)-6 with aryl aldehyde 5 (Scheme 7), exploiting a substrate-controlled aldol reaction. Generation of the Z-boron enolate of (+)-6, achieved by treatment of (+)-6 with dichlorophenylborane,<sup>35</sup> followed by addition of aldehyde 5 furnished (+)-29, the desired *syn*-aldol product, in 90% yield. The stereochemical outcome was dictated by 1,4-substrate stereoinduction.<sup>36</sup> Subsequent chelation-controlled reduction<sup>37</sup> followed by saponification of the methyl ester with concomitant lactonization provided dihydroisocoumarin (+)-**30** in 83% yield for the two steps.

At this juncture we called upon a Curtius rearrangement to install the *N*,*O*-aminal (Scheme 7). Acid (+)-**30** was converted to the corresponding acyl azide, followed by thermal rearrangement in toluene (ca. 80 °C) to provide the isocyanate, which was intercepted by the addition of 2-trimethylsilylethanol to furnish the desired *N*,*O*-aminal, with complete retention of configuration at the methyl ether carbon (NMR). Protection of the remaining free hydroxyl group as the TBS ether was then achieved in 91% yield to complete the construction of advanced amide (+)-**4**, the coupling partner for acid side chain (-)-**3**.

Final fragment union of *N*,*O*-aminal (+)-4 with side chain (-)-3 required considerable experimentation. Eventually we discovered that deprotonation of the Teoc-protected amine (+)-4 with LiHMDS, followed by addition of the side chain, activated as the pivalate anhydride 31, would lead to the desired amide (+)-32 in 79% yield (Scheme 8).

Turning our attention to conditions that would achieve global deprotection while retaining the delicate *N*,*O*-aminal moiety, model studies revealed that TAS-F<sup>38,39</sup> was the reagent of choice. Treatment of the fully protected irciniastatin A (+)-**32** with TAS-F in DMF at 50 °C resulted in two major products (Scheme 8). After purification via preparative TLC, the more polar of the two congeners proved to be (+)-ircinaistatin A (1), while the less polar compound retained one phenolic SEM group. Subjecting

## Scheme 5. Synthesis of Pyran Precursor (+)-26



Scheme 6. Completion of Tetrahydropyran (+)-6



the latter to magnesium bromide<sup>40</sup> resulted in (+)-irciniastatin A (1), furnishing a combined yield of 74% for the two steps. Pleasingly, the spectral data (<sup>1</sup>H and <sup>13</sup>C NMR) of totally synthetic (+)-irciniastatin A (1) proved to be identical in all respects with the spectra of natural (+)-irciniastatin A (1) reported by Pettit<sup>1</sup> and Crews.<sup>2</sup> The total synthesis of (+)-irciniastatin A (a.k.a. psymberin) (1) had thus been achieved with a longest linear sequence of 30 steps (ca. 2.2% overall yield).

Second-Generation Synthesis of (+)-Irciniastatin A (a.k.a. Psymberin). Although we had achieved the total synthesis of (+)-irciniastatin A (1) (a.k.a. psymberin), construction of the core 2,6-*trans*-tetrahydropyran required 22 steps and proceeded with an overall yield of only 7.6%. To provide material for future biological evaluation, as well as to access a series of synthetic analogues, an improved second-

Scheme 7. Fragment Union and Elaboration to Amidation Precursor



generation approach to tetrahydropyran (+)-6 would be required. The new strategy to (+)-6 (Scheme 9) was anticipated

Scheme 8. Completion of (+)-Irciniastatin A (1) (a.ka. Psymberin)



Scheme 9. Revised Retrosynthetic Strategy of Tetrahydropyran (+)-6



to retain the efficient 6-*exo*-tet cyclization of acyclic epoxide precursor 33, which would be constructed via two reagentcontrolled asymmetric transformations from alcohol 34, that in turn would derive via union of aldehyde 35 and ketene acetal 36, exploiting a vinylogous Mukayaima aldol reaction.<sup>41</sup> In this strategy, the *gem*-dimethyl moiety would arise from commercially available 2,2-dimethyl-1,3-propanediol, instead of the epoxide opening strategy employed in our first-generation synthesis. Importantly, the second-generation route would set the requisite stereogenecity in tetrahydropyran (+)-6 via three reagent-controlled asymmetric reactions.

We began the second-generation synthesis of (+)-6 via monoprotection of commercially available 2,2-dimethyl-1,3propanediol 37 (Scheme 10), followed by oxidation of the second hydroxyl employing the Parikh–Doering<sup>22</sup> protocol to provide aldehyde 35. Treatment of aldehyde 35 and ketene acetal  $36^{41}$  employing the chiral oxazaborolidinone derived from Ltryptophan led to a vinylogous Mukaiyama aldol reaction,<sup>41</sup> thereby installing the first stereocenter to furnish (+)-34 as a single enantiomer. Mosher's ester analysis demonstrated the desired (*R*)-isomer was obtained.<sup>42,43</sup> Alcohol (+)-**34** was then protected as the TBS ether, followed by reduction of the methyl ester with DIBAL-H to furnish the corresponding allylic alcohol. Asymmetric epoxidation via the Sharpless<sup>30</sup> protocol next provided the desired  $\beta$ -epoxide (+)-**38** with 13:1 diastereose-lectivity, which in turn was converted directly to the corresponding acid via a one-pot TEMPO<sup>32</sup> oxidation; subsequent methylation led to methyl ester (+)-**39**.

Chemoselective deprotection of the primary TBS ether was then achieved by treatment of (+)-39 with hydrogen fluoride, buffered with pyridine (Scheme 10). The resulting primary alcohol was oxidized<sup>22</sup> to aldehyde (+)-40, and the final stereocenter required for the tetrahydropyran core was introduced via Paterson aldol union<sup>44,45</sup> with 2-butanone, employing (-)-B-chlorodiisopinocampheylborane (DIP-Cl) as the chiral Lewis acid. A 5:1 ( $\beta$ : $\alpha$ ) diastereometric mixture resulted. Cyclization employing a catalytic amount of CSA furnished (+)-41 and (-)-42 exclusively via the 6-exo-tet pathway, again without formation of the seven-membered ring construct. Fortunately, the trans- and cis-diastereomers could now be readily separated by column chromatography to yield 2.6-trans-tetrahydropyran (+)-41 in 74% yield for the 2 steps. Methylation of the secondary hydroxyl group was achieved (92% yield) by treatment with  $Me_3O \cdot BF_4$  and proton sponge [1,8bis(dimethylamino)naphthalene] to complete the secondgeneration synthesis of the 2,6-trans-tetrahydropyran core (+)-6. Pleasingly, the new route to (+)-6 proceeded with a longest linear sequence of 13 steps (a 9-step improvement) and with an overall vield of 17.8% vield, thus more than doubling the overall yield for the longest linear sequence compared to the firstgeneration synthesis.

Total Synthesis of (-)-Irciniastatin B. With an effective route to (+)-irciniastatin A (1), and an improved route to the common tetrahydropyran core, we turned to the construction the biologically more active congener, (-)-irciniastatin B (2). To achieve the requisite ketone oxidation state at C(11), the C(15)secondary hydroxyl in (+)-43, a late-stage intermediate employed in our synthesis of (+)-irciniastatin A (1),<sup>9</sup> was envisioned to be protected as a SEM ether, instead of the TBS ether employed earlier (Scheme 11). This protecting group was selected to permit the critical, selective deprotection of the neopentyl secondary TBS ether at C(11). The secondary alcohol would be oxidized to the requisite ketone, followed by global deprotection to provide access to (-)-irciniastatin B (2). Chemical modification of the late stage C(11) ketone would also permit access to a series of analogues possessing varying substitution at C(11), thus enabling a structure activity relationship study (SAR) to be carried out at this key center in the irciniastatin chemotype.

Surprisingly, protection of the C(15) hydroxyl group in (+)-43 as the SEM ether proved particularly challenging. In particular, manipulation of the resultant SEM ether during workup and purification routinely resulted in the unanticipated loss of the phenolic SEM ethers. Attempts at reprotection proved ineffective, even at elevated temperatures. After extensive experimentation with several model systems, we discovered that the phenolic 3,4-dimethoxybenzyl ether (DMB) would prove durable in the late-stage synthetic sequence with the orthogonally protected C(11) TBS ether.

Synthesis of the revised aryl fragment **45** thus began with protection of bis-phenol  $14^{24}$  with DMB-Br (Scheme 12). Reduction of the resulting ester with DIBAL-H furnished the aryl

Scheme 10. Second-Generation Synthetic Route towards Tetrahydropyran (+)-6



Scheme 11. Divergent Strategy Toward the Synthesis of (–)-Irciniastatin B



aldehyde **45**. From here, the synthetic route continued in similar fashion to the sequence leading to (+)-irciniastatin A (1).<sup>9</sup> Aldol union<sup>35</sup> between aldehyde **45** and ketone (+)-**6** pleasingly furnished  $\beta$ -hydroxyketone (+)-**46** in 70% yield, with minor concomitant lactonization (10:1) and excellent diastereoselectivity (>20:1).<sup>36</sup> Chelation-controlled reduction<sup>37</sup> resulted in a mixture of the desired *syn* diol and the corresponding lactone (ca. 8:1). The mixture was treated under saponification conditions to provide acid (+)-**47** in 69% yield for the 2 steps. With acid (+)-**47** 

in hand, the corresponding acyl azide was generated and subjected to the conditions for the Curtius rearrangement<sup>15</sup> to furnish the Teoc-protected *N*,*O*-aminal in 67% yield, again with complete retention of stereochemical configuration at C(8). The resulting secondary alcohol was then protected as the SEM ether (+)-48 in 82% yield. Importantly, workup and/or purification proceeded without the formation of undesired side products, compared to the earlier SEM protection strategy.

Achieving the requisite amide union to provide (+)-50 once again proved challenging (Scheme 13). The conditions employed in our earlier (+)-irciniastatin A (1) synthesis,<sup>9</sup> involving LiHMDS as the base with the mixed anhydride 31, resulted in low yields (ca. 15%). After significant screening, the conditions employed by Crimmins and co-workers<sup>10</sup> in their synthesis of (+)-irciniastatin A (1), namely, the use of *i*-PrMgCl as the base and acid chloride 49, provided the desired amide (+)-50 in 72% yield.

Having arrived at the full carbon skeleton of (-)-irciniastatin B (2), we set out to effect selective deprotection of the hindered neopentyl secondary C(11) TBS ether (Scheme 13). The TBS ether (+)-**50** was treated with TBAF at room temperature, which resulted in hydrolysis of the Teoc carbamate. Subsequent warming the reaction mixture to 50 °C then led to selective removal of the C(11) TBS group in an overall yield of 79%. Oxidation with Dess–Martin periodinane<sup>34</sup> provided ketone (-)-**51**.

In order to remove the two pairs of protecting groups in (-)-51 and complete the synthesis of (2), a two-stage deprotection protocol was required. Introduction of the ketone

Article





moiety in the tetrahydropyran, however, greatly enhances the sensitivity of the molecular structure. For example, treatment of (-)-51 with base readily initiates a retro-Michael/Michael addition, effecting epimerization of C(9) of the tetrahydropyran core,<sup>46</sup> while acid treatment results in hydrolysis of the *N*,*O*-aminal.<sup>47</sup> Fortunately, treatment of ketone (-)-51 with DDQ



did provide the desired bis-phenol without observable decomposition. Removal of the two remaining SEM groups with either TAS- $F^{38,39}$  or TBAF, however, resulted only in complex mixtures, highlighting the base sensitivity of ketone (-)-**51**. Again, after significant experimentation, we discovered that treatment of the bis-phenol with a premixed solution of MgBr<sub>2</sub>, *n*-butanethiol, and nitromethane in ether<sup>48</sup> removed the SEM ethers cleanly to provide (-)-irciniastatin B (**2**) in 78% yield for the two steps. Pleasingly, synthetic (-)-irciniastatin B (**2**) proved to be identical in all regards (<sup>1</sup>H and <sup>13</sup>C NMR) with the spectral data obtained by Pettit and co-workers<sup>1</sup> and thus constituted the first total synthesis of (-)-irciniastatin B (**2**).

In order to verify the structural relationship of (-)-irciniastatin B (2) with (+)-irciniastatin A (1), we carried out a chemical interconversion of (2) to (1) (Scheme 14). To this end, 2 was

Scheme 14. Structural Confirmation of (-)-Irciniastatin B (2) by Chemical Conversion to (+)-Irciniastatin A (1) (a.k.a. Psymberin)





treated with NaBH<sub>4</sub>, which resulted in a mixture (1:1) of (+)-irciniastatin A 1 and *epi*-C(11)-irciniastatin A (**52**). The two diastereomers were separated via preparative TLC, and the spectral data of the faster moving diastereomer (TLC) was identical with the spectral data of (+)-irciniastatin A (i.e., <sup>1</sup>H, <sup>13</sup>C NMR and HRMS), thereby confirming the structural relationship of (+)-irciniastatin A (1) and (-)-irciniastatin B (2).

# **SUMMARY**

The evolution of an effective unified synthetic strategy for the construction (+)-irciniastatin A (1) (a.k.a. psymberin) and (-)-irciniastatin B (2) has been achieved. The first-generation synthesis of (+)-ircinastatin A required 30 steps and proceeded in 2.2% overall yield. Of these 30 steps, 22 were required to construct the 2,6-trans-tetrahydropyran, which proceeded in 7.6% overall yield. A significantly improved second-generation synthesis of the 2,6-trans-tetrahydropyran core was subsequently achieved, which now permits an efficient total synthesis of both (+)-irciniastatin A (1) and (-)-irciniastatin B (2), the latter requiring a chemoselective deprotection/oxidation sequence. Finally, the structural relationship of the two similar metabolites has been confirmed via chemical conversion of (-)-irciniastatin B (2) to (+)-irciniastatin A (1) and the corresponding C(11) epimer (52). Importantly, the successful synthesis leading to (-)-irciniastatin B (2) now holds the promise for the elaboration of C(11)-irciniastatin analogues, currently ongoing in our laboratory.

### EXPERIMENTAL SECTION

**Materials and Methods.** Reactions were carried out in flame-dried or oven-dried glassware under a nitrogen atmosphere unless noted otherwise. Anhydrous diethyl ether ( $Et_2O$ ), tetrahydrofuran (THF), dichloromethane ( $CH_2Cl_2$ ) and toluene were obtained from a solvent purification system. All commercially available reagents were used without purification unless otherwise noted. Triethylamine, diisopropylethylamine, and pyridine were freshly distilled from calcium hydride under a nitrogen atmosphere. Reactions were magnetically stirred unless stated otherwise and monitored by thin layer chromatography (TLC) with 0.25 mm Silacycle precoated silica gel plates. Silica gel chromatography was performed utilizing ACS grade solvents and silica gel from either Silacycle or Sorbent Technologies.

Infrared spectra were obtained using an FT/IR spectrometer. Optical rotations were obtained using a polarimeter. All melting points were obtained on a melting point apparatus and are uncorrected. <sup>1</sup>H NMR spectra (500 MHz field strength) and <sup>13</sup>C NMR spectra (125 MHz field strength) were obtained on 500 MHz spectrometer or a cryomagnet (500 MHz/52 mm) with a 5 mm dual cryoprobe. Chemical shifts are reported relative to chloroform ( $\delta$  7.26) or methanol ( $\delta$  4.78) for <sup>1</sup>H NMR spectra and chloroform ( $\delta$  77.23), methanol ( $\delta$  49.3), or benzene ( $\delta$  128.0) for <sup>13</sup>C spectra. High-resolution mass spectra (HRMS) were measured on an LC-TOF mass spectrometer

**Pivalate Ester (+)-12.** A 2 N HCl solution (10 mL) was added to acetonide (+)-11 (0.836 g, 4.18 mmol) at rt. After 30 min, the reaction was quenched with sat. NaHCO<sub>3</sub> until all gas evolution had ceased. The reaction was then extracted with EtOAc, and the combined organic layers were dried over MgSO<sub>4</sub> and concentrated. Flash chromatography (40% EtOAc/hexanes) provided diol (+)-**53** (0.650 g, 97% yield) as a colorless oil:  $[\alpha]_D^{20}$  +21.8 (*c* 1.5, CHCl<sub>3</sub>); IR (neat) 3416, 2932, 1646, 1456, 1092, 1050 cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  4.84 (q, *J* = 1.6 Hz, 1H), 4.80 (m, 1H), 3.82–3.77 (m, 1H), 3.72–3.67 (m, 2H), 3.57–3.53 (ddd, *J* = 2.4, 6.5, 13.1 Hz, 1H), 3.43 (s, 3H), 2.61 (d, *J* = 6.7 Hz, 1H), 2.39 (dd, *J* = 14.0, 6.8 Hz, 1H), 2.32 (dd, *J* = 7.2, 3.2 Hz, 1 H), 2.18 (dd, *J* = 14.3, 6.4 Hz, 1H), 1.79 (ap t, *J* = 1.1 Hz, 3H); <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  142.49, 113.2, 81.7, 72.8, 63.3, 58.4, 38.9, 22.9; high resolution mass spectrum (CI+) *m/z* 161.1177 [(M + H)<sup>+</sup>; calcd for C<sub>8</sub>H<sub>17</sub>O<sub>3</sub> 161.1178].

Diol (+)-53 (0.243 g, 1.52 mmol) in pyridine (4 mL) was cooled to 0 °C, followed by addition of trimethylacetyl chloride (0.205 mL, 1.67 mmol). The reaction was allowed to warm to rt, and after 1 h, the reaction was cooled to 0 °C, and H<sub>2</sub>O (5 mL) was added. The reaction was diluted with Et<sub>2</sub>O (5 mL) and washed successively with 5 mL each of sat. NaHCO<sub>3</sub>, 1 N HCl, and brine. The organic layer was then dried over MgSO<sub>4</sub> and concentrated. Flash chromatography (10% EtOAc/ hexanes) provided (+)-12 (0.325 g, 88% yield) as colorless oil:  $\left[\alpha\right]_{D}^{20}$ +23.9 (c 1.1, CHCl<sub>3</sub>); IR (neat) 3467, 2971, 2935, 1728, 1457, 1286, 1163, 1103 cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ 4.84 (s, 1H), 4.81 (s, 1H), 4.23 (dd, J = 11.7, 3.5 Hz, 1H), 4.16 (dd, J = 11.6, 6.8 Hz, 1H), 3.92-3.82 (m, 1H), 3.48-3.39 (m, 1H), 3.40 (s, 3H), 2.39 (d, J = 4.8 Hz, 1H), 2.29 (ddd, I = 19.4, 14.4, 6.2 Hz, 2H), 1.79 (s, 3H), 1.22 (s, 9H); <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  179.0, 142.6, 113.3, 80.6, 71.4, 65.6, 58.4, 39.0, 38.4, 27.4, 23.0; high resolution mass spectrum (ES+) m/z 245.1743 [(M + H)<sup>+</sup>; calcd for C<sub>13</sub>H<sub>25</sub>O<sub>4</sub> 245.1753].

Alcohol (+)-13. A solution of (+)-12 (0.221 g, 0.905 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (3 mL) was cooled to 0 °C, and *i*-Pr<sub>2</sub>NEt (0.395 mL, 2.5 equiv) was added, followed by dropwise addition of SEMCl (0.320 mL, 2.0 equiv). The reaction was allowed to warm to rt, and after 2 h, sat. NH<sub>4</sub>Cl was added. The reaction was extracted with  $CH_2Cl_2$  (3 × 10 mL), and the combined organic layers were dried over MgSO<sub>4</sub> and concentrated. Flash chromatography (5% EtOAc/hexanes) provided SEM ether (-)-54 (338 mg, 99% yield) as colorless oil:  $[\alpha]_{D}^{20}$  -6.3 (*c* 2.9, CHCl<sub>3</sub>); IR (neat) 3457, 2925, 1728, 1480, 1283, 1249, 1159, 1107, 1031 cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  4.81 (s, 1H), 4.78 (s, 1H), 4.74 (dd, J = 14.7, 6.9 Hz, 2H), 4.29 (dd, J = 11.8, 3.9 Hz, 1H), 4.11 (dd, J = 11.8, 6.2 Hz, 1H), 3.84–3.79 (m, 1H), 3.68 (ddd, J = 10.1, 6.3, 6.3 Hz, 1H), 3.59 (ddd, J = 10.0, 6.5, 6.5 Hz, 1H), 3.51-3.46 (m, 1H), 3.39 (s, 3H), 2.32-2.19 (m, 2H), 1.77 (s, 3H), 1.19 (s, 9H) 0.99-0.82 (m, 2H), 0.00 (s, 9H); <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>) δ 178.4, 142.7, 113.0, 95.0, 80.3, 76.8, 65.6, 64.0, 58.6, 39.4, 39.0, 27.4, 23.0, 18.2, -1.2; high resolution mass spectrum (ES+) m/z 397.2372 [(M + Na)<sup>+</sup>; calcd for  $C_{19}H_{38}O_{5}SiNa 397.2387$ ].

A solution of SEM ether (–)-54 (0.339g, 0.905 mmol) in  $CH_2Cl_2$ (4.5 mL) was cooled to -78 °C, and DIBAL-H (2.0 mL, 1 M in toluene, 2.2 equiv) was added dropwise. After 5 min, the reaction was quenched with MeOH (0.5 mL). The reaction was allowed to warm to rt before EtOAc (5 mL) and sat. Rochelle's salt (5 mL) were added. After 1 h, the organic layer transitioned from cloudy to clear. The layers were separated, and the aqueous layer extracted with EtOAc  $(3 \times 10 \text{ mL})$ . The combined organic layers were dried over MgSO<sub>4</sub> and concentrated. Flash chromatography (10% EtOAc/hexanes) provided (+)-13 (0.249 g, 95% yield) as colorless oil:  $[\alpha]_{D}^{20}$  +31.7 (c 1.2, CHCl<sub>3</sub>); IR (neat) 3457, 2952, 2925, 2892, 1650, 1457, 1378, 1249, 1102, 1054, 1025 cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  4.82 (s, 1H), 4.78 (s, 1H), 4.76 (dd, J = 14.3, 7.0 Hz, 2H), 3.79–3.68 (m, 3H), 3.65–3.56 (m, 2H), 3.49 (ddd, J = 7.6, 4.8, 4.8 Hz, 1H), 3.41 (s, 3H), 3.21 (dd, J = 8.3, 4.2 Hz, 1H), 2.31 (dd, J = 14.4, 7.6 Hz, 1H), 2.21 (dd, J = 14.4, 5.3 Hz, 1H), 1.77 (s, 3H), 0.99-0.92 (m, 2H), 0.02 (s, 9H); <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>) δ 142.6, 113.2, 95.6, 82.6, 81.0, 66.0, 62.7, 58.6, 39.5, 23.0, 18.3, -1.3; high resolution mass spectrum (ES+) m/z 313.1821 [(M + Na)<sup>+</sup>; calcd for C<sub>14</sub>H<sub>30</sub>O<sub>4</sub>SiNa 313.1811].

Acid (-)-3. A solution of (+)-13 (0.117 g, 0.404 mmol) in DMSO (0.29 mL, 10 equiv) and CH<sub>2</sub>Cl<sub>2</sub> (4 mL) was cooled to 0 °C, and i- $Pr_2NEt (0.212 \text{ mL}, 3 \text{ equiv})$  was added followed by  $SO_3$  pyridine (0.193) g, 3 equiv) in one portion. After 5 min, brine (10 mL) and  $H_2O$  (2 mL) were added, and the reaction was warmed to rt. The layers were separated, and the aqueous layer was extracted with  $CH_2Cl_2$  (3 × 10 mL). The combined organic layers were dried over MgSO4 and concentrated. Flash chromatography (5% EtOAc/hexanes) provided aldehyde (-)-55 (0.113 g, 97%) as a colorless oil:  $[\alpha]_{\rm D}^{20}$  -9.1 (c 0.9, CHCl<sub>3</sub>); IR (neat) 2952, 2892, 2825, 1732, 1450, 1376, 1249, 1108, 1060, 1029 cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  9.66 (d, J = 1.4 Hz, 1H), 4.85 (s, 1H), 4.84 (s, 1H), 4.80 (dd, J = 19.4, 6.9 Hz, 2H), 4.11 (dd, *J* = 2.8, 1.5 Hz, 1H), 3.75–3.69 (m, 2H), 3.65 (ddd, *J* = 9.8, 9.8, 6.9 Hz, 1H), 3.41 (s, 3H), 2.33 (dddd, J = 14.1, 14.1, 14.1, 6.9 Hz, 2H), 1.72 (s, 3H), 0.92 (ddd, J = 10.1, 6.6, 3.3 Hz, 2H), 0.02 (s, 9H); <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  202.4, 141.7, 114.4, 96.3, 82.5, 81.8, 66.1, 58.1, 38.9, 22.8, 18.2, 1.2; high resolution mass spectrum (ES+) m/z 311.1666 [(M + Na)<sup>+</sup>; calcd for C<sub>14</sub>H<sub>28</sub>O<sub>4</sub>SiNa 311.1655].

Aldehyde (-)-55 (0.203 g, 0.704 mmol) was dissolved in t-BuOH (7.5 mL) and H<sub>2</sub>O (7.5 mL). The solution was cooled to 0 °C followed by addition of 2-methyl-2-butene (6 mL),  $NaH_2PO_4 \cdot H_2O$  (0.550 g, 5 equiv), and NaClO<sub>2</sub> (0.483 g, 80 wt %, 5 equiv). After 15 min, the reaction was poured onto sat. NH<sub>4</sub>Cl (10 mL) and extracted thoroughly with EtOAc. The combined organic layers were dried over MgSO4 and concentrated. Flash chromatography (25% EtOAc/hexanes to 40% EtOAc/hexanes) provided (-)-3 (0.196 g, 92% yield) as a colorless oil:  $\left[\alpha\right]_{D}^{20}$  -18.6 (c 1.1, CHCl<sub>3</sub>); IR (neat) 2953, 2925, 1725, 1649, 1376, 1252, 1110, 1060, 1030 cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  4.83 (s, 1H), 4.81 (s, 1H), 4.78 (s, 2H), 4.39 (d, J = 3.1 Hz, 1H), 3.75 (ddd, J = 8.2, 5.2, 3.2 Hz, 1H), 3.68 (dd, J = 8.5, 8.5 Hz, 2H), 3.43 (s, 3H), 2.39 (dd, *J* = 14.6, 8.0 Hz, 1H), 2.28 (dd, *J* = 14.6, 5.2 Hz, 1H), 1.76 (s, 3H), 0.96-0.87 (m, 2H), 0.01 (s, 9H); <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>) δ 175.0, 141.9, 113.5, 95.1, 81.0, 76.2, 66.3, 58.3, 38.7, 22.9, 18.2, -1.3; high resolution mass spectrum (ES+) m/z 327.1614 [(M + Na)<sup>+</sup>; calcd for C14H28O5SiNa 327.1604].

Aldehyde 5. A solution of homophthalate 14 (0.305 g, 1.20 mmol) in THF (6 mL) was cooled to 0 °C, and NaH (0.106 g, 60 wt %, 2.2 equiv) was added. After 5 min, SEMCl (0.53 mL, 2.5 equiv) was added dropwise. The reaction was allowed to warm to rt, and after 30 min the reaction was quenched with slow addition of MeOH (0.5 mL). Sat. NaHCO<sub>2</sub> (5 mL) was added, and the reaction was extracted with  $CH_2Cl_2$  (3 × 10 mL). The combined organic layers were then dried over MgSO<sub>4</sub> and concentrated. Flash chromatography (10% EtOAc/ hexanes) provided bis-SEM ether 56 (0.611 g, 99% yield) as a colorless oil: IR (neat) 2952, 2898, 1736, 1597, 1314, 1266, 1250, 1158, 1067 cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  6.94 (s, 1H), 5.22 (s, 2H), 5.17 (s, 2H), 3.85 (s, 3H), 3.77-3.69 (m, 4H), 3.67 (s, 5H), 2.12 (s, 3H), 0.98-0.91 (m, 4H), 0.00 (s, 9H), 0.00 (s, 9H); <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>) δ 171.1, 168.6, 157.4, 153.8, 132.3, 120.7, 119.0, 101.9, 94.0, 93.4, 66.5, 66.5, 52.2, 52.2, 36.3, 18.3, 18.2, 11.7, -1.2; high resolution mass spectrum (ES+) m/z 537.2303 [(M + Na)<sup>+</sup>; calcd for C<sub>24</sub>H<sub>42</sub>O<sub>8</sub>Si<sub>2</sub>Na 537.2316].

A solution of bis-SEM ether 56 (0.104 g, 0.202 mmol) in  $CH_2Cl_2$  (2.0 mL) was cooled to -78 °C, and DIBAL-H (0.22 mL, 1 M solution in toluene, 1.1 equiv) was added dropwise over 15 min. After 5 min, the reaction was quenched with MeOH. The reaction was allowed to warm to rt before EtOAc (5 mL) and sat. Rochelle's salt (5 mL) were added. After 1 h, the organic layer transitioned from cloudy to clear. The layers were separated, and the aqueous layer extracted with  $3 \times 20$  mL EtOAc. The combined organic layers were dried over MgSO<sub>4</sub> and concentrated. Flash chromatography (10% EtOAc/hexanes) provided 5 (0.64 g, 66% yield) as a light yellow oil: IR (neat) 2953, 2901, 1726, 1595, 1477, 1272, 1251, 1157, 1110, 1065 cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) 9.62 (dd, J =1.8, 1.8 Hz, 1H), 6.97 (s, 1H), 5.24 (s, 2H), 5.19 (s, 2H), 3.85 (s, 3H), 3.79–3.71 (m, 4H), 3.64 (d, J = 1.8 Hz, 2H), 2.08 (s, 3H), 0.99–0.91 (m, 4H), 0.00 (s, 9H), 0.00 (s, 9H);  $^{13}$ C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$ 198.7, 168.7, 157.7, 154.0, 130.5, 120.8, 119.3, 102.1, 93.9, 93.4, 66.6, 66.6, 52.3, 46.1, 18.3, 18.2, 11.9, -1.2; high resolution mass spectrum (ES+) m/z 507.2210 [(M + Na)<sup>+</sup>; calcd for C<sub>23</sub>H<sub>40</sub>O<sub>7</sub>Si<sub>2</sub>Na 507.2211].

Alkyne (-)-16. To solution of THF (600 mL), cooled to -78 °C, was added propyne (100 g, 4 equiv) via subsurface cannula to dissolve the gas in the THF. Next, a solution of n-BuLi (570 mL, 2.2 M in hexanes, 2 equiv) was cooled to -78 °C and added to the propyne solution over 1 h via cannula. After stirring for 1.5 h at -78 °C (S)-1,2epoxybutane [(-)-15] (45 g, 0.625 mol) was added via cannula over 25 min followed by addition of BF3·OEt2 over 45 min via cannula. After stirring for 1.5 h at -78 °C, the reaction was quenched with sat. NaHCO<sub>3</sub> until all gas evolution ceased (ca. 3 h). The layers were separated, and the aqueous layer extracted with  $Et_2O$  (3 × 500 mL). The combined organic layers were dried over MgSO4 and concentrated in a 0 °C bath (to minimize loss of product due to slight volatility) to provide alcohol (+)-57 (55.6 g, 79%) as a colorless oil:  $[\alpha]_{D}^{20}$  +11.0 (c 1.3, CHCl<sub>3</sub>); IR (neat) 3382, 2964, 2922, 1461, 1435, 1335, 1245, 1112, 1063, 1020 cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  3.69–3.55 (m, 1H), 2.37 (ddddd, J = 16.4, 4.9, 2.5, 2.5, 2.5 Hz, 1H), 2.24 (ddddd, J = 16.4, 7.2, 2.5, 2.5, 2.5 Hz, 1H), 1.95 (d, J = 4.8 Hz, 1H), 1.80 (dd, J = 2.5, 2.5 Hz, 3H), 1.64–1.48 (m, 2H), 0.94 (dd, J = 7.5, 7.5 Hz, 3H); <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  78.3, 75.5, 71.7, 29.2, 27.3, 10.1, 3.6; low resolution mass spectrum (ES+) m/z 112.10 [(M + )<sup>+</sup>; calcd for C<sub>7</sub>H<sub>12</sub>O 112.0888].

To a solution of alcohol (+)-57 (18.5 g, 0.165 mol) in CH<sub>2</sub>Cl<sub>2</sub> (184 mL) and cyclohexane (368 mL) at rt was added benzyl-2,2,2trichloroacetimidate (50.0 g, 1.2 equiv) followed by dropwise addition of TfOH (0.73 mL, 0.05 equiv) The solution turned cloudy and light brown as the reaction progressed. After 6 h, hexane (600 mL) was added, and the solution stirred 30 min to precipitate solids, and then the suspension was filtered. The filtrate was washed with sat. NaHCO<sub>3</sub> (200 mL), which was then back extracted with  $CH_2Cl_2$  (3 × 100 mL). The combined organic layers were then dried over MgSO<sub>4</sub> and concentrated. The concentrate was diluted with 10% EtOAc/hexanes (200 mL) to facilitate further precipitation of solids and then filtered. The filtrate was then concentrated and purified by flash chromatography (0.5% EtOAc/ hexanes to 1% EtOAc/hexanes) to provide alkyne (-)-16 (30.4 g, 91%) yield) as a light yellow oil:  $[\alpha]_{D}^{20} - 27.6$  (c 1.4, CHCl<sub>3</sub>); IR (neat) 2964, 2919, 2873, 1496, 1454, 1348, 1207, 1108, 1071, 1028 cm<sup>-1</sup>; <sup>1</sup>H NMR  $(500 \text{ MHz}, \text{CDCl}_3) \delta 7.44 - 7.22 \text{ (m, 5H)}, 4.68 \text{ (dab, } J = 11.9 \text{ Hz}, 1\text{H}),$ 4.57 (dab, J = 11.8 Hz, 1H), 3.45 (dddd, J = 6.7, 6.7, 5.1, 5.1 Hz, 1H), 2.45 (ddddd, J = 16.5, 5.1, 2.5, 2.5, 2.5 Hz, 1H), 2.41-2.32 (m, 2H), 1.81 (dd, J = 2.5, 2.5 Hz, 3H), 1.77–1.60 (m, 2H), 0.97 (dd, J = 7.4, 7.4 Hz, 3H); <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>) δ 138.9, 128.5, 127.9, 127.7, 79.1, 77.2, 76.1, 71.4, 26.8, 23.8, 9.8, 3.7; high resolution mass spectrum (ES +) m/z 202.1354 [(M)<sup>+</sup>; calcd for C<sub>14</sub>H<sub>19</sub>O 202.1358].

Vinyl lodide (-)-17. To a flask protected from light was added bis(cyclopentadienyl) zirconium(IV) chloride (0.313 g, 2.0 equiv) and THF (1 mL). The solution was cooled to 0 °C, and DIBAL-H (1.07 mL, 1 M in hexanes, 2.0 equiv) was added dropwise. After 30 min at 0 °C, alkyne (-)-16 (0.108g, 0.534 mmol) was dissolved in THF (0.3 mL) and added to the in situ generated Schwartz reagent. The flask and syringe were then flushed with THF (0.3 mL) into the reaction. The reaction flask was then placed in a 50 °C oil bath. After 1 h, the reaction was cooled to 0 °C, and N-iodosuccinimide (0.265 g, 2.2 equiv) was added. After 10 min, the reaction was quenched with sat. NaHCO<sub>3</sub> (10 mL) and then filtered through a 1 cm plug of silica gel. The silica gel was rinsed with EtOAc, and the layers separated. The aqueous layer was then extracted with EtOAc ( $3 \times 10$  mL), and the combined organic layers were dried over  $MgSO_4$  and concentrated. Flash chromatography (10%) EtOAc/hexanes) provided vinyl iodide (-)-17 (135 mg, 77% yield, >20:1 selectivity) as a colorless oil:  $[\alpha]_D^{20}$  –4.4 (*c* 1.0, CHCl<sub>3</sub>); IR (neat) 3030, 2963, 2931, 2872, 1454, 1351, 1092, 1065, 1028 cm<sup>-1</sup>; <sup>1</sup>H NMR  $(500 \text{ MHz}, \text{CDCl}_3) \delta 7.42 - 7.27 \text{ (m, 5H)}, 6.24 \text{ (dd, } J = 7.5, 6.8 \text{ Hz}, 1\text{H}),$ 4.54 (s, 2H), 3.37 (dddd, J = 5.9, 5.9, 5.9, 5.9 Hz, 1H), 2.39 (s, 3H), 2.36–2.17 (m, 2H), 1.62–1.52 (m, 2H), 0.95 (dd, *J* = 7.4, 7.4 Hz, 3H).;  $^{13}\mathrm{C}$  NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  138.9, 137.7, 128.6, 127.9, 127.7, 95.3, 79.3, 71.4, 34.8, 27.9, 26.8, 9.9.; high resolution mass spectrum (ES+) m/z 353.0386 [(M + Na)<sup>+</sup>; calcd for C<sub>14</sub>H<sub>19</sub>IONa 353.0378].

Diene (-)-18. Vinyl iodide (-)-17 (6.02 g, 18.2 mmol) was dissolved in anhydrous DMF (70 mL) in a sealable tube. Methyl acrylate (2.35 g, 1.5 equiv) was added followed by  $Pd(OAc)_2$  (0.817 g, 0.20 equiv), NaHCO<sub>3</sub> (3.06 g, 2.0 equiv), and Bu<sub>4</sub>NI (6.73 g, 1.0 equiv). The tube was flushed with argon, sealed, and heated to 100 °C over 1 h. After 7 h at 100 °C, the reaction was cooled to 0 °C, and sat. NH<sub>4</sub>Cl (100 mL) added. The reaction was then filtered and extracted with EtOAc (3  $\times$ 100 mL). The combined organic layers were washed with H<sub>2</sub>O, sat. NaHCO<sub>3</sub>, and brine (100 mL each). The organic layer was then dried over MgSO<sub>4</sub> and concentrated. Flash chromatography (3% EtOAc/ hexanes to 5% EtOAc/hexanes to 8% EtOAc/hexanes) provided diene (-)-18 (4.11 g, 78% yield) as a light yellow oil:  $[\alpha]_D^{20}$  -11.7 (c 1.0, CHCl<sub>3</sub>); IR (neat) 3033, 2963, 2873, 1719, 1624, 1310, 1194, 1169, 1123, 1097 cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  7.34–7.26 (m, 5H), 5.97 (dd, *J* = 8.0, 8.0 Hz, 1 H), 5.81 (d, *J* = 15.5 Hz, 1H), 4.53 (s, 3H), 3.77 (s, 3H), 3.47 (ddd, J = 12.0, 6.0, 6.0 Hz, 1H), 2.51–2.43 (m, 2H), 1.78 (s, 3H), 1.57–1.64 (m, 3H), 0.94 (dd, J = 7.5, 7.5 Hz, 3H); <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  168.2, 149.9, 138.9, 138.4, 134.3, 128.6, 127.9, 127.8, 115.6, 79.8, 71.4, 51.7, 33.3, 27.0, 12.6, 9.9; high resolution mass spectrum (ES+) m/z 289.1823 [(M + H)<sup>+</sup>; calcd for C<sub>18</sub>H<sub>25</sub>O<sub>3</sub> 289.1804].

**Epoxide (+)-20.** Diene (-)-18 (2.94 g, 10.2 mmol) was dissolved in CH<sub>3</sub>CN (460 mL). Na<sub>2</sub>B<sub>4</sub>O<sub>7</sub> buffer (0.05 M, 115 mL) was added followed  $Bu_4NHSO_4$  (0.425 g) and ketone catalyst (-)-19 (1.32 g, 0.50 equiv). Oxone (8.77g, 1.4 equiv) was dissolved in Na2EDTA buffer (59 mL,  $4 \times 10^{-4}$  M), and K<sub>2</sub>CO<sub>3</sub> (8.31 g, 5.9 equiv) was dissolved in H<sub>2</sub>O (59 mL). The two solutions were then added simultaneously over 3 h via a dual-syringe pump. After the addition was complete, water was added to dissolve any solids that had formed, and the reaction extracted with EtOAc (3  $\times$  300 mL). The combined organic layers were washed with brine (200 mL), dried over MgSO4, and concentrated. Flash chromatography (5% EtOAc/hexanes to 8% EtOAc/hexanes) provided epoxide (+)-20 (2.27 g, 73% yield,  $\beta:\alpha = 14:1$ ) as a colorless oil:  $\left[\alpha\right]_{D}^{20}$ +2.2 (c 1.7, CHCl<sub>3</sub>); IR (neat) 2966, 2932, 2876, 1726, 1654, 1436, 1311, 1170, 1094 cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ 7.34–7.31 (m, 5H), 6.75 (d, J = 15.5 Hz, 1H), 6.00 (d, J = 16.0 Hz, 1H), 4.6 (dab, J = 11.5 Hz, 1 H), 4.52 (dab, J = 11.5 Hz, 1 H), 3.75 (s, 3H), 3.59-3.55 (m, 1H), 3.00 (dd, J = 6.5, 6.5 Hz, 1H), 1.85 (ddd, J = 14.5, 8.0, 5.5 Hz, 1H), 1.76 (ddd, J = 4.5, 6.5, 4.5 Hz, 1H), 1. 68–1.60 (m, 2H), 1.41 (s, 3H), 0.95 (dd, J = 7.5, 7.5 Hz, 3H); <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  166.8, 150.4, 138.8, 128.6, 128.0, 127.9, 121.3, 78.0, 71.6, 63.8, 58.9, 51.9, 33.2, 27.0, 15.7, 9.5; high resolution mass spectrum (ES+) *m/z* 327.1578 [(M + Na)<sup>+</sup>; calcd for  $C_{18}H_{24}O_4Na$  327.1572].

**PMB Ether (+)-21.** To a solution of compound (+)-20 (60 mg, 0.20 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (3 mL) was added H<sub>2</sub>O (21  $\mu$ L, 6 equiv) The reaction was cooled to -40 °C, and Me<sub>3</sub>Al (0.98 mL, 10 equiv, 2 M in hexanes) was added. After 30 min, the reaction was warmed to 0 °C, and the reaction quenched with slow addition of H<sub>2</sub>O (1 mL) followed by addition of 1 N HCl (3 mL) to break up the emulsion. The reaction was extracted with  $CH_2Cl_2$  (3 × 10 mL), and the combined organic layers were dried over MgSO<sub>4</sub> and concentrated. Flash chromatography (5% EtOAc/hexanes to 10% EtOAc/hexanes) provided alcohol (+)-58 (53 mg, 84% yield) as a colorless oil:  $[\alpha]_D^{20}$  +31.7 (c 1.6, CHCl<sub>3</sub>); IR (neat) 3490, 2964, 2875, 1723, 1651, 1435, 1314, 1201, 1173, 1060 cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz,  $CDCl_3$ ) 7.38–7.26 (m, 5H), 7.04 (d, J = 16.1 Hz, 1H), 5.81 (d, J = 16.1 Hz, 1H), 4.60 (dab, J = 11.6 Hz, 1H), 4.52 (dab, J = 11.6 Hz, 1H), 3.73 (s, 3H), 3.70 (d, J = 8.9, 1H), 3.65–3.59 (m, 1H), 2.75 (s, 1H), 1.91–1.66 (m, 1H), 1.65–1.50 (m, 3H), 1.07 (s, 3H), 1.06 (s, 3H), 0.91 (dd, J = 7.5, 7.5 Hz, 3H); <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  167.5, 155.9, 138.6, 128.7, 128.0, 128.0, 119.2, 78.9, 74.5, 71.6, 51.7, 41.6, 34.2, 26.1, 22.9, 22.9, 10.2; high resolution mass spectrum (ES+) m/z343.1876 [ $(M + Na)^+$ ; calcd for C<sub>19</sub>H<sub>28</sub>O<sub>4</sub>Na 343.1886].

To a solution of (+)-58 (0.906 g, 2.83 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (9 mL) and cyclohexane (18 mL) at rt was added p-methoxybenzyl-2,2,2trichloroacetimidate (3.2 g, 4.0 equiv) followed by addition of TfOH  $(3 \mu L, 0.01 \text{ equiv})$ . The solution turned cloudy and light brown as the reaction progressed. After 30 min, hexane (20 mL) was added to precipitate solids, and the suspension was filtered. The filtrate was washed with sat. NaHCO<sub>3</sub> (25 mL), which was then back-extracted with  $CH_2Cl_2$  (3 × 25 mL). The combined organic layers were then dried over MgSO<sub>4</sub> and concentrated. The concentrate was diluted with 10% EtOAc/hexanes (25 mL) to facilitate further precipitation of solids and then filtered. The filtrate was then concentrated and purified by flash chromatography (3% EtOAc/hexanes to 5% EtOAc/hexanes) to provide PMB ether (+)-21 (0.910 g, 73% yield) as a light yellow oil:  $[\alpha]_{D}^{20}$  +43.5 (c 1.6, CHCl<sub>3</sub>); IR (neat) 2962, 2875, 2360, 2340, 1722, 1613, 1513, 1248, 1173, 1064 cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$ 7.44-7.28 (m, 5H), 7.21 (d, J = 8.5 Hz, 2H), 7.13 (d, J = 16.0 Hz, 1H), 6.87 (d, J = 8.5 Hz, 2H), 5.83 (d, J = 16.0 Hz, 1H), 4.62 (dab, J = 11.6 Hz, 1H), 4.46 (dab, J = 10.9 Hz, 1H), 4.35 (dd, J = 2.4, 11.8 Hz, 2H), 3.80 (s, 3H), 3.75 (s, 3H), 3.66-3.56 (m, 1H), 3.50 (d, J = 9.7, 1H), 1.72-1.58 (m, 3H), 1.56–1.48 (m, 1H), 1.12 (s, 3H), 1.12 (s, 3H), 0.93 (dd, J = 7.4, 7.4 Hz, 3H);  ${}^{13}$ C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  167.5, 159.2, 156.5, 139.2, 131.2, 129.2, 128.5, 127.8, 127.6, 118.6, 113.9, 83.0, 76.9, 74.9, 70.0, 55.4, 51.6, 42.7, 36.5, 26.2, 23.4, 23.4, 9.1; high resolution mass spectrum (ES+) m/z 440.2568 [(M + )<sup>+</sup>; calcd for C<sub>27</sub>H<sub>36</sub>O<sub>5</sub> 440.2563].

**Epoxy Alcohol (+)-22.** A solution of (+)-21 (1.76 g, 4.0 mmol) in  $CH_2Cl_2$  (20 mL) was cooled to -78 °C, and DIBAL-H (1.06 mL, 1.0 M in hexanes, 2.1 equiv) was slowly added. After 5 min, the reaction was quenched with MeOH (5 mL) followed by addition of sat. Rochelle's salt (10 mL). The solution was stirred for 2.5 h, and the layers were

separated. The aqueous layer was extracted with  $CH_2Cl_2$  (3 × 20 mL), and the combined organic layers were dried over MgSO4 and then concentrated. Flash chromatography (25% EtOAc/hexanes) provided allylic alcohol (+)-**59** (1.56 g, 95%) as a colorless oil:  $[\alpha]_{D}^{20}$  +65.0 (c 1.0, CHCl<sub>3</sub>); IR (neat) 3425, 2961, 2931, 2873, 1613, 1513, 1464, 1248, 1087, 1063, 1035 cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ 7.41-7.26 (m, 5H), 7.20 (d, J = 8.5 Hz, 2H), 6.85 (d, J = 8.6 Hz, 2H), 5.77 (d, J = 15.8 Hz, 1H), 5.60 (ddd, J = 15.8, 5.9, 5.9 Hz, 1H), 4.60 (dab, J = 11.6 Hz, 1H), 4.46 (dab, J = 10.8 Hz, 1H), 4.34 (dd, J = 2.6, 10.9 Hz, 2H), 4.10 (dd, J = 5.8, 5.8 Hz, 2H), 3.79 (s, 3H), 3.58 (ddd, J = 6.5, 4.2, 2.3 Hz, 1H), 3.40 (dd, J = 10.0, 1.5 Hz, 1H), 1.74–1.53 (m, 3H), 1.48 (ddd, J = 14.5, 10.0, 2.2 Hz, 1H), 1.07 (s, 3H), 1.06 (s, 3H), 0.90 (dd, J = 7.5, 7.5 Hz, 3H); <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>) δ 159.2, 140.6, 139.3, 131.6, 129.2, 128.6, 127.9, 127.7, 126.5, 113.9, 83.6, 77.2, 74.9, 70.0, 64.4, 55.5, 41.7, 36.5, 26.3, 24.5, 23.8, 9.3; high resolution mass spectrum (ES+) m/ z 435.2514 [(M + Na)<sup>+</sup>; calcd for C<sub>26</sub>H<sub>36</sub>O<sub>4</sub>Na 435.2512].

To freshly activated 3 Å molecular sieves (0.2 g) in CH<sub>2</sub>Cl<sub>2</sub> (2 mL) was added (+)-DIPT (31.8  $\mu$ L, 0.12 equiv). The solution was cooled to -20 °C, and Ti(O-*i*-Pr)<sub>4</sub> (37.3  $\mu$ L, 0.1 equiv) was added followed by *t*-BuOOH (0.687 mL, 5.5 M in decane, 3.0 equiv). The reaction was stirred for 30 min, and then allylic alcohol (+)-59 (0.521 g, 1.26 mmol) dissolved in CH<sub>2</sub>Cl<sub>2</sub> (1.5 mL) was added via syringe. The flask and syringe were rinsed with  $CH_2Cl_2$  (2 × 0.8 mL) into the reaction flask. After 2 h, 10% aq. citric acid (10 mL) was added, and the reaction warmed to rt. After 1 h at rt, the reaction was filtered through Celite, and the Celite washed with  $CH_2Cl_2$  (10 mL). The layers were separated, and the aqueous layer extracted with  $CH_2Cl_2$  (3 × 5 mL). The combined organic layers were dried over MgSO4 and concentrated. Flash chromatography (25% EtOAc/hexanes) provided epoxy alcohol (+)-22 (0.524 g, 97% yield, dr > 20:1) as a colorless oil:  $[\alpha]_{D}^{20} + 56.9$ (c 0.8, CHCl<sub>3</sub>); IR (neat) 3435, 2964, 2932, 2874, 1612, 1514, 1455, 1248, 1088, 1064, 1034 cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz, CDCl<sub>2</sub>)  $\delta$  7.40–7.26 (m, 5H), 7.19 (d, J = 8.6 Hz, 2H), 6.85 (d, J = 8.6 Hz, 2H), 4.62 (dab, J = 11.6 Hz, 1H), 4.46 (dab, J = 11.0 Hz, 1H) 4.35 (dd, J = 11.4, 11.8 Hz, 2H), 3.85–3.80 (m, 1H), 3.79 (s, 3H), 3.63–3.52 (m, 2H), 3.52 (dd, J = 10.0, 1.4 Hz, 1H), 3.05 (dd, J = 4.8, 2.6 Hz, 1H), 2.94 (d, J = 2.4 Hz, 1H), 1.82–1.50 (m, 4H), 0.93 (s, 3H), 0.93 (dd, J = 7.6, 7.6 Hz, 3H), 0.87 (s, 3H);  $^{13}\text{C}$  NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  159.2, 139.2, 131.4, 128.9, 128.6, 128.0, 127.7, 113.9, 82.4, 77.0, 74.6, 70.0, 62.2, 61.4, 55.6, 55.5, 39.2, 36.2, 26.3, 20.4, 19.0, 9.2; high resolution mass spectrum (ES+) m/z451.2444 [ $(M + Na)^+$ ; calcd for C<sub>26</sub>H<sub>36</sub>O<sub>5</sub>Na 451.2460].

Ester (+)-9. To a 0 °C solution of (+)-22 (0.233 g, 0.545 mmol) in DMSO (6 mL) and CH<sub>2</sub>Cl<sub>2</sub> (7 mL) was added Et<sub>3</sub>N (0.76 mL, 10 equiv) followed by SO3 pyridine (0.347 g, 4 equiv). After 1.5 h, NaHCO<sub>3</sub> (4 mL) was added, the layers separated, and the aqueous layer was extracted  $3 \times 10$  mL of Et<sub>2</sub>O. The combined organic layers were then washed with 1 M NaHSO<sub>4</sub>, sat. NaHCO<sub>3</sub>, and brine (10 mL each). The organic layer was then dried over MgSO<sub>4</sub> and concentrated. Flash chromatography (10% EtOAc/hexanes) provided aldehyde (+)-60 (1.43 g, 99% yield) as a colorless oil:  $[\alpha]_D^{20}$  +115.0 (c 1.2, CHCl<sub>3</sub>); IR (neat) 2964, 2931, 2876, 1728, 1613, 1514, 1464, 1248, 1090, 1064, 1035 cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  9.01 (d, J = 6.1 Hz, 1H), 7.44–7.27 (m, 5H), 7.19 (d, J = 8.4 Hz, 2H), 6.86 (d, J = 8.5 Hz, 2H), 4.64 (dab, J = 11.7 Hz, 1H), 4.43 (dab, J = 11.0 Hz, 1H), 4.38 (dab, J = 11.2 Hz, 1H), 4.35 (dab, J = 11.6 Hz, 1H), 3.80 (s, 3H), 3.64-3.58 (m, 1H), 3.56 (d, J = 9.8 Hz, 1H), 3.27 (dd, J = 6.1, 1.8 Hz, 1H), 3.21 (d, J = 1.8, 1H), 1.83–1.49 (m, 4H), 0.95 (s, 3H), 0.93 (dd, J = 7.5, 7.5 Hz, 3H), 0.88 (s, 3H); <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>) δ 199.1, 159.4, 139.1, 131.1, 129.2, 128.6, 127.9, 127.8, 114.0, 82.1, 76.9, 74.8, 70.0, 62.0, 56.9, 55.5, 39.6, 36.2, 26.2, 20.3, 19.1, 9.1; high resolution mass spectrum (ES+) m/ z 449.2305 [(M + Na)<sup>+</sup>; calcd for C<sub>26</sub>H<sub>34</sub>O<sub>5</sub>Na 449.2304].

A solution of aldehyde (+)-60 (0.445 g, 1.04 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (5 mL) was cooled to 0 °C, and a solution of carbomethoxy triphenylphosphonium ylide (23) in CH<sub>2</sub>Cl<sub>2</sub> (5 mL) was added over 1 min. The reaction was allowed to warm to rt, and after 30 min, the solvent was evaporated. Flash chromatography (8% EtOAc/hexanes) provided ester (+)-9 (0.453 g, 90% yield) as a light yellow oil:  $[\alpha]_{D}^{20}$  +48.7 (*c* 1.4, CHCl<sub>3</sub>); IR (neat) 2962, 2876, 1725, 1612, 1513, 1463, 1304, 1249, 1172, 1090, 1064 cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  7.41–7.26 (m, SH), 7.17 (d, *J* = 8.5 Hz, 2H), 6.85 (d, *J* = 8.5 Hz, 2H),

6.66 (dd, *J* = 15.7, 7.0 Hz, 1H), 6.05 (d, *J* = 15.7 Hz, 1H), 4.62 (dab, *J* = 11.6 Hz, 1H), 4.43 (dab, *J* = 10.9 Hz, 1H), 4.37 (dab, *J* = 11.2 Hz, 1H), 4.34 (dab, *J* = 11.8 Hz, 1H), 3.80 (s, 3H), 3.74 (s, 3H), 3.63–3.56 (m, 1H), 3.53 (d, *J* = 9.2 Hz, 1H), 3.35 (dd, *J* = 6.9, 1.7 Hz, 1H), 2.88 (d, *J* = 1.9 Hz, 1H), 1.81–1.50 (m, 4H), 0.94 (s, 3H), 0.93 (dd, *J* = 7.4, 7.4 Hz, 3H), 0.86 (s, 3H); <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  166.4, 159.3, 145.6, 139.2, 131.3, 129.0, 128.6, 127.9, 127.7, 123.2, 114.0, 82.1, 76.9, 74.6, 70.0, 67.0, 55.5, 53.8, 51.9, 39.9, 36.1, 26.3, 20.7, 18.6, 9.2; high resolution mass spectrum (ES+) *m*/*z* 505.2553 [(M + Na)<sup>+</sup>; calcd for C<sub>29</sub>H<sub>38</sub>O<sub>6</sub>Na 505.2566].

TBS Ether (+)-24. To freshly distilled dioxane (2 mL) was added  $Pd_2(dba)_3$ ·CHCl<sub>3</sub> (46 mg, 0.05 equiv) and *n*-Bu<sub>3</sub>P (13  $\mu$ L, 0.06 equiv). Next, a solution of HCO<sub>2</sub>H (0.20 mL, 6.0 equiv) and Et<sub>3</sub>N (0.25 mL, 2 equiv) in dioxane (1 mL) was added. This mixture was stirred at rt for 10 min, and then a solution of epoxide (+)-9 (0.452 g, 0.937 mmol) in dioxane (1.5 mL) was added via cannula. The flask and cannula were rinsed with dioxane  $(2 \times 1 \text{ mL})$  into the reaction flask. After 4.5 h, the reaction mixture was filtered through a 1 cm plug of silica gel, and the silica gel washed with CH<sub>2</sub>Cl<sub>2</sub> (20 mL). The filtrate was concentrated and subjected to flash chromatography (5% EtOAc/hexanes to 10% EtOAc/hexanes to 15% EtOAc/hexanes) to provide alcohol (+)-61 (0.421 g, 93% yield) as a colorless oil:  $[\alpha]_{D}^{20}$  +95.5 (c 0.7, CHCl<sub>3</sub>); IR (neat) 3464, 2963, 2875, 1722, 1613, 1514, 1249, 1173, 1063, 1036 cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  7.40–7.26 (m, 5H), 7.13 (d, J = 8.2 Hz, 2H), 7.08 (ddd, J = 14.8, 7.2, 7.2 Hz, 1H), 6.83 (d, J = 8.3 Hz, 2H), 5.90 (d, J = 15.6 Hz, 1H), 4.64 (dab, J = 11.7 Hz, 1H), 4.44 (dab, J = 10.7 Hz, 1H) 4.34 (dab, J = 12.0 Hz, 1H), 4.34 (dab, J = 12.0 Hz, 1H), 4.32 (dab, J = 11.1 Hz, 1H), 4.16 (s, 1H), 3.79 (s, 3H), 3.71 (s, 3H), 3.59–3.56 (m, 1H), 3.54 (d, J = 9.2 Hz, 1H), 2.25 (m, 2H), 1.91–1.67 (m, 3H), 1.62 (ddd, J = 14.1, 7.1, 7.1 Hz, 1H), 1.08 (s, 3H), 0.93 (dd, J = 7.4, 7.4 Hz, 3H), 0.83 (s, 3H); <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>) δ 167.1, 159.5, 148.0, 139.1, 130.4, 129.2, 128.6, 128.0, 127.8, 122.6, 114.1, 86.7, 77.1, 75.3, 75.0, 70.1, 55.5, 51.5, 41.5, 35.8, 35.1, 26.3, 23.3, 20.4, 9.2; high resolution mass spectrum (ES+) m/z 485.2888 [(M + H)<sup>+</sup>; calcd for C<sub>29</sub>H<sub>41</sub>O<sub>6</sub> 485.2903].

To a 0 °C solution of alcohol (+)-61 (0.789 g, 1.63 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (16.3 mL) was added 2,6-lutidine (0.38 mL, 2.0 equiv) followed by dropwise addition of TBSOTf (0.45 mL, 1.2 equiv). After 20 min, sat. NaHCO<sub>3</sub> (10 mL) was added, and the layers separated. The aqueous layer was extracted  $CH_2Cl_2$  (3 × 15 mL), and the combined organic layers were dried over MgSO4 and concentrated. Flash chromatography (5% EtOAc/hexanes) provided TBS ester (+)-24 (0.965 g, 99% yield) as a colorless oil:  $[\alpha]_{D}^{20}$  +42.1 (c 0.9, CHCl<sub>3</sub>); IR (neat) 2956, 2931, 2881, 2855, 1725, 1513, 1463, 1249, 1170, 1073 cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz,  $CDCl_3$ )  $\delta$  7.37–7.26 (m, 5H), 7.20 (d, J = 8.5 Hz, 2H), 7.06 (ddd, *J* = 15.7, 8.3, 6.4 Hz, 1H), 6.86 (d, *J* = 8.5 Hz, 2H), 5.76 (d, *J* = 15.7 Hz, 1H), 4.64 (dab, J = 11.5 Hz, 1H), 4.47 (dab, J = 10.9 Hz, 1H), 4.33 (dab, J = 10.8 Hz, 1H), 4.32 (dab, J = 11.7 Hz, 1H), 3.80 (s, 3H), 3.73 (s, 3H), 3.65 (dd, *J* = 6.4, 4.0 Hz, 1H), 3.60–3.55 (m, *J* = 9.9, 5.5 Hz, 1H), 3.52 (dd, J = 6.5, 5.0 Hz, 1H), 2.51 (ddd, J = 14.9, 4.2, 4.2 Hz, 1H), 2.36 (ddd, J = 15.0, 7.5, 7.5 Hz, 1H), 1.74 - 1.64 (m, 1H), 1.65 - 1.58 (m, 1H), 1.56(dd, J = 5.9, 5.9 Hz, 2H), 0.95 (s, 3H), 0.92 (dd, J = 7.5, 7.5 Hz, 3H), 0.90 (s, 9H), 0.89 (s, 3H), 0.05 (s, 3H), 0.03 (s, 3H); <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>) δ 167.1, 159.2, 148.6, 139.2, 131.7, 129.0, 128.6, 127.8, 127.7, 122.2, 113.9, 81.3, 77.3, 76.6, 75.0, 70.3, 55.5, 51.5, 44.7, 36.2, 36.1, 26.3, 26.2, 21.0, 20.1, 18.6, 9.2, -3.2, -3.8; high resolution mass spectrum (ES +) m/z 621.3605 [(M + Na)<sup>+</sup>; calcd for C<sub>35</sub>H<sub>54</sub>O<sub>6</sub>SiNa 621.3588]

**Epoxide (+)-25.** A solution of (+)-24 (1.43 g, 2.38 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (12 mL) was cooled to -78 °C, and DIBAL-H (1.06 mL, 1.0 M in hexanes, 2.1 equiv) was added slowly over 5 min. After 2 min, the reaction was quenched with methanol (1 mL) followed by addition of sat. Rochelle's salt (10 mL). The solution was stirred for 2 h, and the layers were separated. The aqueous layer was extracted with CH<sub>2</sub>Cl<sub>2</sub> (3 × 25 mL), and the combined organic layers were dried over MgSO<sub>4</sub> and then concentrated. Flash chromatography (5% EtOAc/hexanes) provided allylic alcohol (+)-62 (1.32 g, 97%) as a colorless oil:  $[\alpha]_{D}^{20}$  +47.7 (*c* 0.7, CHCl<sub>3</sub>); IR (neat) 3444, 2958, 2930, 2880, 2855, 1613, 1514, 1463, 1249, 1070 cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  7.41–7.27 (m, 5H), 7.21 (d, *J* = 8.5 Hz, 2H), 6.86 (d, *J* = 8.5 Hz, 2H), 5.74 (ddd, *J* = 15.2, 7.2 Hz, 1H), 5.55 (ddd, *J* = 15.4, 5.8, 5.8 Hz, 1H), 4.64 (dab, *J* =

11.4 Hz, 1H), 4.50 (dab, J = 10.9 Hz, 1H), 4.35 (dab, J = 10.8 Hz, 1H), 4.34 (dab, J = 11.2 Hz, 1H), 4.03–3.91 (m, 2H), 3.80 (s, 3H), 3.64–3.59 (m, 1H), 3.58 (dd, J = 9.7, 1.3 Hz, 1H), 3.55 (dd, J = 5.3, 5.3 Hz, 1H), 2.41 (ddd, J = 14.5, 5.4 Hz, 1H), 2.19 (ddd, J = 14.0, 6.7 Hz, 1H), 1.75– 1.64 (m, 1H), 1.64–1.49 (m, 3H), 1.47 (dd, J = 6.0, 6.0 Hz, 1H), 0.98 (s, 3H), 0.92 (dd, J = 7.5, 7.5 Hz, 3H), 0.90 (s, 9H), 0.88 (s, 3H), 0.04 (s, 6H); <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  159.2, 139.1, 131.8, 131.4, 130.7, 129.0, 128.6, 128.1, 127.7, 113.9, 80.9, 77.3, 75.0, 70.5, 63.8, 55.5, 45.0, 36.4, 36.0, 26.3, 26.3, 21.0, 19.9, 18.6, 9.2, -3.1, -3.9; high resolution mass spectrum (ES+) m/z 593.3651 [(M + Na)<sup>+</sup>; calcd for C<sub>34</sub>H<sub>54</sub>O<sub>5</sub>SiNa 593.3639].

To freshly activated 3 Å molecular sieves (0.4 g) in CH<sub>2</sub>Cl<sub>2</sub> (4.6 mL) was added (–)-DIPT (58  $\mu$ L, 0.12 equiv). The solution was cooled to -20 °C, and Ti(Oi-Pr)<sub>4</sub> (68  $\mu$ L, 0.1 equiv) was added followed by t-BuOOH (1.26 mL, 5.5 M in decane, 3.0 equiv). The reaction was stirred for 30 min, and then allylic alcohol (+)-62 (1.32 g, 2.31 mmol) dissolved in CH<sub>2</sub>Cl<sub>2</sub> (2 mL) was added via cannula. The flask and cannula were rinsed with  $CH_2Cl_2$  (2 × 1.5 mL) into the reaction flask. After 4 h, 10% aq. citric acid (10 mL) was added, and the reaction warmed to rt. After 2 h, the layers were separated, and the aqueous layer extracted with  $CH_2Cl_2$  (3 × 25 mL). The combined organic layers were dried over MgSO<sub>4</sub> and concentrated. Flash chromatography (10% EtOAc/hexanes to 15% EtOAc/hexanes) provided epoxide (+)-25 (1.24 g, 92% yield, dr > 20:1) as a colorless oil:  $[\alpha]_D^{20}$  +72.4 (c 1.9, CHCl<sub>3</sub>); IR (neat) 3444, 2957, 2930, 2881, 2856, 1613, 1514, 1463, 1249, 1071 cm<sup>-1</sup>; <sup>1</sup>H NMR  $(500 \text{ MHz}, \text{CDCl}_3) \delta 7.39 - 7.27 \text{ (m, 5H)}, 7.22 \text{ (d, } J = 8.5 \text{ Hz}, 2\text{H}), 6.86$ (d, J = 8.6 Hz, 2H), 4.61 (dab, J = 11.3 Hz, 1H), 4.50 (dab, J = 10.9 Hz, 1H), 4.37 (dab, J = 10.8 Hz, 1H), 4.31 (dab, J = 11.3 Hz, 1H), 3.80 (s, 3H), 3.78-3.71 (m, 2H), 3.62-3.51 (m, 2H), 3.49 (dd, J = 9.3, 2.0 Hz, 1H), 3.10 (ddd, *J* = 6.8, 4.7, 2.1 Hz, 1H), 2.70 (dd, *J* = 4.3, 2.6 Hz, 1H), 1.80-1.64 (m, 3H), 1.64-1.50 (m, 3H), 0.98 (s, 3H), 0.92 (s, 12H), 0.92 (dd, J = 7.4, 7.4 Hz, 3H), 0.11 (s, 3H), 0.10 (s, 3H); <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>) δ 159.2, 139.3, 131.7, 129.0, 128.5, 127.9, 127.6, 113.9, 81.4, 77.3, 75.1, 75.0, 70.0, 61.7, 59.9, 55.5, 53.7, 44.2, 35.9, 35.1, 26.4, 26.2, 20.9, 20.0, 18.6, 9.2, -3.5, -3.6; high resolution mass spectrum (ES +) m/z 609.3596 [(M + Na)<sup>+</sup>; calcd for C<sub>34</sub>H<sub>54</sub>O<sub>6</sub>SiNa 609.3588].

Alcohol (+)-26. Epoxy alcohol (+)-25 was dissolved in  $CH_3CN$  (12 mL), and then TEMPO (15 mg, 0.08 equiv) was added followed by pH 7 buffer (12 mL). Next, NaClO<sub>2</sub> (0.42 g, 2.5 equiv) was added in one portion followed by dropwise addition of NaOCl (0.32 mL, 5 wt % solution, 0.2 equiv). After 1.5 h, anhydrous Na<sub>2</sub>SO<sub>3</sub> (0.49 g, 3.2 equiv) was added, and the reaction stirred for 30 min, upon which the solution turned from orange to colorless. The reaction was acidified to pH 4 with 10% aq. citric acid solution and then extracted with EtOAc (3 × 25 mL). The combined organic layers were dried over MgSO<sub>4</sub> and concentrated.

The unpurified acid was then dissolved in  $Et_2O$  (24 mL) and cooled to 0 °C. A solution of CH2N2 in Et2O was then added dropwise until gas evolution ceased, and the reaction turned light yellow. Argon was bubbled through the reaction mixture for 15 min to remove any excess CH<sub>2</sub>N<sub>2</sub>, and then the reaction was concentrated. Flash chromatography (10% EtOAc/hexanes) provided methyl ester (+)-63 (0.557 g, 76% yield, 2 steps) as a colorless oil:  $[\alpha]_D^{20}$  +86.8 (c 1.0, CHCl<sub>3</sub>); IR (neat) 2957, 2931, 2882, 2857, 1754, 1513, 1458, 1249, 1065 cm<sup>-1</sup>; <sup>1</sup>H NMR  $(500 \text{ MHz}, \text{CDCl}_3) \delta 7.41 - 7.27 \text{ (m, 5H)}, 7.21 \text{ (d, } J = 8.6 \text{ Hz}, 2\text{H}), 6.86$ (d, J = 8.7 Hz, 2H), 4.62 (dab, J = 11.5 Hz, 1H), 4.50 (dab, J = 10.9 Hz, 1H), 4.36 (dab, J = 10.9 Hz, 1H), 4.30 (dab, J = 11.4 Hz, 1H), 3.80 (s, 3H), 3.75 (s, 3H), 3.75-3.70 (m, 1H), 3.60-3.53 (m, 1H), 3.49-3.42 (m, 1H), 3.31 (ddd, J = 6.5, 4.5, 1.8 Hz, 1H), 3.04 (d, J = 1.8 Hz, 1H), 1.80 (ddd, J = 14.5, 7.7, 4.5 Hz, 1H), 1.71–1.64 (m, 2H), 1.64–1.56 (m, 1H), 1.54 (dd, *J* = 7.6, 4.5 Hz, 2H), 0.96 (s, 3H), 0.93–0.90 (m, 15H), 0.12 (s, 3H), 0.09 (s, 3H); <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>) δ 169.9, 159.2, 139.3, 131.6, 129.0, 128.6, 127.8, 127.7, 113.9, 81.4, 77.1, 75.1, 74.9, 70.0, 56.6, 55.5, 54.7, 52.5, 44.2, 35.9, 34.8, 26.4, 26.1, 21.0, 20.0, 18.6, 9.1, -3.6, -3.6; high resolution mass spectrum (ES+) m/z 637.3521  $[(M + Na)^+; calcd for C_{35}H_{54}O_7SiNa 637.3537].$ 

A solution of (+)-63 (1.32 g, 2.14 mmol) in  $CH_2Cl_2$  (22 mL) and pH 7 buffer (5.4 mL) was cooled to 0 °C, and DDQ (0.542 g, 1.1 equiv) was added in three portions over 1 min. After 45 min, the reaction was diluted with  $CH_2Cl_2$  (20 mL) and filtered through Celite. The Celite was then washed with sat. NaHCO<sub>3</sub> (10 mL). The layers were separated,

and the aqueous layer extracted with  $CH_2Cl_2$  (3 × 20 mL). The combined organic layers were then dried with MgSO4 and concentrated. Flash chromatography (5% EtOAc/hexanes until the anisaldehyde eluted then 10% EtOAc/hexanes) provided alcohol (+)-26 (0.998g, 94% yield) as a light yellow oil:  $[\alpha]_D^{20}$  +54.3 (c 0.7, CHCl<sub>3</sub>); IR (neat) 3494, 2957, 2930, 2880, 2855, 1755, 1452, 1254, 1204, 1058 cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  7.42–7.23 (m, 5H), 4.63 (dab, J = 11.6 Hz, 1H), 4.53 (dab, J = 11.6 Hz, 1H) 3.91 (dd, J = 8.8, 3.2 Hz, 1H), 3.84 (s, 1H), 3.82–3.76 (m, 1H), 3.77 (s, 3H), 3.69 (ddd, J = 11.4, 5.9, 5.9 Hz, 1H), 3.32 (ddd, J = 7.9, 3.3, 1.9 Hz, 1H), 3.23 (d, J = 1.8 Hz, 1H), 1.98 (ddd, J = 14.8, 8.2, 3.4 Hz, 1H), 1.69–1.52 (m, 3H), 1.50–1.45 (m, 2H), 0.95 (s, 3H), 0.94–0.91 (m, 12H), 0.72 (s, 3H), 0.16 (s, 3H), 0.15 (s, 3H); <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>) δ 169.6, 139.5, 128.5, 127.9, 127.6, 79.5, 78.2, 72.2, 71.4, 56.6 54.4, 52.7, 40.9, 36.1, 35.2, 27.5, 26.3, 22.5, 20.0, 18.5, 9.9, -3.7, -4.1; high resolution mass spectrum (ES+) m/z495.3125 [(M + H)<sup>+</sup>; calcd for  $C_{27}H_{47}O_6Si$  495.3142].

Alcohol (+)-27. To a solution of (+)-26 (0.998 g, 2.02 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (40 mL) was added camphorsulfonic acid (94 mg, 0.2 equiv). After stirring for 5 h at rt, sat. NaHCO<sub>3</sub> (20 mL) was added, and the reaction mixture stirred for 10 min. The layers were separated, and the aqueous layer extracted with  $CH_2Cl_2$  (4 × 15 mL). The combined organic layers were then dried over MgSO4 and concentrated. Flash chromatography (5% EtOAc/hexanes) provided alcohol (+)-27 (0.917g, 92%) as a colorless oil:  $[\alpha]_D^{20} + 38.1$  (*c* 1.6, CHCl<sub>3</sub>); IR (neat) 3461, 2956, 2930, 2857, 1741, 1471, 1437, 1360, 1256, 1080 cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  7.44–7.17 (m, 5H), 4.54 (dd, J = 41.3, 11.5 Hz, 2H), 4.23 (dd, J = 5.8, 4.0 Hz, 1H), 4.07 (dd, J = 9.9, 3.5 Hz, 1H), 3.78 (s, 3H), 3.80-3.75 (m, 1H), 3.63 (dd, J = 3.8, 3.8 Hz, 1H), 3.56-3.48 (m, 1H), 2.87 (d, J = 6.2 Hz, 1H), 2.31 (dd, J = 13.0, 13.0 Hz, 1H), 2.06 (ddd, *J* = 13.4, 10.0, 3.1 Hz, 1H), 1.71–1.48 (m, 3H), 1.36 (ddd, *J* = 13.6, 3.9, 3.9 Hz, 1H), 1.01 (s, 3H), 0.92 (dd, J = 7.5, 7.5 Hz, 3H), 0.90 (s, 9H), 0.86 (s, 3H), 0.05 (s, 3H), 0.03 (s, 3H); <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>) δ 173.2, 139.5, 128.5, 128.4, 128.0, 127.5, 77.9, 73.8, 73.7, 71.6, 67.6, 52.6, 37.3, 33.0, 30.1, 27.2, 26.5, 26.0, 21.5, 18.2, 9.6, -4.4, -4.9; high resolution mass spectrum (ES+) m/z 517.2970 [(M + Na)<sup>+</sup>; calcd for C<sub>27</sub>H<sub>46</sub>O<sub>6</sub>SiNa 517.2962].

Methyl Ether (+)-28. A solution of (+)-27 (0.236g, 0.478 mmol) in THF (5 mL) was cooled to 0 °C, and NaH (29 mg, 60% in mineral oil, 1.5 equiv) added in one portion. After 20 min, Me<sub>2</sub>SO<sub>4</sub> (59 µL, 1.3 equiv) was added dropwise, and the reaction allowed to warm to rt. After 1.5 h, the reaction was quenched with sat. NaHCO<sub>3</sub> (5 mL). The layers were separated, and the aqueous layer extracted with  $CH_2Cl_2$  (5 × 15 mL). The combined organic layers were then dried over MgSO4 and concentrated. Flash chromatography (3% EtOAc/hexanes to 5% EtOAc/hexanes) provided methyl ether (+)-28 (0.229g, 94% yield) as a colorless oil:  $[\alpha]_{D}^{20}$  +20.0 (c 2.8, CHCl<sub>3</sub>); IR (neat) 2956, 2931, 2857, 1751, 1462, 1359, 1256, 1195, 1126, 1083, 1006 cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  7.44–7.14 (m, 5H), 4.57 (dab, J = 11.5 Hz, 1H), 4.51 (dab, *J* = 11.4 Hz, 1H), 4.08 (ddd, *J* = 7.1, 4.2, 4.2 Hz, 1H), 3.89 (d, *J* = 6.4 Hz, 1H), 3.74 (s, 3H), 3.70 (dd, J = 11.3, 1.5 Hz, 1H), 3.59 (dd, J = 6.5, 3.4 Hz, 1H), 3.50–3.44 (m, 1H), 3.39 (s, 3H), 2.08 (dd, J = 12.7, 12.7 Hz, 1H), 1.98 (ddd, J = 13.7, 7.6, 3.5 Hz, 1H), 1.70–1.48 (m, 4H), 1.00 (s, 3H), 0.92 (dd, J = 7.6, 7.6 Hz, 3H), 0.90 (s, 9H), 0.85 (s, 3H),  $0.05 (s, 3H), 0.04 (s, 3H); {}^{13}C NMR (125 MHz, CDCl_3) \delta 171.8, 139.8,$ 128.4, 127.8, 127.4, 83.2, 78.0, 77.9, 73.5, 71.6, 68.4, 58.8, 52.1, 37.9, 33.7, 30.7, 27.4, 26.0, 25.9, 19.2, 18.2, 9.6, -4.3, -4.9; high resolution mass spectrum (ES+) m/z 531.3139 [(M + Na)<sup>+</sup>; calcd for C<sub>28</sub>H<sub>48</sub>O<sub>6</sub>SiNa 531.3118].

**Tetrahydropyran** (+)-6. To a solution of (+)-28 (0.229 g, 0.450 mmol) in EtOAc (4.5 mL) was added 10% Pd/C (0.025 g). The reaction flask was purged with H<sub>2</sub>, and then a balloon of H<sub>2</sub> was attached to the flask. After 5 h at rt, the reaction mixture was filtered through a pad of Celite, and the Celite rinsed with CH<sub>2</sub>Cl<sub>2</sub> (10 mL). The reaction was concentrated, and then flash chromatography (10% EtOAc/hexanes) provided alcohol (+)-64 (0.182 g, 97%) as a colorless oil:  $[\alpha]_{20}^{20}$  +12.7 (*c* 3.6, CHCl<sub>3</sub>); IR (neat) 3557, 2955, 2930, 2857, 1730, 1463, 1285, 1254, 1127, 1082 cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  4.08 (d, *J* = 9.0 Hz, 1H), 4.04–4.00 (m, 1H), 3.81 (s, 3H), 3.59–3.53 (m, 2H), 3.53–3.46 (m, 1H), 3.41 (s, 3H), 2.92 (d, *J* = 4.1 Hz, 1H), 1.93 (ddd, *J* = 13.8, 4.2, 2.8 Hz, 1H), 1.74 (ddd, *J* = 13.8, 10.4, 5.7 Hz, 1H), 1.60 (dd, *J* = 12.3,

12.3 Hz, 1H), 1.53–1.34 (m, 3H), 0.94 (dd, J = 7.4, 7.4 Hz, 3H), 0.89 (s, 12H), 0.83 (s, 3H), 0.05 (s, 3H), 0.04 (s, 3H); <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  173.1, 81.0, 76.1, 72.6, 72.2, 68.8, 58.8, 52.6, 39.0, 36.5, 30.4, 30.2, 26.0, 23.8, 18.2, 14.2, 10.6, -4.1, -4.8; high resolution mass spectrum (ES+) m/z 441.2642 [(M + Na)<sup>+</sup>; calcd for C<sub>21</sub>H<sub>42</sub>O<sub>6</sub>SiNa 441.2649].

A solution of alcohol (+)-64 (0.187g, 0.446 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (4.5 mL) was cooled to 0 °C, and NaHCO<sub>3</sub> (56 mg, 1.5 equiv) was added followed by Dess-Martin periodinane (0.568 g, 3 equiv). After 1 h at 0 °C, H2O, sat. NaHCO3, and CH2Cl2 (5 mL each) were added. The solution was stirred until the organic layer went clear (ca. 30 min). The layers were separated, and the aqueous layer extracted with  $CH_2Cl_2$  (3 × 10 mL). The combined organic layers were then dried over MgSO<sub>4</sub> and concentrated. Flash chromatography (10% EtOAc/hexanes) provided tetrahydropyran (+)-6 (0.181g, 97% yield) as a colorless oil:  $\left[\alpha\right]_{D}^{20}$  +16.5 (c 1.7, CHCl<sub>3</sub>); IR (neat) 2954, 2934, 2886, 2858, 1754, 1722, 17.112, 1462, 1361, 1255, 1119, 1073 cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  4.10 (dd, J = 10.6, 2.5 Hz, 1H), 4.04 (ddd, J = 11.9, 4.5, 2.3 Hz, 1H), 3.80 (d, J = 4.5 Hz, 1H), 3.70 (s, 3H), 3.53 (dd, J = 2.7, 2.7 Hz 1H), 3.40 (s, 3H), 2.60–2.38 (m, 3H), 2.23 (dd, J = 14.5, 2.5 Hz, 1H), 1.96 (ddd, J = 14.1, 12.0, 2.4 Hz, 1H), 1.40 (ddd, J = 13.8, 3.1, 2.5 Hz, 1H), 1.04 (dd, J = 7.3, 7.3 Hz, 3H), 0.90 (s, 9H), 0.89 (s, 3H), 0.80 (s, 3H), 0.03 (s, 3H), 0.01 (s, 3H);  $^{13}$ C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  210.4, 171.7, 82.5, 77.4, 73.1, 69.8, 58.8, 52.2, 42.6, 38.1, 37.2, 30.3, 26.0, 25.1, 18.2, 17.9, 7.8, -4.2, -4.8; high resolution mass spectrum (ES+) m/z 439.2491 [(M + Na)<sup>+</sup>; calcd for C<sub>21</sub>H<sub>40</sub>O<sub>6</sub>SiNa 439.2492].

 $\beta$ -Hydroxy Ketone (+)-29. A solution of ketone (+)-6 (0.140 g, 0.336 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (3.4 mL) was cooled to -78 °C, and Cl<sub>2</sub>BPh  $(52 \,\mu\text{L}, 1.2 \text{ equiv})$  was added. After stirring for 20 min, *i*-Pr<sub>2</sub>NEt (88  $\mu$ L, 1.5 equiv) was added dropwise. The reaction was stirred for 1 h at -78°C, warmed to 0 °C over 10 min, and then stirred for 1 h at 0 °C. After cooling back to -78 °C, aldehyde 5 (0.237g, 1.45 equiv) was dissolved in CH<sub>2</sub>Cl<sub>2</sub> (1.5 mL) and added to the boron enolate dropwise over 15 min. After 1 h at -78 °C, the reaction was quenched with a 1:1 mixture of MeOH/pH 7 buffer (6 mL). After warming to 0 °C, the reaction was neutralized to pH 7 with pH 8 buffer and stirred for 1 h at 0 °C. The layers were separated, and the aqueous layer extracted with  $CH_2Cl_2$  (3 × 15 mL). The combined organic layers were then dried over MgSO<sub>4</sub> and concentrated. Flash chromatography (15% EtOAc/hexanes) provided  $\beta$ -hydroxy ketone (+)-29 (0.273 g, 90% yield, single diastereomer) as a colorless oil: [a]<sup>20</sup><sub>D</sub> +30.0 (c 0.9, CHCl<sub>3</sub>); IR (neat) 3467, 2952, 2896, 1731, 1593, 1463, 1253, 1064 cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ 6.88 (s, 1H), 5.21 (s, 2H), 5.15 (d, J = 2.3 Hz, 2H), 4.09 (dd, J = 11.1, 5.7 Hz, 1H), 4.07–4.00 (m, 2H), 3.94 (d, J = 6.1 Hz, 1H), 3.87 (s, 3H), 3.78– 3.70 (m, 4H), 3.72 (s, 3H), 3.64 (dd, J = 7.8, 3.8 Hz, 1H), 3.43 (d, J = 5.6 Hz, 1H), 3.40 (s, 3H), 3.06 (dd, J = 16.4, 9.4 Hz, 1H), 2.84 (dd, J = 14.2, 3.2 Hz, 1H), 2.75–2.63 (m, 1H), 2.61–2.50 (m, 2H), 2.16 (s, 3H), 1.95 (ddd, J = 13.8, 5.8, 3.9 Hz, 1H), 1.57 (ddd, J = 13.3, 7.8, 4.9 Hz, 1H), 1.19 (d, J = 7.1 Hz, 3H), 0.95 (s, 3H), 0.98–0.92 (m, 4H), 0.90 (s, 9H), 0.84 (s, 3H), 0.04 (s, 6H), 0.01 (s, 9H), 0.00 (s, 9H); <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>) δ 212.5, 171.6, 170.7, 157.8, 153.7, 136.6, 120.3, 119.1, 101.1, 93.9, 93.3, 82.4, 76.7, 73.1, 71.6, 70.1, 66.5, 66.5, 58.8, 53.2, 52.7, 52.2, 42.5, 38.1, 35.9, 30.1, 26.0, 24.9, 18.3, 18.2, 17.6, 11.8, 11.4, -1.2, -4.3, -4.8; high resolution mass spectrum (ES+) m/z 923.4782 [(M + Na)<sup>+</sup>; calcd for C44H80O13Si3Na 923.4805].

Acid (+)-30. A solution (+)-29 (0.116 g, 0.129 mmol) in THF (1.4 mL) and MeOH (0.46 mL) was cooled to -78 °C, and Et<sub>2</sub>BOMe (0.167 mL, 1 M in THF, 1.3 equiv) was added dropwise. After 25 min, NaBH<sub>4</sub> (10 mg, 2.0 equiv) was added, and the reaction was strirred at -78 °C. After 1 h, the reaction was warmed to 0 °C over 10 min and then stirred for 30 min. EtOAc (2 mL) was added followed by H<sub>2</sub>O (2 mL) and a 1:1 solution of MeOH/30% aq. H<sub>2</sub>O<sub>2</sub> (5 mL). After 1 h, the reaction was extracted with EtOAc (3 × 10 mL), and the combined organic layers were treated with solid Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub> to destroy any remaining peroxide. The organic layer was then filtered and washed with sat. Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub>, dried over MgSO<sub>4</sub>, and concentrated. Flash chromatography (15% EtOAc/hexanes) provided diol (+)-65 (0.109 g, 95% yield, *dr* > 20:1) as a colorless oil:  $[\alpha]_{20}^{20}$  +10.3 (*c* 1.0, CHCl<sub>3</sub>); IR (neat) 3500, 2953, 2896, 2857, 1734, 1593, 1253, 1157, 1115, 1065 cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  6.87 (s, 1H), 5.21 (s, 2H), 5.16 (s, 2H),

4.16–4.10 (m, 1H), 4.02 (dd, *J* = 7.5, 4.9 Hz, 1H), 3.97 (d, *J* = 7.9 Hz, 1H), 3.88 (s, 3H), 3.84 (dd, *J* = 12.9, 7.1 Hz, 1H), 3.78 (s, 3H), 3.76– 3.70 (m, 4H), 3.65 (s, 1H), 3.56–3.50 (m, 2H), 3.39 (s, 3H), 2.85–2.80 (m, 2H), 2.20 (s, 3H), 1.93 (ddd, *J* = 13.8, 4.0, 4.0 Hz, 1H), 1.83 (ddd, *J* = 14.2, 10.3, Hz, 1H), 1.72–1.63 (m, 1H), 1.51 (d, *J* = 7.0 Hz, 1H), 1.45 (d, *J* = 14.7 Hz, 1H), 0.98 (d, *J* = 6.9, 3H), 0.96–0.92 (m, 4H), 0.90 (s, 9H), 0.88 (s, 3H), 0.85 (s, 3H), 0.04 (s, 3H), 0.04 (s, 3H), 0.01 (s, 9H), 0.00 (s, 9H); <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  171.3, 169.9, 157.2, 153.1, 136.9, 120.2, 119.2, 100.6, 93.6, 93.1, 81.7, 80.9, 76.2, 75.5, 72.1, 71.4, 66.2, 66.2, 58.5, 52.2, 52.2, 41.9, 38.9, 35.8, 33.4, 29.8, 25.7, 24.0, 18.0, 17.9, 15.1, 11.7, 5.8, -1.5, -4.4, -5.0; high resolution mass spectrum (ES+) *m*/z 925.5000 [(M + Na)<sup>+</sup>; calcd for C<sub>44</sub>H<sub>82</sub>O<sub>13</sub>Si<sub>3</sub>Na 925.4961].

A solution of diol (+)-65 (20.6 mg, 0.023 mmol) was dissolved in MeOH (1.1 mL) and cooled to 0 °C. Next, H<sub>2</sub>O (8 µL, 20 equiv) was added followed by LiOH (11 mg, 20 equiv). The cold bath was removed, and the reaction allowed to warm to rt. After 28 h, because of bis-acid formation, the reaction was quenched by diluting with EtOAc and acidified to pH 5 with 5% aq. AcOH. Brine (2 mL) was added, and the reaction extracted with EtOAc  $(5 \times 3 \text{ mL})$ . The combined organic layers were then dried over MgSO4 and concentrated. Flash chromatography (0.1% AcOH in 30% EtOAc/hexanes to 0.2% AcOH in 60% EtOAc/ hexanes) provided acid (+)-30 (17 mg, 87% yield) as a colorless oil:  $[\alpha]_{D}^{20}$  +31.4 (c 1.1, CHCl<sub>3</sub>); IR (neat) 3467, 2953, 2898, 2858, 1720, 1592, 1250, 1107, 1065 cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  6.95 (s, 1H), 5.30 (dd, J = 20.0, 7.0 Hz, 2H), 5.27 (s, 2H), 4.33 (ddd, J = 12.1, 5.8, 2.5 Hz, 1H), 4.23 (dd, J = 10.9, 6.6 Hz, 1H), 4.15 (d, J = 7.3 Hz, 1H), 3.87-3.79 (m, 3H), 3.78-3.72 (m, 2H), 3.64-3.57 (m, 2H), 3.45 (s, 3H), 3.03 (dd, J = 16.5, 2.4 Hz, 1H), 2.91 (dd, J = 16.4, 12.2 Hz, 1H), 2.24–2.07 (m, 1H), 2.12 (s, 3H), 2.01 (dd, J = 10.6, 7.0 Hz, 1H), 1.97– 1.92 (m, 1H), 1.66 (ddd, J = 13.7, 6.8, 4.7 Hz, 1H), 1.54 (d, J = 15.0 Hz, 1H), 1.08 (d, J = 7.0 Hz, 3H), 0.97 (s, 3H), 0.99-0.94 (m, 4H), 0.91 (s, 9H), 0.88 (s, 3H), 0.06 (s, 6H), 0.01 (s, 9H), 0.00 (s, 9H); <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>) δ 172.3, 164.1, 160.1, 159.2, 141.9, 117.45, 109.0, 102.1, 94.4, 93.1, 82.2, 81.9, 79.0, 72.7, 71.9, 69.8, 66.8, 66.8, 58.7, 41.7, 38.7, 33.0, 30.6, 29.5, 26.0, 25.0, 18.3, 18.3, 18.2, 11.4, 9.7, 9.7, -1.2, -1.2, -4.2, -4.8; high resolution mass spectrum (ES+) m/z 879.4569  $[(M + Na)^+; calcd for C_{42}H_{76}O_{12}Si_3Na 879.4543].$ 

Silyl Ether (+)-4. A solution of (+)-30 (54 mg, 0.063 mmol) in acetone (3.2 mL) was cooled to 0 °C, and *i*-Pr<sub>2</sub>NEt (24  $\mu$ L, 2.2 equiv) was added followed by dropwise addition of isobutyl chloroformate (20  $\mu$ L, 2.4 equiv). After 45 min at 0 °C, NaN<sub>3</sub> (21 mg, 5 equiv) was dissolved in H<sub>2</sub>O (0.4 mL) and added to the reaction over 2 min. After 20 min, cold H<sub>2</sub>O (2 mL) was added, and the reaction extracted with cold EtOAc (3  $\times$  5 mL). The combined organic layers were dried thoroughly over MgSO<sub>4</sub> and concentrated. The residue was azeotroped with benzene  $(3 \times 5 \text{ mL})$  and placed on the vacuum pump for 30 min. The unpurified acyl azide was dissolved in toluene (3.2 mL), the reaction flask fitted with a reflux condenser, and then heated to 80 °C. After 45 min, 2-trimethylsilyl ethanol (0.181 mL, 20 equiv) was added through the top of the reflux condenser. After 2 h, the reaction was cooled to rt, and the solvent evaporated. Flash chromatography (10% EtOAc/ hexanes to 25% EtOAc/hexanes) provided N,O-aminal (+)-43 (45 mg, 74% yield) as a colorless oil:  $[\alpha]_D^{20}$  +8.4 (*c* 0.5, CHCl<sub>3</sub>); IR (neat) 3501, 3315, 2953, 2901, 2857, 1719, 1595, 1249, 1108, 1064 cm<sup>-1</sup>; <sup>1</sup>H NMR  $(500 \text{ MHz}, \text{CDCl}_3) \delta 6.94 (s, 1\text{H}), 5.37 (d, J = 10.1 \text{ Hz}, 1\text{H}), 5.31 (dab, J)$ = 7.0 Hz, 1H), 5.27 (dab, J = 6.7 Hz, 1H), 5.26 (s, 1H), 4.89 (dd, J = 10.0, 2.6 Hz, 1H), 4.31 (ddd, J = 11.6, 6.9. 2.4 Hz, 1H), 4.24–4.08 (m, 2H), 3.97 (d, J = 9.4 Hz, 1H), 3.88–3.72 (m, 4H), 3.65 (s, 1H), 3.62 (d, J = 10.4 Hz, 1H), 3.56 (dd, J = 4.9, 3.2 Hz, 1H), 3.36 (s, 3H), 3.10 (dd, J = 16.4, 2.1 Hz, 1H), 2.82 (dd, J = 16.4, 11.9 Hz, 1H), 2.45–2.28 (m, 1H), 2.12 (s, 3H), 1.91–1.80 (m, J = 12.5, 8.0 Hz, 2H), 1.53–1.37 (m, 2H), 1.11 (d, J = 6.9 Hz, 3H), 1.00 (s, 3H), 0.98-0.93 (m, 6H), 0.90 (s, 9H), 0.88 (s, 3H), 0.05 (s, 3H), 0.04 (s, 3H), 0.02 (s, 9H), 0.00 (s, 9H), 0.00 (s, 9H); <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  163.5, 159.9, 159.1, 157.2, 141.9, 117.4, 109.4, 102.2, 94.4, 93.1, 83.8, 83.6, 79.4, 77.4, 73.0, 72.7, 68.0, 66.8, 66.7, 63.7, 55.8, 43.4, 38.1, 32.9, 31.1, 29.7, 26.1, 26.0, 18.4, 18.3, 18.2, 17.8, 11.4, 10.2, -1.1, -1.2, -1.3, -4.3, -4.8; high resolution mass spectrum (ES+) m/z 994.5354 [(M + Na)<sup>+</sup>; calcd for C<sub>47</sub>H<sub>89</sub>NO<sub>12</sub>Si<sub>4</sub>Na 994.5360].

A solution of N,O-aminal (+)-43 (19.6 mg, 0.020 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (0.2 mL) was cooled to 0 °C, and 2,6-lutidine (9.5  $\mu$ L, 4.0 equiv) was added followed by dropwise addition of TBSOTf (10  $\mu$ L, 2.1 equiv). After 45 min, CH<sub>2</sub>Cl<sub>2</sub> (1 mL) and sat. NaHCO<sub>3</sub> (2 mL) added, and the reaction allowed to warm to rt. The layers were separated, and the aqueous layer extracted with  $CH_2Cl_2$  (3 × 5 mL). Preparative TLC  $(25\% \text{ EtOAc/hexanes}, 500 \,\mu\text{m} \text{ plate})$  provided silvl ether (+)-4 (20 mg, 91%) as a colorless oil:  $[\alpha]_D^{20}$  +36.0 (c 0.8, CHCl<sub>3</sub>); IR (neat) 2954, 2928, 2896, 2857, 1725, 1594, 1472, 1250, 1108, 1063  $\rm cm^{-1}; \ ^1H$  NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  6.93 (s, 1H), 5.45 (d, J = 9.7 Hz, 1H), 5.32 (dab, J = 7.0 Hz, 1H), 5.27 (dab, J = 6.7 Hz, 1H), 5.27 (s, 1H), 4.82 (d, J = 9.6 Hz, 1H), 4.23-4.07 (m, 3H), 4.00 (d, J = 8.7 Hz, 1H), 3.88-3.78 (m, 2H), 3.75 (dd, J = 8.7, 7.9 Hz, 2H), 3.58 (dd, J = 3.8, 3.8 Hz, 1H), 3.37 (s, 3H), 3.34 (d, J = 11.6 Hz, 1H), 3.09 (d, J = 15.3 Hz, 1H), 2.64 (dd, J = 16.4, 12.1 Hz, 1H), 2.32–2.22 (m, 1H), 2.12 (s, 3H), 2.03–1.92 (m, 1H), 1.87–1.77 (m, 1H), 1.62 (dd, *J* = 11.6, 11.6 Hz, 1H), 1.48 (ddd, *J* = 13.7, 8.2, 8.2 Hz, 1H), 1.07 (d, J = 6.7 Hz, 3H), 0.97 (s, 3H), 0.9–0.93 (m, 6H), 0.91 (s, 9H), 0.86 (s, 3H), 0.80 (s, 9H), 0.09 (s, 3H), 0.07 (s, 3H), 0.06 (s, 3H), 0.01 (s, 3H), 0.00 (s, 9H), 0.00 (s, 18H); <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>) δ 163.5, 159.8, 159.1, 157.2, 141.4, 117.2, 109.5, 102.3, 94.5, 93.1, 84.4, 79.9, 77.7, 77.2, 73.6, 68.9, 67.8, 66.8, 66.7, 63.6, 56.1, 39.9, 37.6, 32.9, 31.7, 29.9, 26.8, 26.2, 26.0, 18.4, 18.3, 18.3, 18.2, 17.8, 11.4, 9.1, -1.1, -1.2, -1.3, -3.3, -4.2, -4.7, -4.7; high resolution mass spectrum (ES+) m/z 1108.6233 [(M + Na)<sup>+</sup>; calcd for C53H103NO12Si5Na 1108.6225].

Amide (+)-32. A solution of acid (-)-3 (47 mg, 4 equiv) in CH<sub>2</sub>Cl<sub>2</sub> (1.5 mL) was cooled to 0 °C, and *i*-Pr<sub>2</sub>NEt (30  $\mu$ L, 1.1 equiv) was added followed by trimethylacetyl chloride  $(20 \,\mu\text{L}, 1.05 \text{ equiv})$ . After 30 min at 0 °C, sat. NH<sub>4</sub>Cl (3 mL) was added followed by enough H<sub>2</sub>O to dissolve any solids. The layers were separated, and the aqueous layer extracted with  $CH_2Cl_2$  (3 × 5 mL). The combined organic layers were then dried over MgSO<sub>4</sub> and concentrated. The unpurified mixed anhydride 31 was azeotroped with benzene  $(3 \times 3 \text{ mL})$ , placed on the vacuum pump for 30 min, and then dissolved in THF (0.4 mL). A solution of (+)-4 (41.5 mg, 0.038 mmol) in THF (0.4 mL) was cooled to -78 °C, and LiHMDS (0.152 mL, 0.5 M in THF, 2.0 equiv) was added dropwise over 1 min. The solution was stirred for 30 min at -78 °C, and then the solution of mixed anhydride 32 was added dropwise over 5 min. The reaction was stirred at -78 °C for 45 min and then warmed to -60 °C and stirred for 30 min. Sat. NH<sub>4</sub>Cl (2 mL) was added, and the reaction allowed to warm to rt. The reaction was extracted with  $CH_2Cl_2$  (4 × 5 mL). The combined organic layers were then dried over MgSO4 and evaporated. Preparative chromatography (20% EtOAc/hexanes, 1000  $\mu$ m plate) provided amide (+)-32 (41.2 mg, 79% yield) as a colorless oil:  $\left[\alpha\right]_{D}^{2\ell}$ +40.2 (c 0.4, CHCl<sub>3</sub>); IR (neat) 2953, 2896, 2857, 1726, 1591, 1467, 1250, 1110, 1063 cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ 6.92 (s, 1H), 5.66 (d, J = 4.6 Hz, 1H), 5.34–5.25 (m, 4H), 5.17 (d, J = 4.9 Hz, 1H), 4.77 (d, J = 11.5 Hz, 2H), 4.63 (dd, J = 23.1, 6.6 Hz, 2H), 4.33-4.27 (m, 2H), 4.20 (ddd, *J* = 12.1, 8.4, 2.0 Hz, 1H), 4.12 (ddd, *J* = 10.0, 2.9, 2.9 Hz, 1H), 3.90-3.78 (m, 2H), 3.78-3.73 (m, 2H), 3.65-3.59 (m, 1H), 3.59–3.49 (m, 3H), 3.34 (s, 3H), 3.28 (s, 3H), 3.14 (d, J = 9.6 Hz, 1H), 3.10 (dd, J = 16.5, 2.0 Hz, 1H), 2.93 (dd, J = 16.4, 12.6 Hz, 1H), 2.30 (dd, J = 14.7, 9.0 Hz, 1H), 2.22 (d, J = 12.7 Hz, 1H), 2.19 (s, 3H), 2.07–1.96 (m, 2H), 1.81 (dd, J = 14.0, 10.3 Hz, 1H), 1.75 (s, 3H), 1.69 (ddd, J =13.2, 10.8, 6.5 Hz, 1H), 1.61 (ddd, J = 13.6, 9.5, 3.2 Hz, 2H), 1.10 (d, J = 6.6 Hz, 3H), 1.10-1.04 (m, 2H), 1.00-0.93 (m, 4H), 0.90 (s, 12H), 0.87-0.80 (m, 2H), 0.83 (s, 3H), 0.78 (s, 9H), 0.05 (s, 15H), 0.04 (s, 3H), 0.01 (s, 9H), 0.00 (s, 9H), -0.02 (s, 3H), -0.05 (s, 9H); <sup>13</sup>C NMR  $(125 \text{ MHz}, \text{CDCl}_3) \delta 174.9, 163.7, 159.8, 159.1, 154.5, 142.9, 142.1,$ 117.5, 113.0, 109.5, 102.2, 95.3, 94.5, 93.1, 88.5, 81.3, 80.2, 75.7, 74.9, 73.0, 69.3, 66.8, 66.8, 66.3, 66.1, 58.3, 56.8, 40.7, 39.2, 39.1, 35.3, 30.5, 30.0, 28.7, 26.1, 26.0, 24.2, 23.1, 18.3, 18.3, 18.3, 18.2, 17.8, 13.6, 11.4, 9.1, -1.2, -1.2, -1.3, -1.4, -3.1, -4.0, -4.5, -4.7.; high resolution mass spectrum (ES+) m/z 1394.7849 [(M + Na)<sup>+</sup>; calcd for C<sub>67</sub>H<sub>129</sub>NO<sub>16</sub>Si<sub>6</sub>Na 1394.7825].

**Irciniastatin A (+)-1.** Compound (+)-**32** (9.8 mg, 7.1  $\mu$ mol) was dissolved in DMF (0.16 mL), and TASF (30 mg, 15 equiv) was added. The reaction was heated to 50 °C. After 48 h, the reaction was diluted with EtOAc, and sat. NH<sub>4</sub>Cl (1 mL) was added followed by enough H<sub>2</sub>O (0.5 mL) to dissolve any solids. The reaction was extracted with

EtOAc  $(4 \times 2 \text{ mL})$ . The combined organic layers were then dried over MgSO<sub>4</sub> and evaporated. Preparative TLC (70% EtOAc/hexanes, 500  $\mu$ m plate) provided (+)-1, irciniastatin A (1.7 mg), along with a monoprotected compound (SEM group on one of the phenolic oxygens). The monoprotected compound was dissolved in  $Et_2O$  (0.3 mL), and MeNO<sub>2</sub> (20  $\mu$ L) was added followed by MgBr<sub>2</sub> (13 mg, 20 equiv). After stirring for 15 min at rt, the reaction was diluted with EtOAc (2 mL), and sat. NaHCO<sub>3</sub> (1 mL) was added. The reaction was extracted with EtOAc  $(5 \times 1 \text{ mL})$ , and the combined organic layers were then dried over MgSO4 and concentrated. Preparative TLC (60% EtOAc/hexanes, 500  $\mu$ m plate) provided an additional 1.5 mg of the natural product to give a total of 3.2 mg (74% yield, 2 steps) of (+)-1 (irciniastatin A):  $[\alpha]_D^{20}$  +21.0 (c 0.1, CHCl<sub>3</sub>); IR (neat) 3366, 2965, 2928, 2853, 1654, 1515, 1459, 1381, 1254, 1173, 1107, 1067 cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz, CD<sub>3</sub>OD)  $\delta$  6.25 (s, 1H), 5.38 (d, *J* = 8.2 Hz, 1H), 4.74 (s, 1H), 4.72 (s, 1H), 4.49 (ddd, J = 12.2, 5.9, 3.1 Hz, 1H), 4.35 (d, J = 2.6 Hz, 1H), 3.97–3.92 (m, 2H), 3.67 (ddd, J = 9.3, 3.4, 2.7 Hz, 1H), 3.59 (dd, J = 10.9, 4.5 Hz, 1H), 3.50 (dd, J = 10.1, 1.3 Hz, 1H), 3.35 (s, 3H), 3.21 (s, 3H), 3.13 (dd, J = 16.7, 3.1 Hz, 1H), 2.86 (dd, J = 16.7, 12.2 Hz, 1H), 2.35 (dd, J = 14.6, 9.3 Hz, 1H), 2.11 (dd, J = 14.1, 3.6 Hz, 2H), 2.10 (s, 3H), 2.02 (ddd, J = 13.6, 4.3, 2.8 Hz, 1H), 1.91 (ddddd, J = 6.9, 6.9, 6.9, 6.9, 2.7 Hz, 1H), 1.77 (ddd, J = 13.7, 11.1, 6.3 Hz, 2H), 1.72 (s, 3H), 1.68 (ddd, J = 14.5, 3.2, 1.9 Hz, 2H), 1.10 (d, J = 7.0, 3H), 0.97 (s, 3H), 0.90 (s, 3H); <sup>13</sup>C NMR (125 MHz, CD<sub>3</sub>OD) δ 176.3, 172.5, 164.7, 163.8, 144.0, 141.3, 115.4, 113.1, 101.6, 101.5, 82.8, 82.3, 82.1, 79.9, 74.2, 73.6, 73.4, 72.2, 57.8, 56.7, 43.4, 39.9, 38.8, 34.5, 30.6, 29.6, 23.8, 23.0, 14.1, 10.9, 9.3; high resolution mass spectrum (ES+) *m/z* 632.3038  $[(M + Na)^+; calcd for C_{31}H_{47}NO_{11}Na 632.3047].$ 

Aldehyde 35. To a 0 °C solution of diol 37 (0.50 g, 4.8 mmol) in THF (24 mL) was added 60% NaH (0.19 g, 1.0 equiv) in three equal portions over 5 min. After 10 min, TBSCl (0.72 g, 1 equiv) dissolved in THF (5 mL) was added dropwise. After 4 h, the reaction was quenched with MeOH (2 mL). Sat. NaHCO<sub>3</sub> (15 mL) was added, and the reaction extracted with CH<sub>2</sub>Cl<sub>2</sub> (3 × 10 mL). The combined organic layers were then dried over MgSO<sub>4</sub> and concentrated. Flash chromatography (10% EtOAc/hexanes) provided TBS ether **66** (0.93 g, 89% yield) as a colorless oil: IR (neat) 3389, 2955, 2929, 2858, 1472, 1254, 1096, 1047 cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  3.46 (s, 4H), 2.86 (dd, *J* = 5.5, 5.5 Hz, 1H), 0.90 (s, 9H), 0.88 (s, 6H), 0.06 (s, 6H); <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  73.0, 72.5, 36.6, 26.0, 21.7, 18.4, -5.4; high resolution mass spectrum (CI+) *m/z* 219.1794 [(M + H)<sup>+</sup>; calcd for C<sub>11</sub>H<sub>27</sub>O<sub>2</sub>Si 219.1780].

To a 0 °C solution of TBS ether **66** (1.51 g, 6.91 mmol) in DMSO (4.9 mL, 10 equiv) and CH<sub>2</sub>Cl<sub>2</sub> (35 mL) was added Et<sub>3</sub>N (2.89 mL, 3 equiv) followed by SO<sub>3</sub>·pyridine (3.25 g, 3 equiv). After 1 h, brine (20 mL) and H<sub>2</sub>O (5 mL) were added. The layers were separated, and the aqueous layer was extracted with CH<sub>2</sub>Cl<sub>2</sub> (3 × 20 mL). The combined organic layers were then dried over MgSO<sub>4</sub> and concentrated. Flash chromatography (5% EtOAc/hexanes to 10% EtOAc/hexanes) provided aldehyde **35** (1.43 g, 96% yield) as a light yellow oil: IR (neat) 2956, 2930, 2893, 2858, 1733, 1472, 1257, 1104 cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  9.55 (s, 1H), 3.58 (s, 2H), 1.03 (s, 6H), 0.85 (s, 9H), 0.02 (s, 6H); <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  206.3, 68.6, 48.3, 26.0, 18.7, 18.4, -5.4; high resolution mass spectrum (CI+) *m/z* 215.1453 [(M – H)<sup>+</sup>; calcd for C<sub>11</sub>H<sub>23</sub>O<sub>2</sub>Si 215.1468].

**Alcohol (+)-34.** To a suspension of *N*-Ts-L-tryptophan (5.15 g, 1.2 equiv) in CH<sub>2</sub>Cl<sub>2</sub> (150 mL) was added Cl<sub>2</sub>BPh (1.87 mL, 1.2 equiv) dropwise. After 2 h at rt, the mixture was concentrated in vacuo, *n*-butyronitrile (50 mL) added, and the solution cooled to -78 °C. A mixture of aldehyde **35** (2.6 g, 12.0 mmol), ketene acetal **36** (3.86 g, 1.5 equiv), and isopropanol (1.38 mL, 1.5 equiv) in *n*-butyronitrile (15 mL) was added over 2 h via syringe pump. After an additional 2 h at -78 °C, the solution was warmed to 0 °C over 30 min, followed by addition of sat. NaHCO<sub>3</sub> (20 mL) and brine (20 mL). The resulting solution was extracted with Et<sub>2</sub>O (3 × 30 mL). The combined organic layers were then dried over MgSO<sub>4</sub> and evaporated. Flash chromatography (5% EtOAc/hexanes to 10% EtOAc/hexanes to 15% EtOAc/hexanes) provided (+)-**34** (2.49 g, 66% yield) as a colorless oil and a single enantiomer, as determined by Mosher ester analysis: [a]<sup>2D</sup><sub>20</sub>+27.9 (*c* 0.6, CHCl<sub>3</sub>); IR (neat) 3486, 2953, 2929, 2858, 1726, 1657, 1471, 1258,

1092 cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  7.10 (ddd, *J* = 15.6, 7.2, 7.2, Hz, 1H), 5.91 (ddd, *J* = 15.7, 1.5, 1.5, Hz, 1H), 3.75 (d, *J* = 3.2, 1H), 3.71 (s, 3H), 3.63 (ddd, *J* = 10.0, 2.9, 2.9 Hz, 1H), 3.49 (d, *J* = 2.0 Hz, 2H), 2.37 (dd, *J* = 14.5, 6.8 Hz, 1H), 2.26 (dddd, *J* = 14.6, 10.0, 7.5, 1.4 Hz, 1H), 0.91 (s, 3H), 0.89 (s, 9H), 0.85 (s, 3H), 0.07 (s, 6H); <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  167.1, 147.9, 122.6, 78.3, 73.6, 51.6, 38.6, 35.5, 26.0, 22.6, 19.0, 18.3, -5.4, -5.5; high resolution mass spectrum (ES+) *m*/*z* 317.2137 [(M + H)<sup>+</sup>; calcd for C<sub>16</sub>H<sub>33</sub>O<sub>4</sub>Si 317.2148].

**Epoxide (+)-38.** To a 0 °C solution of (+)-34 (0.226 g, 0.714 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (7.1 mL) was added 2,6-lutidine (0.164 mL, 2.0 equiv) followed by dropwise addition of TBSOTf (0.246 mL, 1.5 equiv). After 1 h, sat. NaHCO<sub>3</sub> (5 mL) was added, and the layers separated. The aqueous layer was extracted with  $CH_2Cl_2$  (3 × 10 mL), and the combined organic layers were dried over MgSO4 and concentrated. Flash chromatography (2% EtOAc/hexanes to 5% EtOAc/hexanes) provided TBS ether (+)-67 (0.269 g, 96% yield) as a colorless oil:  $[\alpha]_{\rm D}^{20}$ +3.6 (c 1.0, CHCl<sub>3</sub>); IR (neat) 2955, 2930, 2886, 2857, 1730, 1657, 1472, 1257, 1090 cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ 7.12–6.92 (m, 1H), 5.81 (ddd, J = 15.7, 1.5, 1.5 Hz, 1H), 3.74 (dd, J = 6.4, 4.4 Hz, 1H), 3.72 (s, 3H), 3.28 (dab, J = 9.5 Hz, 1H), 3.26 (dab, J = 9.6 Hz, 1H), 2.48 (dddd, J = 15.0, 6.4, 4.4, 1.7 Hz, 1H), 2.36–2.27 (m, 1H), 0.89 (s, 9H), 0.83 (s, 3H), 0.79 (s, 3H), 0.04 (s, 3H), 0.02 (s, 3H), 0.01 (s, 6H); <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>) δ 167.1, 148.9, 122.0, 75.1, 69.7, 51.6, 41.1, 36.6, 26.2, 26.1, 21.3, 20.9, 18.5, 18.4, -3.5, -4.1, -5.2, -5.3; high resolution mass spectrum (ES+) m/z 431.2997 [(M + H)<sup>+</sup>; calcd for C<sub>22</sub>H<sub>47</sub>O<sub>4</sub>Si<sub>2</sub> 431.3013].

A solution of TBS ether (+)-67 (0.217 g, 0.504 mmol) in  $CH_2Cl_2$  (10 mL) was cooled to -78 °C, and DIBAL-H (1.06 mL, 1.0 M in hexanes, 2.1 equiv) was slowly added. After 5 min, the reaction was quenched with MeOH (0.5 mL) and allowed to warm to rt. Sat. Rochelle's salt (10 mL) and EtOAc (10 mL) were added, and the solution stirred for 1 h.  $H_2O(10 \text{ mL})$  was then added to dissolve the suspended solids, and the layers were separated. The aqueous layer was extracted with EtOAc (3  $\times$ 10 mL), and the combined organic layers were dried over MgSO<sub>4</sub> and concentrated. Flash chromatography (5% EtOAc/hexanes to 10% EtOAc/hexanes) provided allylic alcohol (-)-68 (0.237 g, 99%) as a colorless oil:  $\left[\alpha\right]_{D}^{20}$  -3.4 (c 1.0, CHCl<sub>3</sub>); IR (neat) 3327, 2955, 2929, 2892, 2857, 1715, 1472, 1254, 1091, 1006 cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz,  $CDCl_3$ )  $\delta$  5.82–5.57 (m, 2H), 4.09 (d, J = 5.7, 2H), 3.64 (dd, J = 6.3, 4.3 Hz, 1H), 3.33 (dab, J = 9.5 Hz, 1H), 3.28 (dab, J = 9.5 Hz, 1H), 2.36 (ddd, *J* = 14.7, 4.8, 4.8 Hz, 1H), 2.16 (ddd, *J* = 14.5, 7.1, 7.1 Hz, 1H), 0.89 (s, 21H), 0.80 (s, 3H), 0.04 (s, 3H), 0.02 (s, 9H); <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>) *δ* 132.3, 130.1, 75.9, 69.9, 64.1, 41.2, 36.5, 26.3, 26.2, 21.4, 20.8, 18.5, 18.4, -3.2, -4.1, -5.2, -5.3; high resolution mass spectrum (ES+) m/z 403.3065 [(M + H)<sup>+</sup>; calcd for C<sub>21</sub>H<sub>47</sub>O<sub>3</sub>Si<sub>2</sub> 403.3063].

To freshly activated 3 Å molecular sieves (0.2 g) was added (-)-DIPT (15  $\mu$ L, 0.24 equiv) in CH<sub>2</sub>Cl<sub>2</sub> (0.5 mL). The solution was cooled to -20 °C, and Ti(O*i*-Pr)<sub>4</sub> (18  $\mu$ L, 0.2 equiv) in CH<sub>2</sub>Cl<sub>2</sub> (0.5 mL) was added followed by t-BuOOH (0.176 mL, 5 M in decane, 3.0 equiv). The reaction was stirred for 30 min, and then allylic alcohol (-)-68 (0.118 g, 0.293 mmol) dissolved in CH<sub>2</sub>Cl<sub>2</sub> (1 mL) was added via syringe. The flask and syringe were rinsed with  $CH_2Cl_2$  (0.4 mL) into the reaction flask. After 4 h, 10% aq. citric acid (2 mL) was added, and the reaction warmed to rt. After 1 h at rt, brine (3 mL) was added, the layers were separated, and the aqueous layer extracted with EtOAc (3  $\times$ 5 mL). The combined organic layers were dried over MgSO4 and concentrated. Flash chromatography (5% EtOAc/hexanes to 10% EtOAc/hexanes to 15% EtOAc/hexanes) provided epoxide (+)-38 (0.108 g, 88% yield, 13:1 inseparable mixture of epoxides) as a colorless oil: [α]<sup>20</sup><sub>D</sub> +24.3 (*c* 1.0, CHCl<sub>3</sub>); IR (neat) 3430, 2955, 2929, 2895, 2857, 1472, 1254, 1091, 1009 cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ 3.91 (ddd, *J* = 12.5, 5.4, 2.4 Hz, 1H), 3.80 (dd, *J* = 7.4, 3.8 Hz, 1H), 3.60 (ddd, *J* = 12.5, 7.2, 4.4 Hz, 1H), 3.35 (dab, J = 9.6 Hz, 1H), 3.28 (dab, J = 9.6 Hz, 1H), 3.12 (ddd, J = 6.9, 4.5, 2.3 Hz, 1H), 2.93 (ddd, J = 4.6, 2.4, 2.4 Hz, 1H), 1.80 (dd, J = 7.1, 5.8 Hz, 1H), 1.79–1.72 (m, 1H), 1.65 (ddd, J = 14.6, 7.0, 3.8 Hz, 1H), 0.90 (s, 9H), 0.89 (s, 9H), 0.83 (s, 3H), 0.79 (s, 3H), 0.09 (s, 3H), 0.08 (s, 3H), 0.01 (s, 6H); <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>) & 73.7, 69.7, 61.9, 60.1, 54.1, 40.7, 35.5, 26.3, 26.1, 21.5, 20.5, 18.6, 18.5, -3.8, -4.0, -5.2, -5.3; high resolution mass spectrum (ES+) m/z 441.2846 [(M + Na)<sup>+</sup>; calcd for C<sub>21</sub>H<sub>46</sub>O<sub>4</sub>Si<sub>2</sub>Na 441.2833].

**Ester (+)-39.** Alcohol (+)-38 (0.199 g, 0.474 mmol) in CH<sub>3</sub>CN (4.7 mL) was treated with TEMPO (6 mg, 0.08 equiv) followed by addition of pH 7 buffer (4.7 mL). The reaction was then treated with NaClO<sub>2</sub>, followed by dropwise addition of NaOCl (0.18 mL, 5% aq. solution). After 2 h, sodium sulfite (200 mg, 3.3 equiv) was added, and the reaction stirred for 30 min, where the solution turned from a light orange color to colorless. The reaction was acidified to pH 4 with 10% aq. citric acid and extracted with EtOAc ( $5 \times 10$  mL). The combined organic layers were dried over MgSO<sub>4</sub> and concentrated to yield carboxylic acid, which was carried forward without further purification.

The unpurified acid was dissolved in CH2Cl2 (2.4 mL) and MeOH (2.4 mL). The solution was cooled to 0 °C, and TMS-diazomethane (0.28 mL, 2 M in Et<sub>2</sub>O, 1.2 equiv) was added dropwise until the solution remained yellow in color. Argon gas was bubbled through the reaction for 10 min, and then the solvent evaporated. Flash chromatography (5% EtOAc/hexanes) provided ester (+)-39 (0.195 g, 91% yield, 2 steps) as a colorless oil:  $[\alpha]_{D}^{20}$  +15.8 (c 1.0, CHCl<sub>3</sub>); IR (neat) 2955, 2930, 2887, 2857, 1759, 1472, 1254, 1201, 1089, 1006 cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz,  $CDCl_3$ )  $\delta$  3.81 (dd, J = 7.4, 3.8 Hz, 1H), 3.75 (s, 3H), 3.34 (dab, J = 9.6 Hz, 1H), 3.30 (ddd, J = 6.5, 4.5, 1.9 Hz, 1H), 3.28 (dab, J = 9.6 Hz, 1H), 3.22 (d, J = 1.9 Hz, 1H), 1.80 (ddd, J = 14.6, 7.4, 4.5 Hz, 1H), 1.67 (ddd, J = 14.7, 6.8, 3.8 Hz, 1H), 0.90 (s, 9H), 0.88 (s, 9H), 0.83 (s, 3H), 0.78 (s, 3H), 0.10 (s, 3H), 0.08 (s, 3H), 0.01 (s, 6H); <sup>13</sup>C NMR (125 MHz,  $CDCl_3$ )  $\delta$  169.9, 73.6, 69.6, 56.9, 54.6, 52.5, 40.7, 35.5, 26.3, 26.1, 21.3, 20.7, 18.6, 18.5, -3.8, -4.0, -5.2, -5.3; high resolution mass spectrum (ES+) m/z 469.2778 [(M + Na)<sup>+</sup>; calcd for C<sub>22</sub>H<sub>46</sub>O<sub>5</sub>Si<sub>2</sub>Na 469.2782].

Aldehyde (+)-40. To bis-silyl ether (+)-39 (72.0 mg, 0.159 mmol) was added a HF·pyridine/pyridine/THF solution (0.64 mL, 10 equiv HF, stock solution made up of 0.5 mL of HF pyridine, 1.0 mL of pyridine, and 5.0 mL of THF). After stirring 20 h at rt, sat. NaHCO<sub>3</sub> (5 mL) was added slowly. Once all gas evolution had ceased, the reaction was extracted with  $CH_2Cl_2$  (3 × 5 mL). The combined organic layers were dried over MgSO<sub>4</sub> and concentrated. Flash chromatography (5% EtOAc/hexanes to 10% EtOAc/hexanes to 15% EtOAc/hexanes to 20% EtOAc/hexanes) provided alcohol (+)-**69** as a single diastereomer (37.2 mg, 72% yield) as a colorless oil:  $[\alpha]_{\rm D}^{20}$  +46.8 (*c* 0.5, CHCl<sub>3</sub>); IR (neat) 3465, 2955, 2930, 2857, 1744, 1446, 1290, 1253, 1205, 1089, 1051, 1004  $cm^{-1}$ ; <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  3.80 (dd, *J* = 7.4, 3.8 Hz, 1H), 3.77 (s, 3H), 3.59 (d, J = 10.9 Hz, 1H), 3.36–3.27 (m, 2H), 3.26 (d, J = 1.7 Hz, 1H), 2.37 (s, 1H), 1.95 (ddd, J = 14.8, 7.4, 3.9 Hz, 1H), 1.67 (ddd, J = 14.8, 7.3, 3.8 Hz, 1H), 0.99 (s, 3H), 0.91 (s, 9H), 0.81 (s, 3H), 0.14 (s, 3H), 0.14 (s, 3H); <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  169.6, 76.7, 70.0, 56.6, 54.5, 52.7, 39.6, 35.6, 26.2, 23.0, 21.5, 18.4, -3.9, -4.0; high resolution mass spectrum (ES+) m/z 355.1908 [(M + Na)<sup>+</sup>; calcd for C<sub>16</sub>H<sub>32</sub>O<sub>5</sub>SiNa 355.1917].

To a solution of alcohol (+)-69 (0.025 g, 0.075 mmol) in  $CH_2Cl_2$ (0.75 mL) and DMSO (0.75 mL) at 0 °C was added Et<sub>3</sub>N (0.105 mL, 10 equiv) followed by  $SO_3$  pyridine (0.047 g, 4 equiv) in one portion. After 2 h, sat. NaHCO<sub>3</sub> (4 mL) was added, and the layers separated. The aqueous layer was extracted with  $CH_2Cl_2$  (3 × 5 mL), and the combined organic layers were dried over MgSO4 and concentrated. Flash chromatography (5% EtOAc/hexanes to 10% EtOAc/hexanes to 20% EtOAc/hexanes) provided aldehyde (+)-40 (23.5 mg, 95% yield) as a light yellow oil:  $[\alpha]_{D}^{20}$  +22.9 (c 0.5, CHCl<sub>3</sub>); IR (neat) 2955, 2931, 2857, 1754, 1448, 1291, 1254, 1205, 1093 cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  9.56 (s, 1H), 4.06 (dd, J = 7.9, 3.6 Hz, 1H), 3.77 (s, 3H), 3.28 (ddd, J = 7.5, 3.8, 1.9 Hz, 1H), 3.24 (d, J = 1.9 Hz, 1H), 1.90 (ddd, J = 14.6, 7.9, 3.8 Hz, 1H), 1.54 (ddd, J = 14.7, 7.5, 3.6 Hz, 1H), 1.06 (s, 3H), 1.02 (s, 3H), 0.88 (s, 9H), 0.13 (s, 3H), 0.08 (s, 3H); <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>) δ 205.8, 169.4, 73.7, 56.0, 54.4, 52.7, 51.3, 35.7, 26.1, 19.3, 18.4, 17.7, -3.9, -4.0; high resolution mass spectrum (ES+) m/z 353.1753 [(M + Na)<sup>+</sup>; calcd for C<sub>16</sub>H<sub>30</sub>O<sub>5</sub>SiNa 353.1761].

**Tetrahydropyran (+)-41.** To a solution of (-)-DIPCl (0.035 g, 1.8 equiv) in  $\text{Et}_2\text{O}$  (0.4 mL) cooled to -78 °C was added  $\text{Et}_3\text{N}$  (0.021 mL, 2.5 equiv) followed by slow dropwise addition of 2-butanone (0.010 mL, 1.8 equiv). The solution immediately turned cloudy white. After stirring for 2 h at -78 °C, aldehyde (+)-40 (0.020 g, 0.061 mmol) was dissolved in  $\text{Et}_2\text{O}$  (0.4 mL) and added dropwise to the boron enolate over 5 min via syringe. The flask and syringe were then rinsed with  $\text{Et}_2\text{O}$  (0.4 mL), and the rinse added to the reaction over 2 min. After stirring at -78 °C

for 5 h, the solution was warmed to -40 °C over 1 h and held at -40 °C. After 12 h, the reaction was warmed to 0 °C, and a 1:1:1 mixture of MeOH/pH 7 buffer/30% aq. H<sub>2</sub>O<sub>2</sub> (1 mL) was added. After 1 h, sat. Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub> (4 mL) was added very slowly over 30 min to destroy any remaining peroxides. The reaction was then extracted with CH<sub>2</sub>Cl<sub>2</sub> (3 × 10 mL). The combined organic layers were dried over MgSO<sub>4</sub> and concentrated. Flash chromatography (10% EtOAc/hexanes to 15% EtOAc/hexanes to 20% EtOAc/hexanes) provided the aldol product (0.020 g, 86% yield) as a 5:1 mixture of diastereomers.

The aldol products were dissolved in CH<sub>2</sub>Cl<sub>2</sub> (1 mL), and CSA (2.3 mg, 0.2 equiv) was added. After 2 h, sat. NaHCO<sub>3</sub> (2 mL) added, and the reaction stirred for 5 min. The layers were separated, and the aqueous layer was extracted with  $CH_2Cl_2$  (3 × 5 mL). The combined organic layers were dried over MgSO4 and concentrated. Flash chromatography (10% EtOAc/hexanes to 15% EtOAc/hexanes to 20% EtOAc/hexanes) provided (+)-41 (14.8 mg, 60% yield over 2 steps for the major diastereomer) and (-)-42 (2.9 mg, 12% yield over 2 steps for the minor diastereomer). Major diastereomer (+)-41:  $\left[\alpha\right]_{D}^{20}$  +9.8 (c 0.5, CHCl<sub>3</sub>); IR (neat) 2954, 2930, 2861, 1741, 1714, 1461, 1388, 1366, 1258, 1074  $cm^{-1}$ ; <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  4.26 (dd, *J* = 6.3, 4.2 Hz, 1H), 4.15 (ddd, J = 8.0, 3.8, 3.8 Hz, 1H), 4.06 (dd, J = 10.0, 3.5 Hz, 1H), 3.78 (s, 3H), 3.64 (dd, J = 5.0, 3.3 Hz, 1H), 2.93 (d, J = 6.3 Hz, 1H), 2.57 (dd, J = 15.6, 3.6 Hz, 1H), 2.50–2.42 (m, 2H), 2.05 (ddd, J = 13.5, 9.3, 3.2 Hz, 1H), 1.45–1.38 (m, 1H), 1.40–1.36 (m, 1H), 1.05 (dd, J = 7.3, 7.3 Hz, 3H), 1.00 (s, 3H), 0.92 (s, 9H), 0.85 (s, 3H), 0.05 (s, 3H), 0.04 (s, 3H); <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>) δ 210.8, 173.3, 77.7, 73.5, 73.4, 68.8, 52.7, 42.4, 37.5, 36.9, 29.8, 26.0, 25.8, 20.7, 18.2, 7.9, -4.4, -4.8; high resolution mass spectrum (ES+) m/z 425.2330 [(M + Na)<sup>+</sup>; calcd for  $C_{20}H_{38}O_6$ SiNa 425.2336]. Minor diastereomer (-)-42:  $[\alpha]_D^{20}$  -38.7 (c 0.8, CHCl<sub>3</sub>); IR (neat) 3480, 2955, 2930, 2885, 2857, 1741, 1719, 1255, 1096, 1060, 1008 cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  4.24 (dd, J = 6.0, 3.3 Hz, 1H), 4.17 (dd, J = 10.5, 2.3 Hz, 1H), 4.08 (ddd, J = 12.2, 2.7, 2.7) Hz, 1H), 3.75 (s, 3H), 3.55 (dd, J = 2.7, 2.7 Hz, 1H), 2.79 (d, J = 6.1 Hz, 1H), 2.59–2.40 (m, 3H), 2.25 (dd, J = 14.9, 2.4 Hz, 1H), 2.04 (ddd, J = 14.2, 12.2, 2.4 Hz, 1H), 1.22 (ddd, J = 13.4, 3.0, 3.0 Hz, 1H), 1.04 (dd, J = 7.3, 7.3 Hz, 3H), 0.91 (s, 9H), 0.87 (s, 3H), 0.81 (s, 3H), 0.04 (s, 3H), 0.01 (s, 3H); <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  210.8, 172.9, 76.3, 74.9, 73.9, 73.4, 52.6, 42.8, 37.4, 37.2, 30.0, 26.0, 23.7, 19.7, 18.3, 7.8, -4.4, -4.7; high resolution mass spectrum (ES+) m/z 425.2336 [(M + Na)<sup>+</sup>; calcd for C20H38O6SiNa 425.2336].

Tetrahydropyran (+)-6. To a 0 °C solution of (+)-41 (14.8 mg, 0.036 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (0.18 mL) was added proton sponge (95 mg, 12.0 equiv) followed by Me<sub>3</sub>O·BF<sub>4</sub> (54 mg, 10 equiv). The reaction was then stirred at rt. After 4 h, 10% aq. citric acid (2 mL) was added. The layers were separated, and the aqueous layer extracted  $CH_2Cl_2$  (3 × 5 mL). The combined organic layers were washed with 10% aq. citric acid (4 mL) and then brine (4 mL). The organic layer was then dried over MgSO<sub>4</sub> and concentrated. Flash chromatography (10% EtOAc/ hexanes) provided (+)-6 (14.2 mg, 92% yield) as a colorless oil. All spectral data matched that of the ketone fragment constructed during the first-generation approach:  $[\alpha]_D^{20}$  +16.5 (c 1.7, CHCl<sub>3</sub>); IR (neat) 2954, 2934, 2886, 2858, 1754, 1722, 1712, 1462, 1361, 1255, 1119, 1073  $cm^{-1}$ ; <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) 4.08 (dd, J = 11.3, 6.0 Hz, 1H), 3.99 (dd, J = 9.6, 3.0 Hz, 1H), 3.90 (d, J = 6.4 Hz, 1H), 3.73 (s, 3H), 3.62 (dd, *J* = 7.5, 3.6 Hz, 1H), 3.40 (s, 3H), 2.91 (dd, *J* = 15.2, 9.7 Hz, 1H), 2.56– 2.29 (m, 3H), 1.93 (ddd, J = 13.8, 6.1, 3.9 Hz, 1H) 1.60 (ddd, J = 12.7, 9.0, 5.3 Hz, 1H), 1.03 (dd, J = 7.3, 7.3 Hz, 3H), 0.96 (s, 3H), 0.90 (s, 9H), 0.83 (s, 3H), 0.04 (s, 6H);  $^{13}\text{C}$  NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  210.4, 171.7, 82.5, 77.4, 73.1, 69.8, 58.8, 52.2, 42.6, 38.1, 37.2, 30.3, 26.0, 25.1, 18.2, 17.9, 7.8, -4.2, -4.8; high resolution mass spectrum (ES+) m/z439.2491 [ $(M + Na)^+$ ; calcd for  $C_{21}H_{40}O_6$ SiNa 439.2492].

Aldehyde 45. To a solution of bis-phenol 14 (519 mg, 2.043 mmol) in acetone (25.0 mL) was added  $K_2CO_3$  (1.03 g, 7.452 mmol, 3.7 equiv) followed by dropwise addition of a solution of 3,4-dimethoxybenzyl-bromide (2.0 mL, 2.2 M in acetone, 2.2 equiv). The yellow reaction mixture was stirred for 24 h and quenched with  $H_2O$  (20 mL). The layers were separated, and the aqueous layer was extracted with EtOAc (3 × 20 mL). The combined organic layers were dried over MgSO<sub>4</sub>, filtered and concentrated in vacuo. The crude mixture was purified via flash chromatography (40% EtOAc:hexanes to 50% EtOAc:hexanes) to

provide bis-DMB ether **70** (850 mg, 1.532 mmol, 75%) as a colorless solid: mp 102.0–103.0 °C; IR (neat) 2947, 1732, 1594, 1515, 1459, 1264, 1152, 1025 cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) 6.97–6.81 (m, 6 H), 6.52 (s, 1 H), 4.99 (s, 2 H), 4.95 (s, 2 H), 3.89 (s, 3 H), 3.88 (s, 3 H), 3.88 (s, 3 H), 3.87 (s, 3 H), 3.83 (s, 3 H), 3.69 (s, 2 H), 3.67 (s, 3 H), 2.14 (s, 3 H); <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  171.2, 158.7, 155.2, 149.4, 149.3, 149.1, 148.9, 132.7, 129.6, 129.4, 120.0, 119.7, 119.6, 117.8, 111.3, 111.1, 110.9, 110.1, 98.3, 71.2, 70.7, 56.1, 56.1, 56.1, 56.0, 52.2, 36.2, 11.7; HRMS (ES+) *m*/*z* 577.2047 [(M + Na)<sup>+</sup>; calcd for C<sub>30</sub>H<sub>34</sub>O<sub>10</sub>Na 577.2050].

A solution of bis-DMB ether 70 (850 mg, 1.53 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (15.3 mL) was cooled to -78 °C, and DIBAL-H (2.1 mL, 1.0 M in hexanes, 1.4 equiv) was added over 20 min via syringe pump. The reaction mixture was allowed to stir for an additional 5 min before it was quenched by the addition of MeOH (7.0 mL) and was warmed to room temperature and then diluted with EtOAc (2 mL) and a saturated aq. solution of Rochelle's salt (2 mL). The biphasic reaction mixture was allowed to stir for 1 h at room temperature to allow the organic layer to transition from cloudy to clear. The layers were separated, and the aqueous layer was extracted with EtOAc ( $3 \times 3$  mL). The combined organic layers were dried over MgSO4, filtered and concentrated in vacuo. The crude mixture was purified via flash chromatography (40% EtOAc/hexanes to 50% EtOAc/hexanes) to provide aldehyde 45 (697 mg, 1.33 mmol, 87%) as a colorless solid: mp 122.5-124.0 °C; IR (neat) 2944, 2838, 2726, 1722, 1594, 1515, 1459, 1265, 1153, 1030 cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) 9.65 (t, J = 1.8 Hz, 1 H), 6.97 (d, J = 1.7 Hz, 1 H), 6.94–6.83 (m, 5 H), 6.55 (s, 1 H), 5.02 (s, 2 H), 4.97 (s, 2 H), 3.90 (s, 3 H), 3.89 (s, 9 H), 3.84 (s, 3 H), 3.68 (d, J = 1.6 Hz, 2 H), 2.10 (s, 3 H); <sup>13</sup>C NMR (125 MHz, MeOD) δ 198.6, 168.7, 158.9, 155.4, 149.4, 149.4, 149.2, 149.0, 131.0, 129.4, 129.2, 120.1, 119.8, 119.7, 118.0, 111.3, 111.1, 110.0, 110.7, 98.5, 71.2, 70.7, 56.3, 56.1, 56.1, 56.1, 52.3, 46.1, 11.9; HRMS (ES+) m/z 525.2141 [(M + Na)<sup>+</sup>; calcd for C<sub>29</sub>H<sub>33</sub>O<sub>9</sub> 525.2125]

 $\beta$ -Hydroxy Ketone (+)-46. A solution of ketone (+)-6 (114 mg, 0.274 mmol, azeotroped  $\times$  3 with benzene, placed under a high vacuum overnight) in CH<sub>2</sub>Cl<sub>2</sub> (1.35 mL) was cooled to -78 °C, and Cl<sub>2</sub>BPh (0.05 mL, 0.383 mmol, 1.4 equiv) was added dropwise. After 20 min, i-Pr<sub>2</sub>NEt (0.10 mL, 0.574 mmol, 2.1 equiv) was introduced dropwise. After 1 h, the reaction mixture was warmed to 0 °C, where it was stirred for 1 h and then cooled back down to -78 °C. Aldehyde 45 (205 mg, 0.391 mmol, 1.5 equiv) was dissolved in CH<sub>2</sub>Cl<sub>2</sub> (0.7 mL) and was added to the boron enolate solution at -78 °C over 15 min via syringe pump. After 4 h at -78 °C, the reaction mixture was quenched with a 1:1 mixture of MeOH and pH 7 buffer (4 mL). While warming to 0 °C, pH 8 buffer solution was added to neutralize the reaction mixture to pH 7, and the biphasic mixture was stirred for 1 h at 0 °C. The layers were separated, and the aqueous layer was extracted with  $CH_2Cl_2$  (3 × 5 mL). The combined organic layers were dried over MgSO4, filtered and concentrated in vacuo. The resulting crude mixture was purified via flash chromatography on deactivated SiO2 (2% v/v triethylamine, 7.5% EtOAc/CH<sub>2</sub>Cl<sub>2</sub> to 10% EtOAc/CH<sub>2</sub>Cl<sub>2</sub>) to provide a mixture (ca. 10:1) of  $\beta$ -hydroxy ketone (+)-46 and corresponding lactone (176 mg, 0.192 mmol, 70%) as a colorless foam:  $[\alpha]_D^{20}$  +29.8 (c 0.68 CHCl3); IR (neat) 3417, 2924, 2855, 1719, 1590, 1516, 1461, 1264, 1154 cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) Major δ 6.95–6.91 (m, 3 H), 6.88–6.82 (m, 3 H), 6.49 (s, 1 H), 5.02–4.97 (m, 2 H), 4.95 (s, 2 H), 4.10 (dd, J = 5.3, 10.9 Hz, 1 H), 4.06 (dd, J = 2.9, 9.1 Hz, 1 H), 4.06–4.03 (m, 1 H), 3.94 (d, J = 6.3 Hz, 1 H), 3.89 (s, 3 H), 3.89 (s, 3 H), 3.88 (s, 3 H), 3.88 (s, 3 H), 3.85 (s, 3 H), 3.72 (s, 3 H), 3.64 (dd, J = 3.7, 7.7 Hz, 1 H), 3.46 (d, J = 5.6 Hz, 1 H), 3.41 (s, 3 H), 3.06 (dd, J = 9.4, 16.0 Hz, 1 H), 2.87 (dd, J = 3.4, 14.3 Hz, 1 H), 2.68 (dq, J = 7.1, 5.4 Hz, 1 H), 2.62 (dd, J = 10.1, 14.1 Hz, 1 H), 2.54 (dd, J = 3.0, 16.4 Hz, 1 H), 2.20 (s, 3 H), 1.96 (ddd, J = 3.9, 6.0, 13.9 Hz, 1 H), 1.58 (ddd, J = 5.0, 7.9, 13.3 Hz, 1 H), 1.21 (d, J = 6.9 Hz, 3 H), 0.96 (s, 3 H), 0.90 (s, 9 H), 0.85 (s, 3 H), 0.04 (s, 3 H), 0.04 (s, 3 H); Distinct peaks from minor byproduct  $\delta 6.54$  (s), 5.16 (dab, J = 12.2 Hz), 5.10 (dab, J = 11.8 Hz), 1.33 (d, J = 7.0 Hz); <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>) δ 212.5, 171.5, 170.7, 159.0, 155.2, 149.4, 149.3, 149.1, 148.9, 137.1, 129.5, 129.5, 120.0, 119.8, 119.3, 117.9, 111.3, 111.1, 110.8, 110.7, 97.6, 82.3, 77.4, 76.7, 73.0, 71.6, 71.2, 70.6, 70.1, 58.8, 56.1, 56.1, 56.1, 56.1, 53.2, 52.6, 52.1, 42.4, 38.0, 35.8, 30.0, 26.0, 24.9, 18.2, 11.8, 11.4,

-4.3, -4.9; HRMS (ES+) m/z 963.4525 [(M + Na)<sup>+</sup>; calcd for C<sub>50</sub>H<sub>72</sub>O<sub>15</sub>SiNa 963.4538].

Acid (+)-47. To a solution of  $\beta$ -hydroxy ketone (+)-46 (203 mg, 0.215 mmol) in THF (2.20 mL) and MeOH (0.73 mL) cooled to -78  $^{\circ}$ C was added a solution of Et<sub>2</sub>BOMe (0.31 mL, 1 M in THF, 1.4 equiv). The reaction mixture was stirred for 25 min before NaBH<sub>4</sub> (45 mg, 1.189 mmol, 5.4 equiv) was added. After 5.5 h, the reaction was warmed to 0 °C and quenched with a 1:1 mixture of MeOH and pH 7 buffer (4.0 mL) followed by the addition of m-CPBA (0.330 g, 1.31 mmol, 6.0 equiv) portion-wise. After 30 min, the reaction mixture was warmed to room temperature, and dimethylsulfide was added slowly to quench remaining peroxides. After 10 min, aq. solution of  $K_2CO_3$  (5 mL, 0.1 N) was added, and the layers were separated. The aqueous layer was extracted with EtOAc ( $3 \times 5$  mL). The combined organic layers were washed with brine and then dried over MgSO<sub>4</sub>, filtered and concentrated in vacuo. The crude mixture was purified via flash chromatography (50% EtOAc/hexanes to 60% EtOAc/hexanes then flushed with 10% MeOH/ EtOAc) to provide a mixture (ca. 8:1) of diol and lactone (183 mg).

The mixture of diol and lactone was dissolved in MeOH (10.0 mL) and cooled to 0  $^{\circ}\mathrm{C}$  followed by the addition of H\_2O (70  $\mu\mathrm{L}$ , 3.889 mmol, 20.0 equiv) and LiOH (191 mg, 7.975 mmol, 40.0 equiv). The reaction mixture was allowed to warm to room temperature, and after 35 h, the reaction mixture was quenched with 50% aqueous acetic acid solution (5 mL). Brine (6 mL) and H<sub>2</sub>O (2 mL) were added, and the layers were separated. The aqueous layer was extracted with EtOAc ( $5 \times 15$  mL). The combined organic layers were dried over MgSO4, filtered and concentrated in vacuo. Crude mixture was purified via column chromatography on SiO<sub>2</sub> (0.1% acetic acid in 60% EtOAc/hexanes to 0.1% acetic acid in 80% EtOAc/hexanes) to provide acid (+)-47 (136 mg, 0.152 mmol, 69% over two steps) as a colorless foam:  $[a]_{\rm D}^{20}$  +22.4 (c 0.9 CHCl<sub>3</sub>); IR (neat) 3522, 3058, 2937, 2862, 2862, 1712, 1590, 1515, 1461, 1381, 1261, 1153, 1090 cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ 7.33 (d, J = 1.7 Hz, 1 H), 6.92 (m, 3 H), 6.86 (d, J = 8.0 Hz, 1 H), 6.82 (d, J = 8.3 Hz, 1 H), 6.51 (s, 1 H), 5.15 (dab, J = 11.8 Hz, 1 H), 5.09 (dab J = 12.2 Hz, 1 H), 4.99 (s, 2 H), 4.33 (ddd, J = 2.3, 5.5, 8.0 Hz, 1 H) 4.19 (dd, J = 6.0, 11.7 Hz, 1 H), 4.13 (m, 1 H), 3.91 (s, 3 H), 3.89 (s, 3 H), 3.88 (s, 3 H), 3.86 (s, 3 H), 3.85 (d, J = 4.1 Hz, 1 H), 3.60 (d, J = 12.3 Hz, 1 H), 3.57 (dd, J = 4.0, 7.6 Hz, 1 H) 3.41 (s, 3 H), 3.02 (dd, J = 2.7, 17.2 Hz, 1 H), 2.88 (dd, J = 12.3, 16.4 Hz, 1 H), 2.11 (s, 3 H), 2.08 (m, 1 H), 2.01-1.93 (m, 2 H), 1.66 (ddd, J = 5.1, 8.0, 13.6 Hz, 1 H), 1.56 (app d, J = 14 Hz, 1 H), 1.07 (d, J = 6.9 Hz, 3 H), 0.94 (s, 3 H), 0.89 (s, 9 H), 0.87 (s, 3 H), 0.05 (s, 3 H), 0.04 (s, 3 H);  $^{13}$ C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  172.3, 164.4, 161.4, 160.4, 149.5, 149.4, 149.2, 148.7, 142.2, 129.6, 128.9, 120.2, 119.0, 116.3, 111.3, 111.0, 110.9, 110.7, 107.7, 97.9, 82.0, 81.8, 79.0, 72.6, 71.6, 71.1, 70.5, 69.8, 58.7, 56.2, 56.1, 41.5, 38.7, 33.1, 30.6, 29.4, 26.0, 25.0, 18.2, 11.3, 9.5, -4..2, -4.8; HRMS (ES+) m/z 919.4287 [(M + Na)<sup>+</sup>; calcd for  $C_{48}H_{68}O_{14}SiNa 919.4276$ ].

**SEM-Ether (+)-48.** A solution of acid (+)-47 (116 mg, 0.129 mmol) in freshly distilled acetone (6.6 mL) was cooled to 0 °C followed by the dropwise addition of i-Pr<sub>2</sub>NEt (0.05 mL, 0.287 mmol, 2.2 equiv) and a solution of isobutylchloroformate (0.50 mL, 0.64 M in acetone, 2.4 equiv). After 1 h, a solution of NaN<sub>3</sub> (0.85 mL, 0.78 M in H<sub>2</sub>O, 5 equiv) was added to the reaction mixture dropwise. After an additional 20 min at 0 °C, the reaction mixture was diluted with cold H<sub>2</sub>O (15 mL). The layers were separated, and the aqueous layer was extracted with cold EtOAc (3  $\times$  10 mL). The combined organic layers were dried over MgSO<sub>4</sub>, filtered and concentrated in vacuo. The crude acyl azide was azeotroped (benzene  $\times$  3) and placed under a high vacuum (~0.1 mmHg) for 30 min. The acyl azide was dissolved in toluene (6.6 mL) and reaction flask was fitted with a reflux condenser and heated to 80  $^\circ$ C. After 45 min, 2-TMS-ethanol (0.67 mL, 4.674 mmol, 36.2 equiv) was added via syringe through the top of the condenser. After 5 h at 80 °C, the reaction mixture was cooled to room temperature and concentrated in vacuo. The crude mixture was purified via flash chromatography (40% EtOAc/hexanes to 50% EtOAc/hexanes) to provide N,O-aminal (+)-71 (90 mg, 0.087 mmol, 67%) as a colorless foam:  $[\alpha]_{D}^{20}$  +3.7 (c 0.5, CHCl<sub>3</sub>); IR (neat) 3502, 3340, 2947, 2859, 1713, 1589, 1515, 1462, 1253, 1151, 1078, 1034, 840 cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ 7.34 (d, J = 1.8 Hz, 1 H), 6.93 (m, 3 H), 6.87 (d, J = 8.0 Hz, 1 H), 6.82 (d, J =8.3 Hz, 1 H), 6.52 (s, 1 H), 5.38, (d, J = 8.8 Hz, 1 H), 5.16 (dab, J = 12.0,

1 H), 5.09 (dab, *J* = 11.9 Hz, 1 H), 4.99 (s, 2H), 4.89 (d, *J* = 9.6 Hz, 1 H), 4.31 (ddd, *J* = 2.2, 6.6, 10.5 Hz, 1 H), 4.19 (ddd, *J* = 8.3, 10.3, 18.2 Hz, 1 H), 4.12 (m 2 H), 3.98 (d, *J* = 9.1 Hz, 1 H), 3.92 (s, 3 H), 3.89 (s, 3 H), 3.88 (s, 3 H), 3.87 (m, 1 H), 3.86 (s, 3 H), 3.66 (s, 1 H), 3.62 (d, *J* = 11.5 Hz, 1 H), 3.56 (m, 1 H), 3.35 (s, 3 H), 3.10 (app d, *J* = 16.5 Hz, 1 H), 2.83 (dd, *J* = 12.1, 16.0 Hz, 1 H), 2.37 (m, 1 H), 2.12 (s, 3 H), 1.85 (m, 2 H), 1.45–1.41 (m, 2 H), 1.11 (d, *J* = 6.9 Hz, 3 H), 0.10 (s, 3 H), 0.03 (s, 3 H), 0.01 (s, 9 H); <sup>13</sup>C NMR (125 MHz, CDCl3)  $\delta$  163.7, 161.2, 160.3, 157.1, 149.6, 149.4, 149.2, 148.7, 142.2, 129.7, 129.0, 120.1, 119.0, 116.2, 111.3, 111.0, 110.9, 110.8, 108.2, 97.9, 83.7, 83.6, 79.3, 77.4, 73.0, 72.7, 71.1, 70.5, 63.7, 56.2, 56.1, 56.1, 55.8, 43.4, 38.0, 32.9, 31.1, 29.7, 26.1, 26.0, 18.1, 17.8, 11.4, 10.1, -1.3, -4.4, -4.8. HRMS (ES+) *m/z* 1034.5088 [(M + Na)+; calcd for C<sub>53</sub>H<sub>81</sub>NO<sub>14</sub>Si<sub>2</sub>Na 1034.5093].

A solution of N,O-aminal (+)-71 (86 mg, 0.085 mmol) in THF (0.6 mL) was cooled to 0 °C followed by the addition of *i*-Pr<sub>2</sub>NEt (0.13 mL, 0.746 mmol, 9 equiv), SEMCl (0.09 mL, 0.509 mmol, 6 equiv), and TBAI (8 mg, 0.022 mmol, 0.2 equiv). The reaction mixture was stirred at 0 °C for 15 min and was allowed to warm to room temperature and stirred for 23 h. The reaction mixture was quenched with a saturated aq. solution of NaHCO3 (1 mL). The aqueous layer was extracted with EtOAc  $(3 \times 3 \text{ mL})$ . The combined organic layers were dried over MgSO<sub>4</sub>, filtered and concentrated in vacuo. The crude mixture was purified via flash chromatography on SiO<sub>2</sub> (35% EtOAc/hexanes to 40% EtOAc/hexanes) to provide SEM ether (+)-48 (0.080 g, 0.070 mmol, 82%) as a colorless foam:  $[\alpha]_{D}^{20}$  +31.6 (c 0.8, CHCl3); IR (neat) 3336, 2952, 2929, 2858, 1716, 1593, 1518, 1464, 1264, 1249, 1160, 1078, 1027, 836 cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  7.33 (d, J = 1.8 Hz, 1 H), 6.98–6.93 (m, 3 H), 6.88 (d, J = 8.7 Hz, 1 H), 6.83 (d, J = 8.3 Hz, 1 H), 6.54 (s, 1 H), 5.60 (bd, J = 9.2 Hz, 1 H), 5.19 (dab, J = 11.7 Hz, 1 H), 5.10 (dab, J = 11.6 Hz, 1 H), 4.98 (s, 2 H), 4.81 (bd, J = 7.9 Hz, 1 H), 4.69–4.59 (m, 2 H), 4.26 (ddd, J = 2.3, 8.4, 11.4 Hz, 1 H), 4.17 (dd, J = 7.5, 10.5 Hz, 1 H), 4.10 (m, 1 H), 3.98-3.94 (m, 2 H), 3.94 (s, 3 H), 3.93 (s, 3 H), 3.89 (s, 3 H), 3.88 (s, 3 H), 3.58 (dd, J = 4.2 Hz, 1 H), 3.54 -3.48 (m, 1 H), 3.44-3.39 (m, 1 H), 3.36 (s, 3 H), 3.34-3.27 (m, 2 H), 2.65 (dd, J = 7.8, 16.0 Hz, 1 H), 2.47 (m, 1 H), 2.15 (s, 3 H), 2.04 (m, 1 H), 1.82 (ddd, J = 2.4, 9.5, 12.9 Hz, 1 H), 1.66 (m, 1 H), 1.51 (ddd, J = 3.9, 8.5, 14.2 Hz, 1 H), 1.13 (d, J = 7.1 Hz, 3 H), 0.98 (s, 3 H), 0.94 (m, 2 H), 0.90 (s, 9 H), 0.87 (s, 3 H), 0.85-0.77 (m, 1 H), 0.71-0.65 (m, 1 H), 0.05 (s, 3 H), 0.04 (s, 3 H), 0.004 (s, 9 H), -0.13 (s, 9 H); <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  163.6, 161.3, 160.4, 157.2, 149.6, 149.5, 149.3, 148.8, 142.2, 129.7, 129.0, 120.2, 119.2, 116.3, 111.4, 111.2, 111.0, 110.9, 108.3, 98.0, 93.6, 84.4, 79.4, 77.4, 75.0, 73.5, 71.2, 70.5, 67.6, 65.7, 63.6, 56.3, 56.2, 56.2, 56.1, 56.0, 39.3, 37.7, 31.7, 29.9, 29.3, 26.3, 26.0, 18.2, 18.1, 17.9, 11.4, 9.9, -1.3, -1.4, -4.3, -4.8. high resolution mass spectrum (ES+) m/z 1164.5879 [(M + Na)<sup>+</sup>; calcd for C<sub>59</sub>H<sub>95</sub>NO<sub>15</sub>Si<sub>3</sub>Na 1164.5907].

Amide (+)-50. To a solution of carboxylic acid (-)-3 (31 mg, 0.102 mmol) in CH<sub>2</sub>Cl<sub>2</sub> was added a solution of pyridine (0.66 mL, 0.62 M in CH<sub>2</sub>Cl<sub>2</sub>, 4 equiv) and a solution of thionyl chloride (0.64 mL, 0.48 M in  $CH_2Cl_2$ , 3 equiv), and the mixture was stirred at rt for 2 h. The resulting solution was concentrated under a stream of positive N2 and then placed under a vacuum (~0.1 mmHg). The crude mixture was dissolved in toluene (0.4 mL) and transferred to an oven-dried vial via cannula transfer (flask rinsed with  $2 \times 0.4$  mL toluene). Crude acid chloride 49 was concentrated in vacuo, dissolved in THF (1.0 mL, 0.1 M), and used in the next step without further purification. A solution of carbamate (+)-48 (0.030 mg, 0.026 mmol) in THF (0.65 mL) was cooled to -78 °C, and a solution of *i*-PrMgCl (55 uL, 2.0 M in THF, 4 equiv) was added over 2 min. The yellow solution was stirred for 30 min at -78 °C, and then a solution of acid chloride 49 (1.0 mL, 0.1 M in THF, 3.9 equiv) was added dropwise over 20 min. After 2.5 h, the reaction mixture was quenched with a saturated aq. solution of NaHCO<sub>3</sub> (1.5 mL) and warmed to rt. The aqueous layer was extracted with EtOAc  $(3 \times 2 \text{ mL})$ , and the combined organic layers were washed with brine and then dried over MgSO<sub>4</sub>, filtered and concentrated in vacuo. The crude mixture was purified via flash chromatography on deactivated silica gel (1% v/v triethylamine, 25-30% EtOAc/hexanes) to furnish amide (+)-50 (27 mg, 0.019 mmol, 72%) as a white foam:  $[\alpha]_{D}^{20}$  +27.0 (c 0.3, CHCl3); IR (neat) 2928, 2856, 1716, 1593, 1518, 1464, 1250, 1160, 1083, 1029, 859,

837 cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  7.34 (d, J = 1.9 Hz, 1 H), 6.97–6.93 (m, 3 H), 6.88 (d, J = 8.0 Hz, 1 H), 6.83 (d, J = 8.2 Hz, 1 H), 6.54 (s, 1 H), 5.62 (d, I = 5.7 Hz, 1 H), 5.17 (d, I = 4.4 Hz, 1 H), 5.16 (d, I= 12.7 Hz, 1 H), 5.10 (d, J = 11.9 Hz, 1 H), 4.98 (s, 2 H), 4.75 (s, 1 H), 4.73 (s, 1 H), 4.67 (dd, J = 6.4, 12.7 Hz, 2 H), 4.59 (dd, J = 7.1, 25.3 Hz, 2 H), 4.35 (m, 1 H), 4.32 (dd, J = 3.6, 6.8 Hz, 1 H), 4.30–4.25 (m, 2 H), 3.93 (s, 3 H), 3.91 (s, 3 H), 3.90 (s, 3 H), 3.88 (s, 3 H), 3.83 (m, 1 H), 3.64 (m, 1 H), 3.58 (ddd, J = 1.6, 9.8, 13.1 Hz, 2 H), 3.53 (m, 1 H), 3.47–3.38 (m, 2 H), 3.36 (s, 3 H), 3.31 (s, 3 H), 3.24 (dd, J = 1.8, 17.7 Hz, 1 H), 3.20 (m, 1 H), 2.86 (dd, J = 12.4, 15.9 Hz, 1 H), 2.26 (m, 1 H), 2.25 (m, 1 H), 2.21 (s, 3 H), 2.05 (m, 1 H), 1.96 (ddd, J = 4.0, 4.0, 13.7 Hz, 1 H), 1.86 (m, 2 H), 1.73 (s, 3 H), 1.70 (m, 1 H), 1.14 (d, J = 7.6 Hz, 3 H), 1.09 (dd J = 7.1, 9.2 Hz, 2 H), 0.92 (s, 3 H), 0.90 (s, 9 H), 0.86 (s, 3 H), 0.84 (m, 2 H), 0.78 (ddd, J = 5.7, 11.6, 13.6 Hz, 1 H), 0.66 (ddd, J = 5.5, 11.6, 13.6 Hz, 1 H), 0.06 (s, 3 H), 0.05 (s, 3 H), 0.05 (s, 9 H), -0.05 (s, 9 H), -0.15 (s, 9 H); <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>) δ 175.2, 163.9, 161.2, 160.4, 154.5, 149.5, 149.4, 149.2, 142.9, 148.7 142.6, 129.7, 129.0, 120.2, 119.1, 116.4, 112.8, 111.2, 111.0, 110.8, 110.7, 108.2, 97.7, 95.1, 94.1, 88.5, 81.1, 79.6, 77.4, 77.0, 75.9, 73.0, 71.2, 70.4, 66.3, 66.0, 65.7, 58.4, 57.0, 56.2, 56.1, 56.1, 56.1, 39.8, 39.1, 38.8, 31.8, 31.1, 29.9, 29.9, 26.1, 24.7, 23.1, 18.2, 18.2, 18.1, 17.7, 11.5, 9.8, -1.3, -1.4, -1.4, -4.1, -4.8. HRMS (ES+) m/z 1450.7513 [(M + Na)<sup>+</sup>; calcd for C<sub>73</sub>H<sub>121</sub>NO<sub>19</sub>Si<sub>4</sub>Na 1450.7508].

**Ketone** (–)-**51.** To a solution of amide (+)-**50** (7.0 mg, 0.005 mmol) in THF (0.1 mL) was added a solution of TBAF (15 uL, 1 M in THF, 3.0 equiv). The yellow solution was stirred at rt for 1 h and then warmed to 50 °C. After 19 h, additional TBAF (10 uL, 0.010 mmol, 2.0 equiv) was added. The reaction mixture was stirred at 50 °C for an additional 23 h and then cooled to rt, quenched with water and extracted with EtOAc (4  $\times$  0.5 mL). The combined organic layers were dried over MgSO<sub>4</sub>, filtered and concentrated in vacuo. The crude mixture was purified via flash chromatography on SiO<sub>2</sub> (40% EtOAc/hexanes to 60% EtOAc/ hexanes to 70% EtOAc/hexanes) to furnish alcohol (-)-72 (4.5 mg, 0.004 mmol, 79%) as a colorless oil:  $[\alpha]_D^{20}$  -3.2 (c 0.3, CHCl<sub>3</sub>); IR (neat) 3426, 2951, 2835, 1715, 1685, 1593, 1517, 1265, 1248, 1160, 1028 cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  7.32 (d, J = 1.6 Hz, 1 H), 7.25 (d, J = 9.4 Hz, 1 H), 6.97 - 6.92 (m, 3 H), 6.87 (d, J = 8.8 Hz, 1 H),6.83 (d, J = 8.0 Hz, 1 H), 6.53 (s, 1 H), 5.16 (dab, J = 11.7 Hz, 1 H), 5.09 (dab, J = 11.8 Hz, 1 H), 5.09 (dd, J = 2.5, 5.6 Hz, 1 H), 4.98 (s, 2 H), 4.83 (d, I = 6.6 Hz, 1 H), 4.77 (s, 1 H), 4.76 (s, 1 H), 4.71 (dab, I = 6.3 Hz, 1H), 4.70 (dab, J = 6.8 Hz, 1 H), 4.60 (d, J = 7.7 Hz, 1 H), 4.40 (d, J = 2.1 Hz, 1 H), 4.27 (ddd, J = 1.9, 7.8, 11.9 Hz, 1 H), 4.08 (m, 1 H), 3.99 (m, 1 H), 3.93 (s, 3 H), 3.90 (s, 3 H), 3.89 (s, 3 H), 3.87 (s, 3 H), 3.74 (m, 2 H), 3.67 (dd, J = 4.5, 4.5 Hz, 1 H), 3.58 (ddd, J = 6.5, 10.3, 10.3 Hz, 1 H), 3.50–3.43 (m, 2 H), 3.37 (s, 3 H), 3.32 (s, 3 H), 3.27 (dd, J = 2.2, 16.7 Hz, 1 H), 2.65 (dd, J = 12.7, 16.4 Hz, 1 H), 2.46 (m, 1 H), 2.37 (dd, J = 8.5, 14.5 Hz, 1 H), 2.19 (dd, J = 4.7, 14.8 Hz, 1 H), 2.14 (s, 3 H), 2.10 (ddd, J = 2.6, 8.4, 8.4 Hz, 1 H), 2.01 (m, 1 H), 1.81 (ddd, J = 2.54, 9.4, 13.5 Hz, 1 H), 1.73 (m, 1 H), 1.70 (s, 3 H), 1.56 (dd, *J* = 4.7, 9.6 Hz, 1 H), 1.54 (dd, J = 5.0, 9.3 Hz, 1 H), 1.15 (d, J = 6.7 Hz, 3 H), 1.00 (s, 3 H), 0.94 (s, 3 H), 0.91 (m, 2 H), 0.85–0.79 (m, 1 H), 0.71–0.69 (m, 1 H), 0.01 (s, 9 H), -0.13 (s, 9 H); <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  171.2, 163.9, 161.3, 160.4, 149.5, 149.4, 149.2, 148.7, 142.4, 142.2, 129.6, 128.9, 120.2, 119.2, 116.7, 113.0, 111.2, 111.0, 110.9, 110.7, 108.1, 97.7, 94.8, 94.4, 81.8, 81.4, 79.6, 77.4, 75.5, 72.9, 71.1, 70.5, 68.0, 66.2, 65.6, 58.0, 56.3, 56.2, 56.2, 56.1, 56.1, 39.2, 38.3, 37.3, 30.7, 30.0, 29.9, 29.5, 26.0, 22.9, 19.4, 18.2, 18.1, 11.4, 9.6, -1.2, -1.4; HRMS (ES+) m/z1192.6072 [ $(M + Na)^+$ ; calcd for  $C_{61}H_{95}NO_{17}Si_2Na$  1192.6036]

To a solution of alcohol (-)-72 (3.5 mg, 0.003 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (0.05 mL) was added NaHCO<sub>3</sub> (4.2 mg, 16.6 equiv). The reaction mixture was cooled to 0 °C, and Dess–Martin periodinane (6.0 mg, 0.015 mmol, 5.0 equiv) was added, and the resulting mixture was stirred for 3 h. Reaction was quenched with a saturated aq. solution of NaHCO<sub>3</sub> and extracted with EtOAc (4 × 0.5 mL). The combined organic layers were dried over MgSO<sub>4</sub>, filtered and concentrated in vacuo. The crude mixture was purified via flash chromatography on SiO<sub>2</sub> (40% EtOAc/hexanes to 50% EtOAc/hexanes) to furnish ketone (-)-**51** (3.0 mg, 0.0026 mmol, 87%) as a colorless oil:  $[\alpha]_{20}^{20}$  –14.5 (*c* 0.2, CHCl<sub>3</sub>); IR (neat) 3403, 2951, 2928, 2835, 1713, 1687, 1593, 1517, 1463, 1265, 1248, 1159, 1080.9, 1029 cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  7.31 (ap

s, 1 H), 7.27 (ap s, 1 H), 6.97 (d, J = 8.4 Hz, 1 H), 6.93 (m, 2 H), 6.88 (d, J = 8.4 Hz, 1 H), 6.83 (d, J = 8.2 Hz, 1 H), 6.55 (s, 1 H), 5.18 (dab, J =12.0 Hz, 1 H), 5.11 (dab, I = 11.7 Hz, 1 H), 5.09 (d, I = 9.8 Hz, 1 H), 4.98 Hz(s, 2 H), 4.83 (dab, J = 6.4 Hz, 1 H), 4.81 (s, 1 H), 4.79 (s, 1 H), 4.73 (dab, J = 6.6 Hz, 1 H), 4.65 (dab, J = 7.0 Hz, 1 H), 4.61 (dab, J = 7.0 Hz, 1 H), 4.38 (d, J = 2.0 Hz, 1 H), 4.29–4.23 (m, 2 H), 3.99 (m, 1 H), 3.93 (s, 3 H), 3.90 (s, 3 H), 3.89 (s, 3 H), 3.88 (s, 3 H), 3.75 (m, 1 H), 3.72 (m, 1 H), 3.56 (ddd, J = 6.4, 10.0, 10.0 Hz, 1 H), 3.51-3.42 (m, 2 H), 3.40 (s, 3 H), 3.35 (aps, 1 H), 3.34 (s, 3 H), 3.21 (d, J = 15.9 Hz, 1 H), 2.68 (dd, J = 12.7, 16.4 Hz, 1 H), 2.60 (dd, J = 11.4, 14.9 Hz, 1 H), 2.38 (dd, J = 8.9, 14.9 Hz, 1 H), 2.27 (dd, J = 3.8, 10.9, 1 H), 2.24 (dd, J = 5.2, 12.3, 1 H), 2.14 (s, 3 H), 2.17 – 2.09 (m, 1 H), 1.86–1.75 (m, 2 H), 1.73 (s, 3 H), 1.25 (s, 3 H), 1.16 (d, J = 6.6 Hz, 3 H), 1.01 (s, 3 H), 0.96-0.86 (m, 2 H), 0.82-0.76 (m, 1 H), 0.74-0.68 (m, 1 H), 0.01 (s, 9 H), -0.10 (s, 9 H); <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>) δ 210.8, 171.4, 163.7, 161.4, 160.4, 149.5, 149.4, 149.3, 148.7, 142.2, 141.8, 129.5, 128.8, 120.3, 119.2, 116.1, 113.3, 111.2, 111.0, 110.9, 110.7, 108.0, 97.8, 95.0, 94.8, 81.7, 81.4, 79.8, 79.2, 77.6, 74.6, 72.8, 71.1, 70.5, 66.3, 65.8, 58.1, 56.4, 56.2, 56.2, 56.1, 56.1, 49.6, 39.6, 38.8, 38.3, 30.2, 29.9, 24.8, 23.0, 19.5, 18.2, 18.1, 11.4, 9.7, -1.2, -1.4; high resolution mass spectrum (ES+) m/z 1190.5880 [(M + Na)<sup>+</sup>; calcd for  $C_{61}H_{93}NO_{17}Si_2Na$  1190.5880]

(-)-Irciniastatin B (2). To a solution of fully protected irciniastatin B (-)-51 (5.0 mg, 0.0043 mmol) in  $CH_2Cl_2$  (0.05 mL) and  $H_2O$  (15 uL) was added a suspension of 2,3-dichloro-5,6-dicyano-1,4-benzoquinone (0.1 mL, 0.33 M in  $CH_2Cl_2$ , 8 equiv). After 24 h, the reaction mixture was quenched with a saturated aq. solution of NaHCO<sub>3</sub> and extracted with EtOAc (5 × 0.5 mL). The combined organic layers were dried over MgSO<sub>4</sub>, filtered and concentrated in vacuo. The crude mixture was purified via flash chromatography (40% EtOAc/hexanes) to afford a mixture (1:2) of desired bis-phenol and 3,4-dimethoxybenza-lehyde, respectively.

The mixture was treated with a stock solution of MgBr<sub>2</sub>/n-BuSH/ MeNO<sub>2</sub> in Et<sub>2</sub>O (0.21 mL: 25 equiv MgBr<sub>2</sub>, 25 equiv n-BuSH, stock solution made up of 75.4 mg of MgBr<sub>2</sub>, 44  $\mu$ L of *n*-BuSH, 82  $\mu$ L of MeNO<sub>2</sub> and 0.82 mL of Et<sub>2</sub>O). After 10 h, the reaction mixture was diluted with EtOAc, quenched with a saturated aq. solution of NaHCO<sub>3</sub> and extracted with EtOAc ( $5 \times 0.5$  mL). The combined organic layers were dried over MgSO<sub>4</sub>, filtered and concentrated in vacuo. The crude mixture was purified via flash chromatography with water-washed SiO<sub>2</sub>  $[50 \text{ g of SiO}_2 \text{ washed with H}_2O (500 \text{ mL}), \text{ then MeOH} (500 \text{ mL}), \text{ then}$ EtOAc (500 mL), then hexanes (500 mL), and dried under a vacuum overnight, and then deactivated with 5% v/v triethylamine, 35% EtOAc/ hexanes to 80% EtOAc/hexanes] to afford (-)-irciniastatin B (2) (2.0 mg, 0.0033 mmol, 78% over two steps) as a colorless solid:  $\left[\alpha\right]_{\rm D}^{20}$  -28.7 (c 0.2, MeOH) IR (neat) 3356, 2925, 2873, 1710, 1651, 1612, 1510, 1461, 1380, 1266, 1174, 1103 cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$ 11.11 (s, 1 H), 7.37 (d, J = 10.4 Hz, 1 H), 6.64 (bs, 1 H), 6.30 (s, 1 H), 5.20 (dd, J = 6.4 Hz, 10.1 Hz, 1 H), 4.82 (s, 1 H), 4.79 (s, 1 H), 4.55 (ddd, J = 4.2, 4.2, 11.8 Hz, 1 H), 4.47 (ap t, J = 2.9 Hz, 1 H), 4.21 (ap q, J = 6.4 Hz, 1 H), 4.09 (d, J = 8.4 Hz, 1 H), 4.00 (dd, J = 1.8, 11.2 Hz, 1 H), 3.79-3.77 (m, 1 H), 3.77 (bs, 1H), 3.65 (bs, 1 H), 3.39 (s, 3 H), 3.36 (s, 3 H), 2.94–2.83 (m, 2 H), 2.67 (ap d, J = 6.4 Hz, 2 H), 2.36 (dd, J = 9.4, 14.5 Hz, 1 H), 2.14 (dd, J = 3.7, 14.5 Hz, 1 H), 2.04 (s, 3 H), 1.91 (1 H, m), 1.84 (ddd, J = 10.1, 14.6, 24.8 Hz, 1 H), 1.75 (s, 3 H), 1.59 (ap d, J = 10.1, 14.6, 24.8 Hz, 1 H), 1.75 (s, 3 H), 1.59 (ap d, J = 10.1, 14.6, 24.8 Hz, 1 H), 1.75 (s, 3 H), 1.59 (ap d, J = 10.1, 14.6, 24.8 Hz, 1 H), 1.75 (s, 3 H), 1.59 (ap d, J = 10.1, 14.6, 24.8 Hz, 1 H), 1.75 (s, 3 H), 1.59 (ap d, J = 10.1, 14.6, 24.8 Hz, 1 H), 1.75 (s, 3 H), 1.59 (ap d, J = 10.1, 14.6, 24.8 Hz, 1 H), 1.75 (s, 3 H), 1.59 (ap d, J = 10.1, 14.6, 24.8 Hz, 1 H), 1.75 (s, 3 H), 1.59 (ap d, J = 10.1, 14.6, 24.8 Hz, 1 H), 1.75 (s, 3 H), 1.59 (ap d, J = 10.1, 14.6, 24.8 Hz, 1 H), 1.75 (s, 3 H), 1.59 (ap d, J = 10.1, 14.6, 24.8 Hz, 14.8 Hz, 1414.0 Hz, 1 H), 1.16 (s, 3 H), 1.11 (d, J = 7.2 Hz, 3 H), 1.10 (s, 3H); <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>) δ 210.3, 173.2, 170.7, 162.5, 161.3, 142.1, 139.9, 113.5, 113.3, 101.7, 101.5, 83.2, 80.5, 80.5, 80.3, 73.8, 72.7, 72.4, 57.9, 56.6, 49.6, 42.8, 38.8, 37.4, 33.0, 28.3, 22.8, 22.3, 19.4, 10.7, 9.2; HRMS (ES+) m/z 608.3058 [(M + 1)<sup>+</sup>; calcd for C<sub>31</sub>H<sub>46</sub>NO<sub>11</sub> 608.3071]

(+)-Irciniastatin A (1) and *epi*-C(11)-Irciniastatin A (52). Neat (-)-irciniastatin B (2) (1 mg, 1.6  $\mu$ mol) was treated with a solution of NaBH<sub>4</sub> (0.1 mL, 0.024 M in MeOH, 1.5 equiv) at 0 °C. After 15 min, the reaction mixture was quenched with a saturated aq. solution of NaHCO<sub>3</sub> (0.4 mL) and extracted with EtOAc (3 × 0.5 mL). The combined organic layers were dried over MgSO<sub>4</sub>, filtered and concentrated in vacuo. The crude mixture (1:1) of (+)-1 and 52 was purified via preparatory TLC (70% EtOAc/hexanes, 250  $\mu$ m SiO<sub>2</sub> plate) to provide (+)-irciniastatin A (1) (0.3 mg, 0.5  $\mu$ mol 31%) and *epi*-C(11)-irciniastatin A (52) (0.3 mg, 0.5  $\mu$ mol, 31%). All spectral data of

(+)-irciniastatin A (1) matched to the synthetic sample from our firstgeneration approach. Characterization data for (+)-irciniastatin A (1): <sup>1</sup>H NMR (500 MHz, MeOD)  $\delta$  6.24 (s, 1 H), 5.39 (d, J = 8.3 Hz, 1 H), 4.74 (s, 1 H), 4.71 (s, 1 H), 4.51–4.47 (ddd, *I* = 3.0, 5.9, 12.0 Hz, 1 H), 4.35 (d, J = 2.6 Hz, 1 H), 3.94 (m, 2 H), 3.67 (ddd, J = 2.6, 3.5, 9.5 Hz, 1 H), 3.60 (dd, J = 4.4, 10.9 Hz, 1 H), 3.50 (dd, J = 2.0, 10.1 Hz, 1 H), 3.35 (s, 3 H), 3.23 (s, 3 H), 3.13 (dd, J = 3.3, 16.7 Hz, 1 H), 2.86 (dd, J = 12.0, 16.6 Hz, 1 H), 2.35 (dd, J = 9.4, 14.8 Hz, 1 H), 2.11 (m, 1 H), 2.10 (s, 3 H), 2.02 (ddd, J = 2.6, 4.5, 13.4 Hz, 1 H), 1.91 (m, 1 H), 1.86–1.74 (m, 2 H), 1.72 (s, 3 H), 1.68 (ddd, J = 2.1, 3.8, 14.6 Hz, 1 H), 1.10 (d, J = 7.1 Hz, 3 H), 0.97 (s, 3 H), 0.90 (s, 3 H); <sup>13</sup>C NMR (125 MHz, MeOD) Observable peaks & 176.3, 172.5, 163.9, 144.0, 141.2, 115.5, 113.1, 101.6, 82.8, 82.3, 82.1, 79.9, 73.6, 73.3, 72.1, 57.8 56.7, 43.4, 39.9, 38.8, 34.5, 30.6, 29.6, 23.8, 23.0, 14.0, 11.0, 9.3; HRMS (ES+) m/z 632.3033 [(M+ 1)<sup>+</sup>; calcd for  $C_{31}H_{47}NO_{11}Na$  632.3047]. Characterization data for *epi*-C(11)-irciniastatin A (52): <sup>1</sup>H NMR (500 MHz, MeOD) Observable peaks  $\delta$  6.25 (s, 1 H), 5.25 (d, J = 3.6 Hz, 1 H), 4.76 (s, 1 H), 4.73 (s, 1 H), 4.51–4.47 (ddd, J = 3.1, 6.8, 12.2 Hz, 1 H), 4.37 (d, J = 2.8 Hz, 1 H), 4.04 (m, 1 H), 3.97 (dd, J = 4.0, 13.2 Hz, 1 H), 3.78 (dd, J = 3.1, 11.8 Hz, 1 H), 3.73–3.67 (m, 2 H), 3.33 (s, 3 H), 3.17 (dd, *J* = 3.3, 16.7 Hz, 1 H), 2.83 (dd, J = 11.9, 16.2 Hz, 1 H), 2.35 (dd, J = 9.7, 15.0 Hz, 1 H), 2.12 (dd, J = 3.9, 14.2 Hz, 1 H), 2.09 (s, 3 H), 1.96 (m, 2 H), 1.79 (m, 1 H), 1.73 (s, 3 H), 1.63 (m, 2 H), 1.12 (d, J = 7.0 Hz, 3 H), 1.01 (s, 3 H), 0.93 (s, 3 H); <sup>13</sup>C NMR (125 MHz, MeOD) Observable peaks δ 176.0, 172.6, 164.7, 163.8, 144.0, 141.2, 115.4, 113.2, 101.5, 101.5, 83.0, 82.5, 73.0, 72.7, 72.0, 57.9, 56.7, 42.8, 39.0, 38.7, 30.9, 29.6, 23.1, 22.8, 21.3, 10.9, 9.7; HRMS (ES+) m/z 632.3029 [(M + 1)<sup>+</sup>; calcd for C<sub>31</sub>H<sub>47</sub>NO<sub>11</sub>Na 632.3047].

### ASSOCIATED CONTENT

# **Supporting Information**

<sup>1</sup>H and <sup>13</sup>C spectral data of all compounds. This material is available free of charge via the Internet at http://pubs.acs.org.

### AUTHOR INFORMATION

#### Corresponding Author

\*E-mail: smithab@sas.upenn.edu.

#### Notes

The authors declare no competing financial interest.

## ACKNOWLEDGMENTS

Financial support was provided by the National Institute of Health (Institute of General Medical Sciences) through Grant No. CA-19033. We thank Drs. George Furst and Rakesh Kohli (University of Pennsylvania) for help in obtaining the high resolution NMR and mass spectral data, respectively. The authors thank the generous help of Professor G. R. Pettit (Arizona State University) for providing the <sup>1</sup>H and <sup>13</sup>C NMR spectra for (-)-irciniastatin B.

#### REFERENCES

(1) Pettit, G. R.; Xu, J.-P.; Chapuis, J.-C.; Pettit, R. K.; Tackett, L. P.; Doubek, D. L.; Hooper, J. N. A.; Schmidt, J. M. *J. Med. Chem.* **2004**, *47*, 1149–1152.

(2) Cichewicz, R. H.; Valeriote, F. A.; Crews, P. Org. Lett. 2004, 6, 1951–1954.

(3) Jiang, X.; García-Fortanet, J.; De Brabander, J. K. J. Am. Chem. Soc. **2005**, *127*, 11254–11255.

(4) Feng, Y.; Jiang, X.; De Brabander, J. K. J. Am. Chem. Soc. **2012**, 134, 17083–17093.

(5) Huang, X.; Shao, N.; Huryk, R.; Palani, A.; Aslanian, R.; Seidel-Dugan, C. Org. Lett. **2009**, *11*, 867–870.

(6) Chinen, T.; Nagumo, Y.; Watanabe, T.; Imaizumi, T.; Shibuya, M.; Kataoka, T.; Kanoh, N.; Iwabuchi, Y.; Usui, T. *Toxicol. Lett.* **2010**, *199*, 341–346.

- (7) Wu, C.-Y.; Feng, Y.; Cardenas, E. R.; Williams, N.; Floreancig, P. E.; De Brabander, J. K.; Roth, M. G. *J. Am. Chem. Soc.* **2012**, *134*, 18998–19003.
- (8) Huang, X.; Shao, N.; Palani, A.; Aslanian, R.; Buevich, A. Org. Lett. 2007, 9, 2597–2600.
- (9) Smith, A. B., III; Jurica, J. A.; Walsh, S. P. Org. Lett. 2008, 10, 5625–5628.
- (10) Crimmins, M. T.; Stevens, J. M.; Schaaf, G. M. Org. Lett. **2009**, *11*, 3990–3993.
- (11) Watanabe, T.; Imaizumi, T.; Chinen, T.; Nagumo, Y.; Shibuya,
- M.; Usui, T.; Kanoh, N.; Iwabuchi, Y. Org. Lett. **2010**, *12*, 1040–1043. (12) Byeon, S. R.; Park, H.; Kim, H.; Hong, J. Org. Lett. **2011**, *13*, 5816–5819.
- (13) Wan, S.; Wu, F.; Rech, J. C.; Green, M. E.; Balachandran, R.; Horne, W. S.; Day, B. W.; Floreancig, P. E. *J. Am. Chem. Soc.* **2011**, *133*, 16668–16679.
- (14) An, C.; Hoye, A. T.; Smith, A. B., III Org. Lett. 2012, 14, 4350–4353.
- (15) Smith, A. B., III; Safonov, I. G.; Corbett, R. M. J. Am. Chem. Soc. **2002**, 124, 11102–11113.
- (16) Yamamoto, K.; Suzuki, S.; Tsuji, J. Chem. Lett. 1978, 7, 649-652.
- (17) Byrson, T. A.; Dolak, T. M. Org. Synth. 1977, 57, 62.
- (18) Forbes, J. E.; Pattenden, G. Tetrahedron Lett. 1987, 28, 2771–2774.
- (19) Ebenezer, W.; Pattenden, G. Tetrahedron Lett. 1992, 33, 4053–4056.
- (20) Hunter, T. J.; O'Doherty, G. A. Org. Lett. 2002, 4, 4447–4450.
  (21) Smith, A. B., III; Walsh, S. P.; Frohn, M.; Duffey, M. O. Org. Lett.
  2004, 7, 139–142.
- (22) Parikh, J. R.; Doering, W. E. J. Am. Chem. Soc. 1967, 89, 5505-5507.
- (23) Bal, B. S.; Childers, W. E.; Pinnick, H. W. Tetrahedron 1981, 37, 2091–2096.
- (24) Langer, P.; Kracke, B. Tetrahedron Lett. 2000, 41, 4545-4547.
- (25) Duan, J. J. W.; Smith, A. B., III J. Org. Chem. 1993, 58, 3703-3711.
- (26) Huang, Z.; Negishi, E.-i. Org. Lett. 2006, 8, 3675-3678.
- (27) Fischetti, W.; Mak, K. T.; Stakem, F. G.; Kim, J.; Rheingold, A. L.; Heck, R. F. J. Org. Chem. **1983**, 48, 948–955.
- (28) Wang, Z.-X.; Tu, Y.; Frohn, M.; Zhang, J.-R.; Shi, Y. J. Am. Chem. Soc. 1997, 119, 11224–11235.
- (29) Ishibashi, N.; Miyazawa, M.; Miyashita, M. *Tetrahedron Lett.* **1998**, *39*, 3775–3778.
- (30) Katsuki, T.; Sharpless, K. B. J. Am. Chem. Soc. **1980**, 102, 5974–5976.
- (31) Oshima, M.; Yamazaki, H.; Shimizu, I.; Nisar, M.; Tsuji, J. J. Am. Chem. Soc. **1989**, 111, 6280–6287.
- (32) Zhao, M. M.; Li, J.; E, M.; Song, Z. J.; Taschaen, D. M. Org. Synth. 2005, 195.
- (33) Baldwin, J. E. J. Chem. Soc., Chem. Commun. 1976, 734-736.
- (34) Dess, D. B.; Martin, J. C. J. Org. Chem. 1983, 48, 4155-4156.
- (35) Hamana, H.; Sasakura, K.; Sugasawa, T. Chem. Lett. 1984, 13, 1729-1732.
- (36) Evans, D. A.; Calter, M. A. Tetrahedron Lett. 1993, 34, 6871-6874.
- (37) Chen, K.-M.; Hardtmann, G. E.; Prasad, K.; Repič, O.; Shapiro, M. J. *Tetrahedron Lett.* **1987**, *28*, 155–158.
- (38) Noyori, R.; Nishida, I.; Sakata, J.; Nishizawa, M. J. Am. Chem. Soc. **1980**, 102, 1223–1225.
- (39) Scheidt, K. A.; Chen, H.; Follows, B. C.; Chemler, S. R.; Coffey, S.; Roush, W. R. J. Org. Chem. **1998**, 63, 6436–6437.
- (40) Vakalopoulos, A.; Hoffmann, H. M. R. Org. Lett. 2000, 2, 1447–1450.
- (41) Simsek, S.; Horzella, M.; Kalesse, M. Org. Lett. 2007, 9, 5637–5639.
- (42) Ohtani, I.; Kusumi, T.; Kashman, Y.; Kakisawa, H. J. Am. Chem. Soc. **1991**, 113, 4092–4096.
- (43) Hoye, T. R.; Jeffrey, C. S.; Shao, F. Nat. Protoc. 2007, 2, 2451–2458.

- (44) Paterson, I.; Goodman, J. M. Tetrahedron Lett. 1989, 30, 997–1000.
- (45) Paterson, I.; Goodman, J. M.; Anne Lister, M.; Schumann, R. C.; McClure, C. K.; Norcross, R. D. *Tetrahedron* **1990**, *46*, 4663–4684.
- (46) For an example of base-catalyzed epimerization of *trans*tetrahydropyran, see: Allevi, P.; Anastasia, M.; Ciuffreda, P.; Fiecchi, A.; Scala, A. J. Chem. Soc., Perkin Trans. 1 **1989**, 1275–1280.
- (47) For an example of acid catalyzed hydrolysis of *N*,*O*-aminal, see: Mori, M.; Uozumi, Y.; Ban, Y. *J. Chem. Soc., Chem. Commun.* **1986**, 841–842.
- (48) Denmark, S. E.; Kobayashi, T.; Regens, C. S. Tetrahedron 2010, 66, 4745–4759.