

α -Aryl-Substituted Allenamides in an Imino-Nazarov Cyclization Cascade Catalyzed by Au(I)


Zhi-Xiong Ma, Shuzhong He, Wangze Song, and Richard P. Hsung*

Division of Pharmaceutical Sciences, School of Pharmacy, University of Wisconsin,
Madison, Wisconsin 53705, United States

rhsung@wisc.edu

Received October 4, 2012

ABSTRACT



An imino-Nazarov cyclization using α -aryl-substituted allenamides is described. This gold(I)-catalyzed cascade is efficient and regioselective in constructing a diverse array of synthetically useful aromatic-ring fused cyclopentenamides. The success in this transformation represents a solution to the challenge in establishing an imino-Nazarov cyclization process.

The Nazarov cyclization¹ has captured an immense amount of interest from the synthetic community in the past three decades.^{2,3} This classical pericyclic process


(1) Nazarov, I. N.; Torgov, I. B.; Terekhova, L. N. *Izv. Akad. Nauk SSSR, Otd. Khim. Nauk* **1942**, 200.

(2) For earlier reviews, see: (a) Habermas, K. L.; Denmark, S. E.; Jones, T. K. In *Organic Reactions*; Paquette, L. A., Ed.; John Wiley & Sons: New York, 1994; Vol. 45, pp 1–158. (b) Denmark, S. E. In *Comprehensive Organic Synthesis*; Trost, B. M., Fleming, I., Eds.; Pergamon: Oxford, U.K., 1991; Vol. 5, pp 751–784. (c) Santelli-Rouvier, C.; Santelli, M. *Synthesis* **1983**, 429.

(3) For recent reviews, see: (a) Vaidya, T.; Eisenberg, R.; Frontier, A. J. *ChemCatChem* **2011**, 3, 1531. (b) Shimada, N.; Stewart, C.; Tius, M. A. *Tetrahedron* **2011**, 67, 5851. (c) Grant, T. N.; Rieder, C. J.; West, F. G. *Chem. Commun.* **2009**, 45, 5676. (d) Nakanishi, W.; West, F. *Curr. Opin. Drug Discovery Dev.* **2009**, 12, 732. (e) Frontier, A.; Collison, C. *Tetrahedron* **2005**, 61, 7577. (f) Tius, M. A. *Eur. J. Org. Chem.* **2005**, 2193. (g) Pellissier, H. *Tetrahedron* **2005**, 61, 6479. (h) Tius, M. A. *Acc. Chem. Res.* **2003**, 36, 284. (i) Harmata, M. *Chemtracts* **2004**, 17, 416.


(4) Given the large volume, for some reports in 2012 on Nazarov cyclizations, see: (a) Shimada, N.; Stewart, C.; Bow, W. F.; Jolit, A.; Wong, K.; Zhou, Z.; Tius, M. A. *Angew. Chem., Int. Ed.* **2012**, 51, 5727. (b) Lebuf, D.; Gandon, V.; Ciesielski, J.; Frontier, A. J. *J. Am. Chem. Soc.* **2012**, 134, 6296. (c) Kerr, D. J.; Flynn, B. L. *Org. Lett.* **2012**, 14, 1740. (d) Guo, F.; Wang, L.; Mao, S.; Zhang, C.; Yu, J.; Han, J. *Tetrahedron* **2012**, 68, 8367. (e) Alvarez, E.; Miguel, D.; Garcia-Garcia, P.; Fernández-Rodríguez, M. A.; Rodríguez, F.; Sanz, R. *Synthesis* **2012**, 1874. (f) Raja, S.; Ieawsuwan, W.; Korotkov, V.; Rueping, M. *Chem.—Asian J.* **2012**, 7, 2361. (g) Dateer, R. B.; Pati, K.; Liu, R.-S. *Chem. Commun.* **2012**, 48, 7200. (h) Gan, Z.; Wu, Y.; Gao, L.; Sun, X.; Lei, J.; Song, Z.; Li, L. *Tetrahedron* **2012**, 68, 6928. (i) Also see refs 11 and 12.

Scheme 1. Nazarov Cyclizations: Challenge in an Imino Version



proceeds through a conrotatory 4π -electron electrocyclic ring closure of a pentadienyl cation via activation of divinyl ketone **1** or its equivalent, leading to a key oxyallyl cation **2** that can afford cyclopentenone **3** or substituted cyclopentanones **4** through trapping [Scheme 1]. Despite a diverse array of innovative structural designs that have been developed,^{4–6} one aspect that has received very little attention is an imino version of this cascade with the sole exceptions of Tius' seminal work on lithiated imino-Nazarov

Scheme 2. An Approach to Imino-Nazarov Cyclization



reactions^{7a} and, recently, with enamine-iminium ions.^{7b,c} There are examples in which stabilization of the pentadienyl cation is assisted by the presence of an amino or amide substituent, albeit not a formal imino-Nazarov cyclization. Cha's⁸ earlier work on cephalotaxine synthesis and reports by Occhiato/Prandi,⁹ Frontier,¹⁰ and Flynn¹¹ employing ynamides, and recently by West¹² in a clever usage of allenamides, represent such elegant cases. The challenge in developing a successful imino-Nazarov cyclization is to overcome the propensity of the 2-amino-allyl cation **6** to ring open in favor of amino-pentadienyl cation **5**, which receives greater stabilization from the amino nitrogen atom.¹³ We wish to communicate here our solution to this challenge through gold(I) activation of allenamides.

Our long-standing interest in allenamides,^{14–16} and in particular their electrophilic activations,^{17,18} led us to explore their potential use in this underdeveloped aspect of the Nazarov cyclization. As shown in Scheme 2, α -vinyl or

α -aryl allenamides **7** could be employed in an imino-Nazarov cyclization cascade through Brønsted acid activation.¹⁹ This electrophilic activation could give amido-pentadienyl cation **8**, which would undergo an electrocyclic cyclization to afford 2-amido-allyl cation **9**. It is noteworthy that a distinct advantage in achieving a successful imino-Nazarov cyclization is the synthetically useful enamide functionality²⁰ in the end product **10**.

We reasoned that unlike amino-pentadienyl cation **5**, the electrocyclic cyclization of **7** could be more favored given the reduced ability of an *N*-acyl nitrogen atom to stabilize cations relative to an amino group. Such rationalization would find prevalent support in the related chemistry of vinyl *N*-acyl iminium ions²¹ and *aza*-Prins cyclizations.²² However, after many attempts, the Brønsted acid activation failed because we were unable to control and overtake the hydrolysis pathway

(15) Also see: (a) Standen, P. E.; Kimber, M. C. *Curr. Opin. Drug Discovery Dev.* **2010**, *13*, 645. (b) Deagostino, A.; Prandi, C.; Tabasso, S.; Venturello, P. *Molecules* **2010**, *15*, 2667. (c) For general reviews on allenes, see: Krause, N.; Hashmi, A. S. K. *Modern Allene Chemistry*; Wiley-VCH Verlag GmbH & Co. KGaA: Weinheim, 2004; Vols. 1 and 2.

(16) Given the large volume, for reports on allenamide chemistry just in 2012, see: (a) Reference 12. (b) Du, Y.; Krenske, E. H.; Antoline, J. E.; Lohse, A. G.; Houk, K. N.; Hsung, R. P. *J. Org. Chem.* **2012**, ASAP Article (DOI:10.1021/jo3011792). (c) Singh, S.; Elsegood, M. R. J.; Kimber, M. C. *Synlett* **2012**, 23, 565. (d) Heffernan, S. J.; Carbery, D. R. *Tetrahedron Lett.* **2012**, 53, 5180. (e) Hayashi, R.; Ma, Z.-X.; Hsung, R. P. *Org. Lett.* **2012**, 14, 252. (f) X.-X.; Zhu, L.-L.; Zhou, W.; Chen, Z. *Org. Lett.* **2012**, 14, 436. (g) Husinec, S.; Petkovic, M.; Savić, V.; Simić, M. *Synthesis* **2012**, 399. (h) Suárez-Pantiga, S.; Hernández-Díaz, C.; Piedrafita, M.; Rubio, E.; González, J. M. *Adv. Synth. Catal.* **2012**, 354, 1651. (i) Faustino, H.; Paloma Bernal, P.; Luis Castedo, L.; Fernando López, F.; Mascareñas, J. L. *Adv. Synth. Catal.* **2012**, 354, 1658. (j) Francos, J.; Grande-Carmona, F.; Faustino, H.; Iglesias-Sigüenza, J.; Díez, E.; Alonso, I.; Fernández, R.; Lassaletta, J. M.; López, F.; Mascareñas, J. L. *J. Am. Chem. Soc.* **2012**, 134, 14322. (k) Brogini, G.; Borsini, E.; Fasana, A.; Poli, G.; Liron, F. *Eur. J. Org. Chem.* **2012**, 3617.

(17) For an account, see: Lohse, A. G.; Hsung, R. P. *Chem.—Eur. J.* **2011**, 17, 3812.

(18) (a) Via oxidation: Xiong, H.; Hsung, R. P.; Berry, C. R.; Rameshkumar, C. *J. Am. Chem. Soc.* **2001**, 123, 7174. (b) Via Brønsted acid: Berry, C. R.; Hsung, R. P.; Antoline, J. E.; Petersen, M. E.; Rameshkumar, C.; Nielson, J. A. *J. Org. Chem.* **2005**, 70, 4038. (c) Via halogen: Hayashi, R.; Walton, M. C.; Hsung, R. P.; Schwab, J.; Yu, X. *Org. Lett.* **2010**, 12, 5768.

(19) For a Brønsted acid activation of allenol ethers, see: (a) Wu, Y.-K.; West, F. G. *J. Org. Chem.* **2010**, 75, 5410. For an oxidative activation, see: (b) Spencer, W. T., III; Levin, M. D.; Frontier, A. J. *Org. Lett.* **2011**, 13, 414. (c) Malona, J. A.; Cariou, K.; Frontier, A. J. *J. Am. Chem. Soc.* **2009**, 131, 7560. For a gold activation, see: (d) Zhang, L.; Wang, S. *J. Am. Chem. Soc.* **2006**, 128, 1442.

(20) For a leading review on enamides, see: (a) Carbery, D. R. *Org. Biomol. Chem.* **2008**, 9, 3455. Also see: (b) Rappoport, Z. *The Chemistry of Functional Groups*; John Wiley and Sons: New York, 1994. (c) (b) Tracey, M. R.; Hsung, R. P.; Antoline, J.; Kurtz, K. C. M.; Shen, L.; Slafer, B. W.; Zhang, Y. In *Science of Synthesis, Houben-Weyl Methods of Molecular Transformations*; Weinreb, S. M., Ed.; Georg Thieme Verlag KG: Chapter 21.4, 2005.

(21) For reviews, see: (a) Maryanoff, B. E.; Zhang, H.-C.; Cohen, J. H.; Turchi, I. J.; Maryanoff, C. *Chem. Rev.* **2004**, 104, 1431. (b) Royer, J.; Bonin, M.; Micouin, L. *Chem. Rev.* **2004**, 104, 2311.

(22) For reviews, see: Hiemstra, H.; Speckamp, W. N. In *Comprehensive Organic Synthesis*; Brossi, A., Ed.; Academic Press: San Diego, 1988; Vol. 2, pp 271–339. (b) Hanessian, S.; Tremblay, M.; Marzi, M.; Del Valle, J. R. *J. Org. Chem.* **2005**, 70, 5070.

(23) For reviews, see: (a) Boorman, T. C.; Larrosa, I. *Chem. Soc. Rev.* **2011**, 40, 1910. (b) Bandini, M. *Chem. Soc. Rev.* **2011**, 40, 1358. (c) Krause, N.; Winter, C. *Chem. Rev.* **2011**, 111, 1994. (d) Fürstner, A. *Chem. Soc. Rev.* **2009**, 38, 3208. (e) Gorin, D. J.; Sherry, B. D.; Toste, F. D. *Chem. Rev.* **2008**, 108, 3351. (f) Fürstner, A.; Davies, P. W. *Angew. Chem., Int. Ed.* **2007**, 46, 3410. (g) Hashmi, A. S. K. *Chem. Rev.* **2007**, 107, 3180.

(5) For leading references on arrested-Nazarov, see: (a) Scadeng, O.; Ferguson, M. J.; West, F. G. *Org. Lett.* **2011**, 13, 114. (b) Rostami, A.; Wang, Y.; Arif, A. M.; McDonald, R.; West, F. G. *Org. Lett.* **2007**, 9, 703. (c) Song, D.; Rostami, A.; West, F. G. *J. Am. Chem. Soc.* **2007**, 129, 12019. (d) White, T. D.; West, F. G. *Tetrahedron Lett.* **2005**, 46, 5629. (e) Giese, S.; Kastrup, L.; Stiens, D.; West, F. G. *Angew. Chem., Int. Ed.* **2000**, 39, 1970. (f) Bender, J. A.; Arif, A. M.; West, F. G. *J. Am. Chem. Soc.* **1999**, 121, 7443.

(6) Also see: (a) Dhoro, F.; Tius, M. A. *J. Am. Chem. Soc.* **2005**, 127, 12472. (b) Dhoro, F.; Kristensen, T. E.; Stockmann, V.; Yap, G. P. A.; Tius, M. A. *J. Am. Chem. Soc.* **2007**, 129, 7256. (c) Nair, V.; Bindu, S.; Sreekumar, V.; Chiaroni, A. *Org. Lett.* **2002**, 4, 2821.

(7) (a) Tius, M. A.; Chu, C. C.; Nieves-Colberg, R. *Tetrahedron Lett.* **2001**, 42, 2419. (b) Bow, W. F.; Basak, A. K.; Jolit, A.; Vicić, D. A.; Tius, M. A. *Org. Lett.* **2010**, 12, 440. (c) Shimada, N.; Ashburn, B. O.; Basak, A. K.; Bow, W. F.; Vicić, D. A.; Tius, M. A. *Chem. Commun.* **2010**, 46, 3774. For an example postulating an imino-Nazarov reaction pathway, see: (d) Suarez-Pantiga, S.; Rubio, E.; Alvarez-Rua, C.; González, J. M. *Org. Lett.* **2009**, 11, 13.

(8) Kim, S. H.; Cha, J. K. *Synthesis* **2000**, 2113.

(9) (a) Larini, P.; Guarna, A.; Occhiato, E. G. *Org. Lett.* **2006**, 8, 781. (b) Occhiato, E. G.; Prandi, C.; Ferrali, A.; Guarna, A. *J. Org. Chem.* **2005**, 70, 4542.


(10) (a) Bitar, A. Y.; Frontier, A. J. *Org. Lett.* **2009**, 11, 49. (b) Malona, J. A.; Colbourne, J. M.; Frontier, A. J. *Org. Lett.* **2006**, 8, 5661.

(11) Kerr, D. J.; Miletic, M.; Chaplin, J. H.; White, J. M.; Flynn, B. L. *Org. Lett.* **2012**, 14, 1732.

(12) Wu, Y.-K.; Niuz, T.; West, F. G. *Chem. Commun.* **2012**, 48, 9186.

(13) Smith, D. A.; Ulmer, C. W., II. *J. Org. Chem.* **1997**, 62, 5110.

(14) For a leading review on allenamide chemistry, see: Hsung, R. P.; Wei, L.-L.; Xiong, H. *Acc. Chem. Res.* **2003**, 36, 773.

Table 1. Screening for a Suitable Metal Catalyst


entry ^a	catalyst	time (h)	conversion (%) ^b	13a (%) ^c	14 + 15 ^d (%) ^c
1	AuCl ₃	30	>95	—	55
2	AuCl ₃ /AgSbF ₆	1	>95	—	45
3	AuCl ₃ /AgSbF ₆ ^e	22	48	—	36
4	AuCl ₃ /AgSbF ₆ ^f	40	55 ^g	—	36
5	AuCl/AgSbF ₆	5	>95	—	48
6	AuCl(PPh ₃)/AgSbF ₆	60	75 ^g	50 ^h	24
7	JohnphosAuCl/AgSbF ₆	5	>95	91	—
8	IPrAuCl/AgSbF₆	1	>95	97	—
9	IPrAuCl	21	<5	—	—
10	AgSbF ₆ ⁱ	20	64	—	24
11	PtCl ₂	24	<5	—	—
12	PtCl ₄ ⁱ	12	>95	—	82 ^j

^a Unless noted, all reactions were carried out at 0.1 mmol scale in 2 mL of CH₂Cl₂ (concn = 0.05 M) at rt with the addition of 5 mol % catalyst. ^b Conversion determined by ¹H NMR. ^c Isolated yields. ^d **14** and **15** could not be separated by column chromatography. ^e 20 mL of CH₂Cl₂ were used (concn = 0.005 M). ^f 35 mg of 4 Å MS were added. ^g Based on recovered **12a**. ^h Inseparable mixture with **12a**. ⁱ 10 mol % of catalyst was used. ^j An inseparable mixture of **14**, **15**, and unidentified products.

that led to amido ketones **11** through a facile trapping of the initial enone.

Inspired by recent gold catalysis²³ particularly in the area of a Nazarov-type process²⁴ with the report by Zhang,^{19d} we turned our attention to gold catalysts. As shown in Table 1, a trivalent gold catalyst was first investigated. Reactions of **12a** proceeded smoothly in the presence of a catalytic amount of AuCl₃ or cationic AuCl₃/AgSbF₆ (entries 1 and 2) but gave an inseparable mixture of dimers **14** and **15**²⁵ with no desired Nazarov cyclization product **13a**. Attempts to suppress the dimerization by lowering the concentration (entry 3) or addition of 4 Å MS (entry 4) proved to be fruitless.

Nevertheless, we were encouraged by dimers **14** and **15** because they imply that the desired imino-Nazarov


(24) For examples of Nazarov-like processes through activations of vinyl allenes, see: (a) Chen, B.; Fan, W.; Chai, G.; Ma, S. *Org. Lett.* **2012**, *14*, 3616. (b) Lemi re, G.; Gandon, V.; Cariou, K.; Hour, A. *J. Am. Chem. Soc.* **2009**, *131*, 2993. (c) Oh, C. H.; Karmakar, S. *J. Org. Chem.* **2009**, *74*, 370. (d) Bhunia, S.; Liu, R.-S. *J. Am. Chem. Soc.* **2008**, *130*, 16488. (e) Shi, F.-Q.; Li, X.; Xia, Y.; Zhang, L.; Yu, Z.-X. *J. Am. Chem. Soc.* **2007**, *129*, 15503. (f) Lee, J. H.; Toste, F. D. *Angew. Chem., Int. Ed.* **2007**, *46*, 912. (g) Funami, H.; Kusama, H.; Iwasawa, N. *Angew. Chem., Int. Ed.* **2007**, *46*, 909. (h) Marion, N.; Diez-Gonz lez, S.; Fr mont, P.; de; Noble, A. R.; Nolan, S. P. *Angew. Chem., Int. Ed.* **2006**, *45*, 3647. (i) Shi, X.; Gorin, D. J.; Toste, F. D. *J. Am. Chem. Soc.* **2005**, *127*, 5802.

(25) See Supporting Information.

Table 2. A Au(I)-Catalyzed Imino-Nazarov Cyclization


entry ^a	allenamide	product	time (h)	yield (%) ^b
1	12b : R = Bu	13b : R = Bu	36	94
2	12c : R = Bn	13c : R = Bn	30	90
3	12d	13d	12 ^c	81
4	12e : R = 3-OMe	13e : R = 3-OMe	1	97
5	12f : R = 2-OMe	13f : R = 2-OMe	29	16
6	12g : R = 4-Me	13g : R = 4-Me	1	92
7	12h : R = 2-Me	13h : R = 2-Me	24	90
8	12i : R = H	13i : R = H	1	93
9	12j : R = 4-Br	13j : R = 4-Br	12 ^c	— ^d
10	12k	13k	1	91

^a Unless noted, all reactions were carried out at 0.1 mmol scale in 2 mL of CH₂Cl₂ (concn = 0.05 M) at rt with the addition of 5 mol % catalyst. ^b Isolated yields. ^c The reaction was run at 50 °C in DCE. ^d No reaction.




reaction had taken place possibly postdimerization through the intermediacy of **14'** [addition of **12a** to activated-**12a**] Gratifyingly with AuCl(PPh₃)/AgSbF₆, **13a** was obtained for the first time in 50% yield along with the **14/15** mixture in 24% yield. Further optimizations revealed that the best yield was reached when using IPrAuCl/AgSbF₆, leading to **13a** in 97% yield (entry 8) with JohnphosAuCl/AgSbF₆ affording **13a** in 91% yield. Remarkably, neutral IPrAuCl itself turned out to be completely inactive (entry 9), while AgSbF₆ alone gave **14** and **15**, albeit sluggishly (entry 10). Lastly, Pt catalysts were also examined and gave very poor results in this process (entries 11 and 12).

Having established the optimum catalytic system, the scope of the imino-Nazarov process was explored [Table 2]. Notably, for allenamides equipped with more bulky substituents on the nitrogen (*n*-Bu in **12b** and Bn in **12c**), reactions were slower, although yields were still excellent (entries 1 and 2). The structure of **13b** was unambiguously assigned through its X-ray structure [left in Figure 1]. The cyclization of **12d**, bearing a 4-methoxybenzenesulfonyl [MBS] substituent, also worked but at elevated temperature. Lastly, allenamides with different substituents on the aryl

**Figure 1.** X-ray structures of **13b** and **20**.


Scheme 3. Regioselective Imino-Nazarov Cyclizations




ring (entries 4–9) also proved to be quite effective overall with exceptions of 2-MeO and 4-Br substituents (entries 5 and 9). In particular, while the reaction of **12f** is very tardy, the cyclization of **12j** was completely attenuated. We are not clear of the precise rationale at this point. However, it is noteworthy that the high level of regioselectivity found in cases of **12e** and **12k** was remarkable. While these two cyclizations appear to favor the less hindered C6-position with no detectable cyclization at the C2-position, this preference is likely due to direct donation from the C3–OR group.

To further investigate the regioselectivity issue, allenamides **16** and **19** were subjected to the cyclization conditions [Scheme 3]. For allenamide **16** with an α -(2-naphthyl) substituent, cyclization took place selectively at C1 [see **17'**], affording **17** exclusively in 97% yield. This regioselectivity is also likely an electronic preference given that cyclization at C1 would involve the cyclohexadiene portion of the naphthyl ring, thereby leading to greater cation stabilization: Benzylic stabilization in **17'** vs pseudo-benzylic stabilization in **18'** through cyclization from C3. On the other hand, for allenamide **19** with an α -(1-naphthyl) group, a competing 6 π -electron cyclization of the reaction intermediate [see **21'**] could deliver 1*H*-phenalene **21**. However, the reaction exclusively proceeded through the imino-Nazarov cyclization pathway via **20'**, leading to **20** in excellent yield. The structure of **20** was also confirmed by its single crystal X-ray structure [Figure 1]. It is noteworthy that **17** and **20** are structurally almost identical with the difference residing in the position for the respective enamide motif.

Scheme 4. 4 π - vs 6 π -Electron Electrocyclization



Scheme 5. Synthesis of Indole-Fused Cyclopentenamides



Intrigued by this competition, another reaction with potentially competing imino-Nazarov 4 π -electron and 6 π -electron electrocyclization was examined using allenamide **22** [Scheme 4]. The reaction of **22** led to **25** and **26** as the only discernible products with the latter likely coming from hydrolysis of the initial Nazarov cyclization product **25**. This study suggests that while the cationic imino-Nazarov intermediate could exist in two equilibrating conformations **23** and **24** differing only at the 's'-marked C–C bond [or being *trans*- and *cis*-1-aza-dienes, respectively, shown in green], 6 π -electron electrocyclization of 3-aza-triene **24**^{26,27} was also not competitive in this case.

Lastly, while cyclopentenamides are synthetically useful, we recognized that our imino-Nazarov cyclization could provide a concise approach to indole-fused cyclopentenamides [or cyclopenta[*b*]indoles],^{24a} thereby representing an opportunity to develop a new strategy for natural product synthesis.²⁸ Consequently, as shown in Scheme 5, under the optimized conditions, cyclizations of α -indolyl-substituted allenamides **28a** and **28b** proceeded smoothly to afford indole-fused cyclopentenamides **29a** and **29b**, respectively, in good yields.

We have showcased here an imino-Nazarov cyclization using α -aryl-substituted allenamides. This Au(I)-catalyzed cascade represents a successful solution to overcome the challenge in designing an imino-Nazarov cyclization cascade. It is a highly efficient and regioselective transformation in constructing a diverse array of aromatic-ring fused cyclopentenamides. Further mechanistic studies and applications of this cyclization process in total synthesis are currently underway.

Acknowledgment. We thank the NIH [GM066055] for funding. We thank Dr. Vic Young of the University of Minnesota for providing X-ray structural analysis.

Supporting Information Available. Experimental procedures, NMR spectra, X-ray structural files, and characterizations for all new compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.

(26) Okamura, W. H.; de Lera, A. R. In *Comprehensive Organic Synthesis*; Trost, B. M., Fleming, I., Paquette, L. A., Eds.; Pergamon Press: New York, 1991; Vol. 5, pp 699–750.

(27) For leading references, see: (a) Tanaka, K.; Katsumura, S. *J. Am. Chem. Soc.* **2002**, *124*, 9660. (b) Sklenicka, H. M.; Hsung, R. P.; McLaughlin, M. J.; Wei, L.-L.; Gerasuto, A. I.; Brennessel, W. W. *J. Am. Chem. Soc.* **2002**, *124*, 10435.

(28) For some examples, see: (a) Feng, T.; Li, Y.; Wang, Y.-Y.; Cai, X.-H.; Liu, Y.-P.; Luo, X.-D. *J. Nat. Prod.* **2010**, *73*, 1075. (b) He, Y.-L.; Chen, W.-M.; Feng, X.-Z. *Phytochemistry* **1994**, *37*, 1055.

The authors declare no competing financial interest.