The Facile Reduction of Carbon Dioxide to Carbon Monoxide with an Amido-Digermyne**

Jiaye Li, Markus Hermann, Gernot Frenking,* and Cameron Jones*

The steady increase in the atmospheric concentration of the greenhouse gas, CO_2 , since the industrial revolution is thought to be the main cause of recent increases in global temperatures.^[1] The future implications of this phenomenon are driving significant current efforts to develop efficient methods for the sequestration of CO_2 ,^[2] and for its use as a C_1 feedstock in the formation of useful chemicals.^[3] With regard to the latter, the reduction of CO₂ to CO is of particular interest as carbon monoxide can be used as a fuel or as a chemical feedstock in its own right. However, CO₂ reduction is problematic because of the considerable strength of its O=C(O) bond (532 kJ mol⁻¹),^[4] and for kinetic reasons.^[3] With that said, CO₂ reduction has been achieved, either stoichiometrically or catalytically, in nature (e.g. with CO dehydrogenase/acetyl-coenzyme A synthase)^[5] and in the laboratory by use of photolytic,^[6] electrochemical,^[7] or metalbased oxygen abstraction protocols.[8]

While the vast majority of methodologies for the reduction of CO₂ to CO require d- or f-block metal-containing materials to proceed, a handful of low oxidation state p-block compounds have recently been shown to effect CO₂ reductions at room temperature. These include three-coordinate, intramolecularly donor-stabilized $(D\rightarrow)$ silylenes $(R_2(D\rightarrow)Si:),^{[9,10]}$ a disilyne $(R(D\rightarrow)SiSi(\leftarrow D)R),^{[10]}$ and *N*-heterocyclic carbenes.^[11] Moreover, a small number of normal oxidation state p-block systems, for example, Al/P-based "frustrated Lewis pairs", have been reported to reduce CO₂ to CO.^[12] The reduction reactions involving these compounds exemplify the rapidly emerging interest in the "transitionmetal-like" reactivity of main group compounds.^[13] We have become involved in this area with, for example, the preparation of the bulky amido-substituted two-coordinate digermyne, [LGe-GeL] 1 $(L = N(Ar^*)(SiMe_3)),$ Ar*= C₆H₂{C(H)Ph₂}₂Me-2,6,4).^[14] Unlike all previously reported bulky aryl-substituted digermynes which have Ge-Ge multiple bonds,^[15,16] compound 1 has an extremely long (2.7093(7) Å) Ge-Ge single bond, which is thought to be the basis of its very narrow HOMO-LUMO energy gap (0.62 eV, calculated using RI-BP86/def2-SVP; HOMO/ LUMO = highest occupied/lowest unoccupied molecular orbital). The resultant high reactivity of 1 has been demonstrated by the fact that it quantitatively and rapidly activates dihydrogen, to give [LGeGe(H)₂L], at temperatures as low as -10 °C, both in solution and the solid state. It seemed reasonable to us that 1 could participate in other small molecule activations that have not been previously achieved with germanium compounds, and which are normally thought to be the realm of transition-metal systems. In this respect, here, we show that the digermyne facilely and quantitatively reduces CO_2 to CO at temperatures as low as -40 °C. The mechanism of this reduction has been explored using spectroscopic and computational techniques, and the reductions of the CO₂ analogs, CS₂ and tBuNCO, are also described.

A toluene solution of **1** was exposed to an excess of dry CO_2 at -70 °C, and the reaction vessel was sealed. Upon warming, the solution slowly lost the deep purple color of the digermyne at about -60 °C and became orange-brown. From about -40 to -30 °C the color of the solution changed to pale yellow, and remained that color when warmed to room temperature. At that point, an IR spectroscopic analysis of the head space gas of the reaction vessel confirmed the presence of CO ($\tilde{v} = 2143$ cm⁻¹). Subsequent removal of volatiles from the mixture afforded an essentially quantitative yield of the novel bis(germylene) oxide **2** (Scheme 1). To ascertain the yield of generated CO, the method of Baceiredo

[*] J. Li, Prof. C. Jones
School of Chemistry, Monash University
PO Box 23, Melbourne, VIC, 3800 (Australia)
E-mail: cameron.jones@monash.edu
M. Hermann, Prof. Dr. G. Frenking
Fachbereich Chemie, Philipps-Universität Marburg
35032 Marburg (Germany)
E-mail: frenking@chemie.uni-marburg.de

[**] We thank the Australian Research Council (C.J.), The U.S. Air Force Asian Office of Aerospace Research and Development (grant FA2386-11-1-4110 to C.J.), and the Deutsche Forschungsgemeinschaft. The EPSRC is also thanked for access to the UK National Mass Spectrometry Facility.

Supporting information for this article, including full synthetic and spectroscopic details for compounds 2–5, crystallographic data for 2–4, and full details and references for the DFT calculations, is available on the WWW under http://dx.doi.org/10.1002/anie. 201203607.



Angew. Chem. Int. Ed. 2012, 51, 1-5

© 2012 Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim

These are not the final page numbers!

🕅 WILEY 盾

and co-workers was employed.^[10] That is, the reaction was repeated in a vessel that was connected through a gas bridge to another vessel containing half an equivalent of [RhCl-(COD)(IPrMe)] (COD = cycloocta-1,5-diene; IPrMe = :C{N(*i*Pr)C(Me)}₂) in a [D₆]benzene solution. The sealed two-component vessel was allowed to stand for 48 h at 20 °C, after which time an ¹H NMR spectroscopic analysis of the [D₆]benzene solution revealed the quantitative conversion of [RhCl(COD)(IPrMe)] to [RhCl(CO)₂(IPrMe)] and free COD (see the Supporting Information for further details). Accordingly, the reduction of CO₂ to CO by **1** was confirmed to be essentially quantitative.^[17]

A series of related reactions were subsequently investigated for the purpose of comparison (Scheme 1). First, compound 2 was found to be alternatively accessible in high yield by treatment of 1 with excess N₂O. Interestingly, the germanium(II) centers of 2 are not further oxidized by N₂O. This outcome contrasts with the reaction of Power's multiply bonded digermyne, Ar'GeGeAr' $(Ar' = C_6H_3(C_6H_3iPr_2-2,6)_2-$ 2,6), with N_2O , which yields a cyclic germanium(IV) peroxospecies.^[18] The reaction of $\mathbf{1}$ with an excess of CS₂ proceeded rapidly at -70 °C and afforded a high yield of the pale yellow bis(germylene) sulfide 3 after subsequent warming to ambient temperature. Treatment of 1 with the isocyanate, tBuNCO, gave a mixture of products which included significant quantities of 2, but no observable *t*BuNC. It was thought that if tBuNC was generated in this reaction it could compete with tBuNCO for reaction with 1. This proposal was assessed by reacting 1 directly with tBuNC, which gave a good yield of the green reductively coupled product 4. Compound 4 was subsequently found to be a component of the tBuNCO/ digermyne reaction mixture (as determined by NMR spectroscopy). Therefore, it seems likely that tBuNC is generated in that reaction, but is rapidly consumed by reduction with 1. While reductive coupling reactions of isonitriles with low oxidation state main group complexes are not uncommon,^[19] the reaction of tBuNC with Power's digermyne, Ar'GeGeAr', yielded only the 1:1 adduct, [Ar'GeGe(Ar')(CNtBu)].^[18] This could indicate that 1 is both more electrophilic and more reducing towards isonitriles than Ar'GeGeAr'.

The spectroscopic data for 2-4 are fully consistent with their solid-state structures (see the Supporting Information for details). Compounds 2 and 3 represent the first crystallographically characterized examples of two-coordinate, amidosubstituted bis(germylene) chalcogenides (Figure 1), though two related three-coordinate examples, $[{(amid')Ge}_2E] (E =$ O or S, amid' = { $(C_6H_3iPr_2-2,6)N_2CtBu$ }, and a two-coordinate aryl-substituted complex, [Ar'GeOGeAr'], have been reported.^[20] The two compounds in the current study are broadly isostructural, though it is noteworthy that they exhibit markedly different Ge-E-Ge angles $(E = O: 122.30(9)^\circ; S:$ 98.21(3)°), in line with the lesser propensity of sulfur, relative to oxygen, to undergo hybridization. The Ge-E distances in both compounds are in the normal range for single-bond interactions,^[21] while their Ge-N distances are close to those in 1 (1.872(2) Å). It is interesting that the germanium lone pairs in 2 and 3 adopt a *cis* disposition relative to each other, and therefore the compounds have the potential to act as chelating Ge-donor ligands (compare the three-coordinate



Figure 1. Molecular structures of compounds a) **2** and b) **3** (25% ellipsoids; hydrogen atoms are omitted). Relevant bond lengths (Å) and angles (°). **2**: Ge(1)-O(1) 1.8088(16), Ge(1)-N(1) 1.8784(19), Ge(2)-O(1) 1.8154(15), Ge(2)-N(2) 1.8749(19), O(1)-Ge(1)-N(1) 100.95(8), O(1)-Ge(2)-N(2) 97.50(8), Ge(1)-O(1)-Ge(2) 122.30(9). **3**: Ge(1)-N(1) 1.868(2), Ge(1)-S(1) 2.2854(8), Ge(2)-N(2) 1.877(2), Ge(2)-S(1) 2.2869(8), N(1)-Ge(1)-S(1) 99.41(7), N(2)-Ge(2)-S(1) 99.73(7), Ge(1)-S(1)-Ge(2) 98.21(3).

bis(silylene) oxide complex, $[(COD)Ni{Si(amid)}_2O]$, amid = $(tBuN)_2CPh$.^[22] The molecular structure of **4** (Figure 2) reveals the compound to be a dimeric digermabicycle with pyramidal germanium(II) centers, the lone pairs of which are



Figure 2. Molecular structure of compound 4 (25% ellipsoids; hydrogen atoms are omitted). Relevant bond lengths (Å) and angles (°). Ge(1)-N(1) 1.887(3), Ge(1)-C(73) 2.087(11), Ge(1)-N(4) 2.371(7), Ge(2)-N(2) 1.885(3), Ge(2)-C(74) 2.071(11), Ge(2)-N(3) 2.358(6), N(3)-C(73) 1.279(12), N(4)-C(74) 1.286(12), C(73)-C(74) 1.476(17), C(73)-Ge(1)-N(4) 60.1(4), N(1)-Ge(1)-N(4) 107.42(15), N(1)-Ge(1)-C(73) 105.3(3), C(74)-Ge(2)-N(3) 60.0(3), N(2)-Ge(2)-N(3) 108.34(15), N(2)-Ge(2)-C(74) 109.3(2), N(3)-C(73)-C(74) 107.6(12), N(4)-C(74)-C(73) 108.6(11).

www.angewandte.org

© 2012 Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim

These are not the final page numbers!

in *cis* positions relative to each other, as in **2** and **3**. The geometry of the central coupled isonitrile unit is indicative of an essentially localized 1,4-diazabutadiene-2,3-diyl fragment, that is $(N = C_{-})_2$, the N centers of which have markedly longer dative interactions with the Ge atoms than the exocyclic amide nitrogen atoms.

To shed light on the mechanism of the reduction of CO_2 by 1, and the temperature-dependent color changes of the reaction mixture, the reaction was followed by ¹H and ¹³C NMR spectroscopy ($[D_8]$ toluene solution) from -70 to 30°C (see the Supporting Information for NMR spectra). No reaction occurred at -70°C, but upon warming to -60°C signals for a new, unsymmetrical compound with chemically inequivalent amide ligands began to appear. After warming to -50 °C the conversion of 1 to this compound was largely complete. Although no signal was evident for free CO in the ¹³C NMR spectrum of the mixture at this temperature, a lowfield resonance ($\delta = 222.8 \text{ ppm}$) had appeared. The unsymmetrical compound was stable until -40°C, whereupon it began converting to symmetrical 2. This conversion became more rapid at -30 °C, and was complete by -20 °C. There was essentially no further change in the NMR spectra up to 30 °C. At this temperature, the signal at $\delta = 222.8$ ppm in the ¹³C NMR spectrum had vanished, and one for free CO ($\delta =$ 184.5 ppm) had appeared. The results of this NMR experiment strongly suggest that the reduction of CO₂ proceeds through an unsymmetrical, metastable intermediate. Unfortunately, all attempts to isolate crystalline samples of this intermediate at low temperatures proved fruitless.

So as to elucidate the nature of this intermediate, the freeenergy profile of the reaction of $\mathbf{1}$ with CO_2 was calculated using density functional theory with inclusion of dispersion interactions (BP86-D3/def2-TZVPP//BP86/SVP), and is depicted in Figure 3. We located two transition states, TS1-2a and TS1-2b, for the initial stage of the reaction, which involve a side-on approach of the CO₂ molecule to one or both Ge centers. While these transition states lead to the same intermediate, IM2, through the insertion of CO₂ into the Ge-Ge bond, the energetically lowest lying entrance channel (activation barrier of $\Delta G^{+} = 17.0 \text{ kcal mol}^{-1}$) is through **TS1**-2a. Intermediate, IM2, then rearranges through a low-energy transition state (**TS2-3**, $\Delta G^{\pm} = 4.1 \text{ kcalmol}^{-1}$) to give the trans-germacarboxylato germanium(II) amide complex, IM3, which is $5.0 \text{ kcal mol}^{-1}$ lower in energy than IM2. The elimination of CO from this to give 2, then proceeds with an energy barrier ($\Delta G^{\pm} = 16.3 \text{ kcal mol}^{-1}$) similar to that for the entrance channel. The overall CO₂ reduction reaction is exergonic by only -17.7 kcal mol⁻¹. Because of the relatively low energy of intermediate IM3, and the calculated barrier to its elimination of CO, we conclude that the unsymmetrical, spectroscopically observed reaction intermediate resembles that compound (i.e. 5 in Scheme 1). Indeed, insertions of CO_2 into metal-metal bonds to give metalla-carboxylate complexes (compare 5) are not uncommon,^[3] and the low-field ¹³C NMR resonance for **5** is in the expected region for such complexes. This conclusion is supported by calculation of the ¹³C NMR chemical shift for a model compound of **5**, that is IM4, in which the amido groups (NMe₂) have methyl substituents. The calculated value at GIAO/MP2/TZVPP of $\delta = 216.3$ ppm is in good agreement with the experimental value of $\delta = 222.8$ ppm.

In summary, the quantitative reduction of CO_2 to CO by a digermyne at temperatures as low as -40 °C has been achieved. Moreover, strong evidence for the mechanism of the reaction has been gained from a combination of spectroscopic and computational studies. There is little precedent for



Figure 3. Free-energy profile of the reaction of 1 with CO₂ (BP86-D3/def2-TZVPP//BP86/def2-SVP). Selected interatomic distances (Å), bond angles (°), and NGeGeN torsion angles, θ (°), are given with the molecular structures.

Angew. Chem. Int. Ed. 2012, 51, 1-5

© 2012 Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim

These are not the final page numbers!

www.angewandte.org

the generation of CO from CO_2 using p-block compounds, and, as far as we are aware, none for germanium systems.^[23] We are currently exploring the possibility of rendering the reduction reaction described above, catalytic; in addition to examining the activation of other small molecules using **1** and related compounds.

Received: May 9, 2012 Revised: June 21, 2012 Published online:

Keywords: carbon dioxide \cdot density functional calculations \cdot germanium \cdot green chemistry \cdot reduction

- [1] *IPCC, Climate Change 2007: Synthesis Report*, Intergovernmental Panel on Climate Change, Geneva, **2007**.
- [2] K. Sumida, D. L. Rogow, J. A. Mason, T. M. McDonald, E. D. Bloch, Z. R. Herm, T.-H. Bae, J. R. Long, *Chem. Rev.* 2012, *112*, 724–781, and references cited therein.
- [3] a) Carbon Dioxide as a Chemical Feedstock (Ed. M. Aresta), Wiley-VCH, Weinheim, 2010; b) M. Cokoja, C. Bruckmeier, B. Rieger, W. A. Herrmann, F. E. Kühn, Angew. Chem. 2011, 123, 8662-8690; Angew. Chem. Int. Ed. 2011, 50, 8510-8537; c) H. Arakawa, M. Aresta, J. N. Armor, M. A. Barteau, E. J. Beckman, A. T. Bell, J. E. Bercaw, C. Creutz, E. Dinjus, D. A. Dixon, K. Domen, D. L. DuBois, J. Eckert, E. Fujita, D. H. Gibson, W. A. Goddard, D. W. Goodman, J. Keller, G. J. Kubas, H. H. Kung, J. E. Lyons, L. E. Manzer, T. J. Marks, K. Morokuma, K. M. Nicholas, R. Periana, L. Que, J. Rostrup-Nielson, W. M. H. Sachtler, L. D. Schmidt, A. Sen, G. A. Somorjai, P. C. Stair, B. R. Stults, W. Tumas, Chem. Rev. 2001, 101, 953-996; d) X. Lin, J. R. Moss, Coord. Chem. Rev. 1999, 181, 27-59; e) W. Leitner, Coord. Chem. Rev. 1996, 153, 257-284.
- [4] CRC Handbook of Chemistry and Physics, 73rd ed. (Ed.: D. R. Lide), CRC, Boca Raton, 1992.
- [5] E. L. Hegg, Acc. Chem. Res. 2004, 37, 775-783.
- [6] W. Lin, H. Frei, J. Am. Chem. Soc. 2005, 127, 1610-1611.
- [7] E. Simon-Manso, C. P. Kubiak, Organometallics 2005, 24, 96– 102.
- [8] See for example: a) J. S. Silvia, C. C. Cummins, J. Am. Chem. Soc. 2010, 132, 2169–2171; b) D. S. Laitar, P. Müller, J. P. Sadighi, J. Am. Chem. Soc. 2005, 127, 17196–17197; c) I. Castro-Rodriguez, K. Meyer, J. Am. Chem. Soc. 2005, 127, 11242–11243.
- [9] S. Yao, Y. Xiong, M. Brym, M. Driess, J. Am. Chem. Soc. 2007, 129, 7268–7269.
- [10] D. Gau, R. Rodriguez, T. Kato, N. Saffon-Merceron, A. de Cozar, F. P. Cossio, A. Baceiredo, *Angew. Chem.* 2011, 123, 1124–1128; *Angew. Chem. Int. Ed.* 2011, 50, 1092–1096.

- [11] L. Gu, Y. Zhang, J. Am. Chem. Soc. 2010, 132, 914–915. N. B. The validity of the reported generation of CO from the reaction of CO₂ with an NHC was subsequently questioned, see P.-C. Chiang, J. W. Bode, Org. Lett. 2011, 13, 2422–2425.
- [12] G. Ménard, D. W. Stephan, *Angew. Chem.* 2011, *123*, 8546–8549;
 Angew. Chem. Int. Ed. 2011, *50*, 8396–8399. N. B. The low yield reduction of CO₂ to CO using KSiH₃ has been reported. V. A. Williams, D. M. Ritter, *Inorg. Chem.* 1985, *24*, 3278–3280.
- [13] Selected recent reviews: a) P. P. Power, *Nature* 2010, 463, 171–177; b) P. P. Power, *Acc. Chem. Res.* 2011, 44, 627–637; c) M. Asay, C. Jones, M. Driess, *Chem. Rev.* 2011, 111, 354–396; d) S. Yao, Y. Xiong, M. Driess, *Organometallics* 2011, 30, 1748–1767; e) D. Martin, M. Soleilhavoup, G. Bertrand, *Chem. Sci.* 2011, 2, 389–399; f) S. S. Sen, S. Khan, P. P. Samuel, H. W. Roesky, *Chem. Sci.* 2012, 3, 659–682.
- [14] a) J. Li, C. Schenk, C. Goedecke, G. Frenking, C. Jones, J. Am. Chem. Soc. 2011, 133, 18622–18625; b) J. Li, A. Stasch, C. Schenk, C. Jones, Dalton Trans. 2011, 40, 10448–10456.
- [15] a) P. P. Power, Organometallics 2007, 26, 4362-4372; b) E. Rivard, P. P. Power, Inorg. Chem. 2007, 46, 10047-10064; c) Y. Peng, R. C. Fischer, W. A. Merrill, J. Fischer, L. Pu, B. D. Ellis, J. C. Fettinger, R. H. Herber, P. P. Power, Chem. Sci. 2010, 1, 461-468.
- [16] As **1** possesses a Ge–Ge single bond it could be regarded as a 1,2bis(germylene). We use the term digermyne to describe the compound in line with nomenclature established for both single and multiple bonded heavier analogs of alkynes, that is the ditetrelynes, REER, E = Si, Ge, Sn, or Pb. See Refs. [13] and [14].
- [17] In a separate experiment, compound **1** was found to be unreactive towards CO at ambient temperature.
- [18] C. Cui, M. M. Olmstead, J. C. Fettinger, G. H. Spikes, P. P. Power, J. Am. Chem. Soc. 2005, 127, 17530-17541.
- [19] See for example: a) X. Li, X. Cheng, H. Song, C. Cui, Organometallics 2007, 26, 1039–1043; b) W. Uhl, U. Schütz, W. Hiller, M. Heckel, Chem. Ber. 1994, 127, 1587–1592; c) W. Uhl, I. Hahn, U. Schütz, S. Pohl, W. Saak, J. Martens, J. Manikowski, Chem. Ber. 1996, 129, 897–901.
- [20] a) S.-H. Zhang, C.-W. So, *Organometallics* 2011, *30*, 2059–2062;
 b) O. T. Summerscales, M. M. Olmstead, P. P. Power, *Organometallics* 2011, *30*, 3468–3471.
- [21] As determined from a survey of the Cambridge Crystallographic Database, May, **2012**.
- [22] W. Wang, S. Inoue, S. Yao, M. Driess, J. Am. Chem. Soc. 2010, 132, 15890–15892.
- [23] The reduction of CO₂ by germanium(II) hydride complexes, to give hydrogermylation products, has been previously reported;
 a) S. L. Choong, W. D. Woodul, C. Schenk, A. Stasch, A. F. Richards, C. Jones, *Organometallics* 2011, *30*, 5543–5550;
 b) A. Jana, D. Ghoshal, H. W. Roesky, I. Objartel, G. Schwab, D. Stalke, *J. Am. Chem. Soc.* 2009, *131*, 1288–1293.

www.angewandte.org

These are not the final page numbers!

Communications



Taking the fizz out: A digermyne compound with a Ge–Ge single bond has been shown to quantitatively reduce CO_2 to CO at temperatures as low as -40 °C. The mechanism of this unprecedented reaction has been probed by spectroscopic and computational techniques and involves a metastable intermediate (see picture, $Ar^* = C_6H_2\{C(H)Ph_2\}_2Me-$ 2,6,4).

© 2012 Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim

einheim www.angewandte.org 5 These are not the final page numbers!