

Journal of The Chemical Society, Chemical Communications

NUMBER 17/1977

7 SEPTEMBER

Formation of Lactones from Dienes *via* Iron Carbonyl Complexes

By GARY D. ANNIS and STEVEN V. LEY*


(Department of Chemistry, Imperial College, London SW7 2AY)

Summary A number of dienes were monoepoxidised and treated with $\text{Fe}(\text{CO})_5$ to form intermediate complexes which were subsequently oxidised to lactones.

RECENT studies¹ on the preparation of δ -lactones from dienes prompts us to report our results in a related area. We were attracted by the known propensity for carbonyl group insertion into vinyloxirans by pentacarbonyliron² and reasoned that the complexes formed would lead to lactones on oxidation. Irradiation of the monoepoxide (1),³ from 2,3-dimethylbutadiene, in the presence of $\text{Fe}(\text{CO})_5$ gave the iron complex (2), which showed structurally significant i.r. bands at 2050, 1990, and 1660 cm^{-1} . The n.m.r. spectrum was also in agreement with the proposed structure (2). Subsequent oxidation with ceric ammonium nitrate at 60 °C in ethanol-water (50:50) gave the lactone (3) in 38% yield.

The known iron complex (5)² was similarly prepared from 1,3-cyclohexadiene monoepoxide (4); however on oxidation with ceric ammonium nitrate at room temperature (4) afforded the β -lactone (6) in 65–70% yield, and the lactone (7)⁴ was not detected. The β -lactone (6) was readily characterized from its spectral data {e.g. $\nu(\text{C}=\text{O})$ 1820 cm^{-1} , typical of β -lactones; δ 5.6–6.35 (2H, m, vinylic), 4.75–5.0 [1H, m, $-\text{CH}-\text{O}-\text{C}(\text{O})-$], 4.12 [1H, dd J 6 Hz, $-\text{C}(\text{O})-\text{CH}=\text{C}-$], and 1.4–2.5 (4H, m, aliphatic)}. Decoupling experiments are in accord with the above assignments and in addition these data are similar to those of recently prepared β -lactones.⁵

The monoepoxide (8) on treatment with $\text{Fe}(\text{CO})_5$ gave the iron complex (9) (79%) which showed typical carbonyl absorptions at ν_{max} 2050, 2000, 1990, and 1660 cm^{-1} . Oxidation of this complex afforded exclusively the β -lactone (10) in 40% yield. Structural assignment follows from the spectral data, ν_{max} 1820 cm^{-1} , δ 1.0–2.2 (m, 6H,

Ts = $p\text{-MeC}_6\text{H}_4\text{SO}_2$

aliphatic), 4.78 (d, 1H J 2 Hz, $-\text{CH}-\text{O}-$), 5.1–5.55 (m, 2H, methylene), and 5.65–6.15 (m, 1H, vinylic). Very few examples of similar α -vinyl β -lactones appear to be known in the literature.⁶

As both the β -lactones (**6**) and (**10**) were rather unstable, they were reduced with lithium aluminium hydride to the corresponding diols which in turn were characterised as their ditosylates (**11**) and (**12**).

The above procedure constitutes the formation of lactones from dienes *via* iron carbonyl complexes.

We thank the S.R.C. for a Research Studentship (to G. D. A.) and Dr. R. Bonjouklian for spectra of (**3**) for comparison.

(Received, 24th May 1977; Com. 512.)

¹ R. A. Ruden and R. Bonjouklian, *J. Amer. Chem. Soc.*, 1975, **97**, 6892; R. Aumann and H. Ring, *Angew. Chem. Internat. Edn.*, 1977, **16**, 50.

² Cf. R. Aumann, K. Fröhlich, and H. Ring, *Angew. Chem. Internat. Edn.*, 1974, **13**, 275.

³ Cf. W. K. Anderson and T. Veysoglu, *J. Org. Chem.*, 1973, **38**, 2267.

⁴ J. R. Malpass and N. J. Tweddle, *J.C.S. Perkin I*, 1977, 874.

⁵ W. E. Barnett and L. L. Needham, *J. Org. Chem.*, 1975, **40**, 2843; G. W. Holbert, L. B. Weiss, and B. Ganem, *Tetrahedron Letters*, 1976, 4435.

⁶ H. E. Zaugg, *Org. Reactions*, 1954, **8**, 305; L. L. Muller and J. Hamer, '1,2-Cycloaddition Reactions,' Wiley, New York, 1967.