

CrossMark
click for updates

A metal free domino synthesis of 3-aryloindoles via two sp^3 C–H activation†

Anupal Gogoi, Anju Modi, Srimanta Guin, Saroj Kumar Rout, Debapratim Das and Bhisma K. Patel*

Cite this: *Chem. Commun.*, 2014, 50, 10445Received 10th June 2014,
Accepted 18th July 2014

DOI: 10.1039/c4cc04407j

www.rsc.org/chemcomm

A metal free synthesis of 3-aryloindole involving two sp^3 C–H activation has been achieved starting from *o*-alkynyl-*N,N*-dialkylamines using catalyst TBAI and oxidant TBHP.


In modern organic synthesis, C–H functionalisation strategies have brought about atom and step-economy by streamlining various arduous synthetic processes.¹ However, in this forum, much of the attention has been focused on functionalisations of sp and sp^2 C–H bonds.² In contrast high pK_a 's (>30 – 35) as well as large bond dissociation energies (>104 kcal mol⁻¹) of various sp^3 C–H bonds make their selective activation a formidable challenge. The solutions to these challenges led to the evolution of various sp^3 C–H activation strategies, one of them being the cross dehydrogenative coupling (CDC). In most cases, functionalisations of sp^3 C–H bonds α to heteroatoms are attained by the CDC strategy using transition metal catalysts.³ The heteroatoms present in precursors are invariably *N*- or *O*- that generate iminium or oxonium ion intermediates *in situ*.^{3,4} These reactive intermediates are formed by a single electron transfer from the heteroatoms to the metal centre that forms a radical cation. This is then followed by a loss of a hydrogen radical from the adjacent carbon generating either an iminium or an oxonium ion which is amenable for the construction of C–C or C–heteroatom bonds by an inter- or an intramolecular nucleophilic attack. For precursors like tertiary amines, a variety of transition metal catalysts in combination with organic peroxides are conducive for the formation of reactive intermediates.^{3,4a–c} Noteworthy, an intramolecular nucleophilic attack at these iminium ions can generate various nitrogen containing heterocycles.^{4a,5,6}

Among nitrogenous heterocycles, the indole ring system finds wide applications in materials chemistry and pharmaceuticals.⁷ In particular, 3-aryloindoles have gained more importance as

they provide alternate routes to other functionalised indoles because of easy manipulation of their carbonyl groups.⁸ As an obvious consequence, development of a straightforward method for the synthesis of 3-aryloindoles has been of great significance. This motivation led to our Cu catalysed synthesis of 3-aryloindole from 2-alkynyl-*N,N*-dialkylamine *via* cascade C–C and C–O bond forming pathways following the above-mentioned intra-molecular CDC strategy (path-a, Scheme 1).⁵ Exactly the same strategy has been reported by the Liang group using a Pd–Cu catalytic system instead of a simple Cu catalyst.⁶ Prior to these reports, several other unconventional methods (*via* C–H activation) for their synthesis have been developed using transition metal catalysts.^{8g,9}

Although transition metal catalysed processes are advantageous they are invariably associated with high cost and difficulties in removal of residual metals from the final product which limits their utility in pharmaceutical industries. Furthermore, for a copper catalysed reaction often doubt arises whether it is the real catalyst or a co-catalyst due to the traces of Pd present in commercial grade Cu salts. There are instances where the metal impurities present even in traces in commercial grade salts are the actual catalyst.¹⁰

To overcome some of the disadvantages associated with metal catalysts recently photo-catalysts¹¹ and molecular iodine or organo-iodide¹² such as tetrabutylammonium iodide (TBAI) in combination with an external oxidant are appropriate substitutes.^{11,12} The use of designer ligands and at times metals are unavoidable in any photo-catalytic process, thereby making the overall method expensive when applied to large scale reactions. Thus, we envisaged that instead of using metal catalysts the

Scheme 1 Synthesis of 3-aryloindole *via* sp^3 C–H activation.


Department of Chemistry, Indian Institute of Technology Guwahati, 781 039, Assam, India. E-mail: patel@iitg.ernet.in; Fax: +91-3612690762

† Electronic supplementary information (ESI) available. CCDC 1006843. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c4cc04407j


domino transformation of 2-alkynyl-*N,N*-dialkylamines to 3-aryl indoles could be realised under metal free conditions using TBAI. With this inspiration in mind our initial investigation began by treating *N,N*-dimethyl-2-(2-phenylethynyl)benzenamine (**1a**) (1 equiv.) with TBAI (20 mol%) and 70% aq. TBHP (3 equiv.) at 80 °C. To our delight aroylindole (**1'a**) was isolated in 52% yield (entry 1, Table S1, ESI[†]). To the best of our knowledge this is the first metal free synthesis of 3-arylindole involving two sp³ C-H activation (path-b, Scheme 1).

Encouraged by this metal-free cascade synthesis of 3-aryl indole, other reaction parameters such as catalysts, oxidants and solvents were varied to attain the best possible yield. Increasing the amount of TBHP from 3 equiv. to 4 equiv. and 5 equiv. resulted in an improved yield of 59% and 74% respectively (entries 2 and 3, Table S1, ESI[†]). The use of other halogen analogues such as KI, I₂, and tetrabutylammonium bromide (TBAB), instead of TBAI, was found to be less effective (entries 4–6, Table S1, ESI[†]). A decane solution of TBHP (5–6 M) (5 equiv.) in lieu of its aqueous solution (70%) gave a comparable yield of 69%, however, other peroxides such as di-*tert*-butyl peroxide (DTBP) or aq. H₂O₂ (50%) failed to give satisfactory yield of the product (entries 7–9, Table S1, ESI[†]). Furthermore, the use of other solvents such as DMF (62%), toluene (46%), 1,4-dioxane (54%), DCE (37%) was found to be less productive than DMSO (entries 10–13, Table S1, ESI[†]). The yield dropped to 56% when the catalyst loading was reduced to 15 mol% (entry 14, Table S1, ESI[†]). Neither TBAI nor TBHP alone were capable of triggering this domino transformation, suggesting an essential requirement of their combination (entries 15 and 16, Table S1, ESI[†]). Thus, catalyst TBAI (20 mol%), and aqueous TBHP (5 equiv.) in DMSO at 80 °C were found to be the ideal conditions for this transformation (entry 3, Table S1, ESI[†]).

After achieving the optimised conditions, this methodology was applied to different *o*-alkynyl-*N,N*-dialkylamines to afford their respective 3-arylindoles. As shown in Schemes 2 and 3, a series of aroylindoles could be obtained in moderate to excellent yields from their aminoalkyne precursors. At first, effects of substituents on the aryl ring of the alkynes were examined. The electron-donating substituents *viz.* *m*-Me (**1b**), *p*-tBu (**1c**) and *p*-OMe (**1d**) when present in the aryl ring of the alkyne had a positive impact on the product (**1'b–1'd**) yields (in the range of 76–78%). However, when the aryl ring contains electron-withdrawing groups such as *p*-Br (**1e**) and *m*-F (**1f**), the reactions proceeded sluggishly to afford their corresponding aroylindoles (**1'e**) and (**1'f**) in slightly lesser yields (in the range 67–69%). When the amine bearing aryl rings of the *o*-alkynylamines are substituted with weakly electron-donating groups such as *p*-Me and the other aryl ring being unsubstituted (**2a**) or substituted with an electron-withdrawing group (**2f**), good yields of the products were achieved. An excellent yield of 81% was obtained when both the rings contain electron-donating groups as found for (**2'd**). A moderately electron-withdrawing group such as *p*-Cl present in the aryl ring bearing the tertiary amine group resulted in comparatively lesser yields regardless of the nature of the substituents on the other ring as exemplified for (**3'a**), (**3'g**) and (**3'e**). The scope of this methodology was next extended to




Scheme 2 Substrate scope of 3-arylindoles. Reaction conditions: *o*-alkynyl-*N,N*-dialkylamine (0.25 mmol), TBAI (0.05 mmol), TBHP (1.25 mmol) DMSO (1 mL) at 80 °C. Isolated yields.



Scheme 3 Substrate scope of 3-arylindoles. Reaction conditions: *o*-alkynyl-*N,N*-dialkylamine (0.25 mmol), TBAI (0.05 mmol), TBHP (1.25 mmol) DMSO (1 mL) at 80 °C. Isolated yields.

annular tertiary amines. The *o*-alkynyl amine precursors with a six- or a five-membered ring provided their corresponding aroylindoles under the optimised reaction conditions as demonstrated by the synthesis of (**4'a**), (**4'd**), (**4'e**) (**5'a**) and (**5'g**). However, the yields were moderate which may be due to the ring strain associated with the resulting fused aroylindoles. The structure of the aroylindole (**5'g**) was confirmed by X-ray crystallography (Fig. S1, see ESI[†]).



Scheme 4 Proposed mechanism for the formation of 3-aryloindoles.

Based on literature reports^{12a,n,o} and trends in the yields of products obtained for substituted substrates a plausible mechanism has been proposed for this transformation. An aminyl radical cation (**A**) is formed by a single electron transfer (SET) process from the nitrogen atom of the *o*-alkynyl-*N,N*-dialkylamine. Abstraction of a hydrogen radical α to the nitrogen atom gives an iminium intermediate (**B**). The intermediate (**B**) then undergoes annulation by an intramolecular nucleophilic attack of the alkynyl group at the iminium carbon with simultaneous attack of water or TBHP at the alkenyl carbon to give intermediate species (**C**).^{5,6} The reaction of the substrate (**1a**) under the reaction conditions in the presence of 20 equivalent of H₂¹⁸O afforded aroyloindole (**1'a**) without any ¹⁸O incorporation indicating TBHP as the possible oxygen source. Ketonisation of (**C**) provides 3-aryloindoline (**D**) which is finally oxidised/aromatised to its 3-aryloindole (Scheme 4).

In conclusion, we have developed a metal free method for the synthesis of 3-aryloindoles from *o*-alkynyl-*N,N*-dialkylamine through a TBAI catalysed intramolecular oxidative coupling pathway using TBHP as the oxidant. This protocol simultaneously installs C–C and C–O bonds at the expense of two sp³ C–H bonds. The use of an inexpensive and environmentally benign catalytic system and relatively low reaction time and temperature make the present protocol practically more applicable.

B. K. P acknowledges the support for this research by the Department of Science and Technology (DST) (SB/S1/OC-53/2013), New Delhi, and the Council of Scientific and Industrial Research (CSIR) (02(0096)/12/EMR-II).

Notes and references

- (a) G. Dyker, *Handbook of C–H Transformations: Applications in Organic Synthesis*, Wiley-VCH, Weinheim, 2005; (b) J.-Q. Yu and Z.-J. Shi, *C–H Activation*, Springer, Berlin, Germany, 2010; (c) T. Satoh and M. Miura, *Chem. – Eur. J.*, 2010, **16**, 11212; (d) X. Chen, K. M. Engle, D.-H. Wang and J.-Q. Yu, *Angew. Chem., Int. Ed.*, 2009, **48**, 5094; (e) T. W. Lyons and M. S. Sanford, *Chem. Rev.*, 2010, **110**, 1147; (f) L.-M. Xu, B.-J. Li, Z. Yang and Z.-J. Shi, *Chem. Soc. Rev.*, 2010, **39**, 712; (g) D. A. Colby, R. G. Bergman and J. A. Ellman, *Chem. Rev.*, 2010, **110**, 624; (h) J. Wencel-Delord, T. Droge, F. Liu and F. Glorius, *Chem. Soc. Rev.*, 2011, **40**, 4740; (i) S. H. Cho, J. Y. Kim, J. Kwak and S. Chang, *Chem. Soc. Rev.*, 2011, **40**, 5068; (j) C.-L. Sun, B.-J. Li and Z.-J. Shi, *Chem. Rev.*, 2011, **111**, 1293; (k) L. Ackermann, *Chem. Rev.*, 2011, **111**, 1315; (l) S. R. Neufeldt and M. S. Sanford, *Acc. Chem. Res.*, 2012, **45**, 936; (m) K. M. Engle, T.-S. Mei, M. Wasa and J.-Q. Yu, *Acc. Chem. Res.*, 2012, **45**, 788.
- (a) C. S. Yeung and V. M. Dong, *Chem. Rev.*, 2011, **111**, 1215; (b) C. Liu, H. Zhang, W. Shi and A. Lei, *Chem. Rev.*, 2011, **111**, 1780; (c) B. J. Lie and Z. J. Shi, *Chem. Soc. Rev.*, 2012, **41**, 5588.
- (a) C.-J. Li, *Acc. Chem. Res.*, 2009, **42**, 335; (b) W. Yoo and C.-J. Li, *Top. Curr. Chem.*, 2010, **292**, 281; (c) Z. Li, D. S. Bohle and C.-J. Li, *Proc. Natl. Acad. Sci. U. S. A.*, 2006, **103**, 8928; (d) S. A. Girard, T. Knauber and C.-J. Li, *Angew. Chem., Int. Ed.*, 2014, **53**, 74; (e) F. Jia and Z. Li, *Org. Chem. Front.*, 2014, **1**, 194.
- (a) A. Modak, U. Dutta, R. Kancherla, S. Maity, M. Bhadra, S. M. Mobin and D. Maiti, *Org. Lett.*, 2014, **16**, 2602; (b) G. Zhang, Y.-X. Ma, S.-L. Wang, Y.-H. Zhang and R. Wang, *J. Am. Chem. Soc.*, 2012, **134**, 12334; (c) G. Zhang, Y. Zhang and R. Wang, *Angew. Chem., Int. Ed.*, 2011, **50**, 10429; (d) C. A. Correia and C.-J. Li, *Heterocycles*, 2010, **82**, 555; (e) Y. Zhang and C.-J. Li, *Angew. Chem., Int. Ed.*, 2006, **45**, 1949; (f) Y. Zhang and C.-J. Li, *J. Am. Chem. Soc.*, 2006, **128**, 4242.
- A. Gogoi, S. Guin, S. K. Rout and B. K. Patel, *Org. Lett.*, 2013, **15**, 1802.
- X.-F. Xia, L.-L. Zhang, X.-R. Song, Y.-N. Niu, X.-Y. Liu and Y.-M. Liang, *Chem. Commun.*, 2013, **49**, 1410.
- (a) M. Somei and F. Yamada, *Nat. Prod. Rep.*, 2004, **21**, 278; (b) M. Somei and F. Yamada, *Nat. Prod. Rep.*, 2005, **22**, 73; (c) T. Kawasaki and K. Higuchi, *Nat. Prod. Rep.*, 2005, **22**, 761; (d) M. C. Van Zandit, M. L. Jones, D. E. Gunn, L. S. Geraci, J. H. Jones, D. R. Sawicki, J. Sredy, J. L. Jacol, A. T. DiCiccio, T. Petrova, A. Mitschler and A. D. Podjarny, *J. Med. Chem.*, 2005, **48**, 3141; (e) J. A. Joule, *Product Class 13: Indoles and Its Derivatives*, Thieme, Stuttgart, 2001; (f) M. Bandini and A. Eichholzer, *Angew. Chem., Int. Ed.*, 2009, **48**, 9608.
- (a) G. W. Gribble, In *Comprehensive Heterocyclic Chemistry*, ed. A. R. Katritzky, C. W. Rees and E. F. V. Scriven, Pergamon Press, New York, 1996, vol. 2, p. 207; (b) S. Massa, R. Di Santo and M. Artico, *J. Heterocycl. Chem.*, 1990, **27**, 1131; (c) I. Coldham, B. C. Dobson, S. R. Fletcher and A. I. Franklin, *Eur. J. Org. Chem.*, 2007, 2676; (d) R. Lauchli and K. J. Shea, *Org. Lett.*, 2006, **8**, 5287; (e) S. Tohyama, T. Choshi, K. Matsumoto, A. Yamabuki, K. Ikegata, J. Nobuhiro and S. Hibino, *Tetrahedron Lett.*, 2005, **46**, 5263; (f) J. E. Macor, D. H. Blank, C. B. Fox, A. L. Lebel, M. E. Newman, R. J. Post, K. Ryan, A. W. Schmidt, D. W. Schulz and B. K. Koc, *J. Med. Chem.*, 1994, **37**, 2509; (g) W.-L. Wu and W.-P. Su, *J. Am. Chem. Soc.*, 2011, **133**, 11924.
- (a) Y. Ma, J. You and F. Song, *Chem. – Eur. J.*, 2013, **19**, 1189; (b) R. Bernini, G. Fabrizi, A. Sferrazza and S. Cacchi, *Angew. Chem., Int. Ed.*, 2009, **48**, 8078; (c) L. Yu, P. Lia and L. Wang, *Chem. Commun.*, 2013, **49**, 2368; (d) C. Wang, S. Wang, H. Li, J. Yan, H. Chi, X. Chen and Z. Zhang, *Org. Biomol. Chem.*, 2014, **12**, 1721; (e) T. Jiang and G. Wang, *Org. Lett.*, 2013, **15**, 788; (f) Y. Li, D. Xue, W. Lu, X. Fan, C. Wang and J. Xiao, *RSC Adv.*, 2013, **3**, 11463.
- P.-F. Larsson, A. Correa, M. Carril, P.-O. Norrby and C. Bolm, *Angew. Chem., Int. Ed.*, 2009, **48**, 5691.
- (a) D. M. Schultz and T. P. Yoon, *Science*, 2014, **343**, 985; (b) J. Xie, H. Jin, P. Xu and C. Zhu, *Tetrahedron Lett.*, 2014, **55**, 36; (c) M. Reckenthalera and A. G. Griesbeck, *Adv. Synth. Catal.*, 2013, **355**, 2727; (d) J. Xuan, L.-Q. Lu, J.-R. Chen and W.-J. Xiao, *Eur. J. Org. Chem.*, 2013, 6755; (e) Y.-Q. Zou, J.-R. Chen and W.-J. Xiao, *Angew. Chem., Int. Ed.*, 2013, **52**, 11701; (f) C. K. Prier, D. A. Rankic and D. W. C. MacMillan, *Chem. Rev.*, 2013, **113**, 5322; (g) Y. Xi, H. Yi and A. Lei, *Org. Biomol. Chem.*, 2013, **11**, 2387; (h) D. P. Hari and B. König, *Angew. Chem., Int. Ed.*, 2013, **52**, 4734; (i) D. P. Hari and B. König, *Org. Lett.*, 2011, **13**, 3852; (j) P. Zhang, T. Xiao, S. Xiong, X. Dong and L. Zhou, *Org. Lett.*, 2014, **16**, 3264.
- (a) X.-F. Wu, J.-L. Gong and X. Qi, *Org. Biomol. Chem.*, 2014, **12**, 5807; (b) L. Chen, E. Shi, Z. Liu, S. Chen, W. Wei, H. Lei, K. Xu and X. Wan, *Chem. – Eur. J.*, 2011, **17**, 4085; (c) B. Tan, N. Toda and C. F. Barbas III, *Angew. Chem., Int. Ed.*, 2012, **51**, 12538; (d) J. Feng, S. Liang, S.-Y. Chen, J. Zhang, S.-S. Fu and X.-Q. Yu, *Adv. Synth. Catal.*, 2012, **354**, 1287; (e) J. Huang, L.-T. Li, H.-Y. Li, E. Husan, P. Wang and B. Wang, *Chem. Commun.*, 2012, **48**, 10204; (f) L.-T. Li, J. Huang, H.-Y. Li, L.-J. Wen, P. Wang and B. Wang, *Chem. Commun.*, 2012, **48**, 5187; (g) M. Uyanik, D. Suzuki, T. Yasui and K. Ishihara, *Angew. Chem., Int. Ed.*, 2011, **50**, 5331; (h) S. Chen, Y. Xu and X. Wan, *Org. Lett.*, 2011, **13**, 6152; (i) W. Wei, C. Zhang, Y. Xu and X. Wan, *Chem. Commun.*, 2011, **47**, 10827; (j) T. Froehr, C. P. Sindlinger, U. Kloeckner, P. Finkbeiner and B. J. Nachtsheim, *Org. Lett.*, 2011, **13**, 3754; (k) L. Ma, X. Wang, W. Yu and B. Han, *Chem. Commun.*, 2011, **47**, 11333; (l) M. Uyanik and K. Ishihara, *ChemCatChem*, 2012, **4**, 177; (m) G. Majji, S. Guin, A. Gogoi, S. K. Rout and B. K. Patel, *Chem. Commun.*, 2013, **49**, 3031; (n) J. Dhineshkumar, M. Lamani, K. Alagiri and K. R. Prabhu, *Org. Lett.*, 2013, **15**, 1092; (o) T. Nobuta, N. Tada, A. Fujiya, A. Kariya, T. Miura and A. Itoh, *Org. Lett.*, 2013, **15**, 574.