

182. A Convenient Synthesis of Vinyl Sulfides

by Alan R. Katritzky*, Amir S. Afridi, and Wojciech Kuzmierkiewicz

Department of Chemistry and Center for Heterocyclic Compounds, University of Florida,
Gainesville, Florida 32611-2046, USA

(24. VI. 91)


tert-Alkyl sulfides, with an α -(1*H*-benzotriazol-1-yl) group **6** and **13**, are readily prepared from *N*-[(arylthio)methyl]-1*H*-benzotriazoles **3** and *N*-(**11**), respectively, by reaction with BuLi and then with the appropriate electrophile. The *tert*-alkyl sulfides **6** and **13** are smoothly converted by $\text{BF}_3 \cdot \text{OEt}_2$ into vinyl sulfides **5**, **7** or **12**, respectively, in satisfactory yields.

Introduction. – Vinyl sulfides are useful synthetic intermediates [1]. Routes to molecules having $\text{C}=\text{C}-\text{S}$ structural units are thus of considerable interest, and several have been described: *i*) Vinyl sulfides have been prepared by the direct treatment of cyclic ketones with thiols in the presence of strong acids (P_2O_5 , TsOH , AlCl_3) [2–5]; however, acyclic ketones give poor yields. *ii*) The addition of thiols to acetylenes in the presence of a base [6–9]. *iii*) The *Wittig* reaction of suitable phosphoranes with carbonyl compounds [10], as well as the related phosphonate olefins synthesis can lead to vinyl sulfides [11]. Thus, diethyl(methylthio)methyl phosphonate is alkylated by successive treatments with BuLi and an alkyl iodide to give the corresponding 1-(methylthio)alkyl-phosphonate esters. Their lithio derivatives react with aldehydes or ketones to form α -alkoxy-phosphonate adducts, which decompose upon heating at 50° (in THF) to form substituted methyl vinyl sulfides. Vinyl sulfides are accessible also through the reaction of (alkyldiene)triphenylphosphoranes with thioesters [12]. *iv*) Pyrolysis of thioacetals is a well-established route to vinyl sulfides [13]. An alternative procedure involves the conversion of dithioketals by using the $\text{Et}_2\text{Zn}/\text{CH}_2\text{I}_2$ reagent [5] [14].

Recently, we have reported [15] that *tert*-alkyl sulfides can be readily prepared by displacement of 1*H*-benzotriazole from α -benzotriazole sulfides with *Grignard* reagents. In the present paper, we report a new synthetic route to vinyl sulfides of the type $\text{Ar}-\text{S}-\text{C}=\text{C}$ by the elimination of 1*H*-benzotriazole from *tert*-alkyl sulfides.

Results and Discussion. – *Preparation of N-[α -(Arylthio)alkyl]-1H-benzotriazoles 6a–j and 13a–c.* Using the methodology previously developed in our laboratories, *tert*-alkyl sulfides of type **6** were prepared in good yield from 1-[(phenylthio)methyl]-1*H*-benzotriazole (**3a**) [16] or 1-[(4-methylphenylthio)methyl]-1*H*-benzotriazole (**3b**). The benzotriazoles **3a** and **3b** are readily available from 1-(chloromethyl)-1*H*-benzotriazole (**1**) and the appropriate thiophenol **2a**, **b**. Compounds **3a** and **3b** formed anions with BuLi at -78° (*Scheme 1*) which, with PhCH_2Br and 4-methylbenzyl chloride, gave the expected alkylated products **4a–d** in 83–89% yield (*Table 1*). Treatment of *sec*-alkyl sulfides **4a–d** with more BuLi readily afforded the α -lithio derivatives, which reacted with MeI, 1-iodoethane, 1-iodobutane, 1-bromoheptane, benzyl chloride, and 4-methylbenzyl bromide to provide the corresponding *tert*-alkyl sulfides **6a–j** with an α -benzotriazolyl

Scheme 1



5,7	Ar	R ¹	R ²	5,7	Ar	R ¹	R ²
5a	Ph	Ph	H	5h	<i>p</i> -CH ₃ -C ₆ H ₄	Ph	H
5b	Ph	Ph	Me	5i	<i>p</i> -CH ₃ -C ₆ H ₄	Ph	Ph
5c	Ph	Ph	Pr	5j	<i>p</i> -CH ₃ -C ₆ H ₄	<i>p</i> -CH ₃ -C ₆ H ₄	<i>p</i> -CH ₃ -C ₆ H ₄
5d	Ph	Ph	C ₆ H ₁₃	7a	Ph	Ph	CH ₃
5e	Ph	Ph	Ph	7b	Ph	Ph	C ₃ H ₇
5f	Ph	Ph	<i>p</i> -CH ₃ -C ₆ H ₄	7c	Ph	Ph	C ₆ H ₁₃
5g	Ph	<i>p</i> -CH ₃ -C ₆ H ₄	<i>p</i> -CH ₃ -C ₆ H ₄	7d	Ph	Ph	<i>p</i> -CH ₃ -C ₆ H ₄


Table 1. Preparation of the Sulfides **3**, **4**, **6**, **11**, and **13**

Compound (Formula)	Electrophile	Yield [%]	M.p. [°C] ^a	Solvent ^b
3b (C ₁₄ H ₁₃ N ₃ S)	—	80	101–103	EtOH
4a (C ₂₀ H ₁₇ N ₃ S)	PhCH ₂ Br	79	108–110 ^c	MeOH
4b (C ₂₁ H ₁₉ N ₃ S)	<i>p</i> -CH ₃ C ₆ H ₄ CH ₂ Cl	74	98–100	MeOH ^d
c (C ₂₁ H ₁₉ N ₃ S)	PhCH ₂ Br	81	60–62	E-PE
6a (C ₂₁ H ₁₉ N ₃ S)	MeI	88	100–102	MeOH
b (C ₂₂ H ₂₁ N ₃ S)	EtI	70	96–98	MeOH
c (C ₂₄ H ₂₅ N ₃ S)	BuI	86	110–112	MeOH
d (C ₂₇ H ₃₁ N ₃ S)	C ₇ H ₁₅ Br	86	83–85	MeOH
e (C ₂₄ H ₁₉ N ₃ S)	PhCH ₂ Br	83	138–140 ^e	MeOH
f (C ₂₈ H ₂₅ N ₃ S)	<i>p</i> -CH ₃ C ₆ H ₄ CH ₂ Cl	90	136–138	MeOH
g (C ₂₈ H ₂₇ N ₃ S)	<i>p</i> -CH ₃ C ₆ H ₄ CH ₂ Cl	80	111–113	MeOH
h (C ₂₂ H ₂₁ N ₃ S)	MeI	81	60–62	MeOH
i (C ₂₈ H ₂₅ N ₃ S)	PhCH ₂ Br	90	55–57	E-PE
j (C ₃₀ H ₂₉ N ₃ S)	<i>p</i> -CH ₃ C ₆ H ₄ CH ₂ Cl	82	65–67	E-PE ^e
11 (C ₁₉ H ₁₄ N ₃ S)	—	35 ^f	81–82	E
13a (C ₂₁ H ₁₈ N ₃ S)	EtI	75	126–128	MeOH
b (C ₂₃ H ₂₃ N ₃ S)	BuI	80	127–129	MeOH
c (C ₂₆ H ₂₁ N ₃ S)	PhCH ₂ Br	95	128–130	EtOH

^a) All compounds gave satisfactory elemental analyses.^b) E: Et₂O, PE: petroleum ether.^c) [16]: m.p. 103–105°.^d) Initially purified by column chromatography (hexane/CHCl₃ 2:1).^e) [16]: m.p. 140–142°.^f) The yield was calculated only for the 1*H* isomer.

group in good yields. Products **6a–j** were purified by recrystallization or column chromatography followed by recrystallization (Table 1).

In an alternative approach, condensation of benzotriazole (**8**), benzaldehyde (**9**), and benzenethiol (**10**) in the presence of catalytic TsOH in refluxing toluene with azeotropic removal of H₂O, formed α -(benzotriazolyl)alkyl sulfide **11** as the 1*H* isomer contaminated with a trace of the 2*H* isomer. The 1*H* isomer was easily separated from the mixture by recrystallization from Et₂O. The sulfide **11** was lithiated by BuLi at –78°, and the carbanion formed trapped with electrophiles to give *tert*-alkyl sulfides **13** (Table 1 and Scheme 2).



Synthesis of Vinyl Sulfides 5a–j, 7a–d, and 12a–c. The *tert*-alkyl sulfides with an α -(1*H*-benzotriazol-1-yl) group, **6a–j** and **13a–c**, were smoothly transformed into the corresponding vinyl sulfides on stirring in dry THF in the presence of an equivalent amount of BF₃·OEt₂ at 20° for 4 to 8 h. The vinyl sulfides were isolated in good yields (Table 2). In the cases of **6c**, **d** and **6f**, mixtures of two isomeric vinyl sulfides **5** and **7** were formed due to the presence of two different active α -H-atoms (Scheme 1 and Table 2).

Table 2. Preparation of Vinyl Sulfides **5**, **7**, and **12**

Compound ^{a)}	Substrat	Yield [%]	M.p [°C] ^{b)}
5a	6a	60	oil ^{c)}
b , 7a	6b	75	oil
c , 7b ^{d)}	6c	62	oil
d , 7c	6d	70	oil
e	6e	85	60–62
f , 7d	6f	83	54–56
g	6g	80	79–81
h	6h	52	oil
i	6j	91	71–73
j	6j	90	82–84
12a	13a	92	42–43 ^{e)}
b	13b	71	oil
c	13c	71	63–64 ^{f)}

^{a)} Vinyl sulfides **5** and **7** were obtained as a mixture of (*Z*)- and (*E*)-isomers.

^{b)} All compounds gave satisfactory elemental analyses.

^{c)} B.p. 137–140°; [17]: b.p. 138–140°.

^{d)} From compounds **6b–d** and **6f** mixtures of two types of vinyl sulfides **5** and **7** were obtained.

^{e)} [17]: m.p. 42–43°.

^{f)} [17]: m.p. 63–64°.

When $R^2 = H$ or $R^1 = R^2$, precursor **6** gave only a mixture of (*Z*)- and (*E*)-isomers of **5**. *tert*-Alkyl sulfides **13** gave vinyl sulfides **12** in good yield as (*Z/E*)-mixtures.

Compound **6f** was also converted into vinyl sulfide **5h** upon reaction with PhMgBr . After 24-h treatment of **6f** with NaH in boiling THF or toluene, ca. 50% of the starting material was recovered. HCl was also used for the transformation of **6** into vinyl sulfides. However, yields were lower than the BF_3 reactions due to the partial hydrolysis of **6** to the ketones.

Thus, the elimination of 1*H*-benzotriazole from *tert*-alkyl sulfides with an α -benzotriazolyl group in the presence of $\text{BF}_3 \cdot \text{OEt}_2$ provides a convenient route to the synthesis of a variety of vinyl sulfides in good-to-excellent yields.

The most appropriate method for the preparation of vinyl sulfides of the type described in this paper, was the *Emmons-Horner* phosphonate reaction [11], described in the *Introduction*. This also involves a three-step sequence with yields (ca. 50%) which are comparable with our method in several instances, but in some cases (acetophenone, cyclopentanone) only traces (up to 10%) of the products were formed.

¹H- and ¹³C-NMR Spectra. The structure of all the compounds were determined on the basis of their ¹H- and ¹³C-NMR spectra. It is interesting to note that in the ¹H spectra of *sec*-alkyl sulfides **4**, the two methylene protons and the N-CH proton connected to the asymmetric C-atom resonate as *ABX* systems. The CH_2 protons of benzylic substituents in **6** $R^1 \neq R^2$ resonate as two *doublets* with large geminal coupling constants of ca. 14–15 Hz. H-C(4) and H-C(7) of the 1-substituted 1*H*-benzotriazole moieties are shifted to ca. 7.95 and 7.60 ppm in the case of **4** and to ca. 8.30 and 8.10 ppm for **6**, respectively. The ¹³C-NMR spectra revealed typical patterns for 1-substituted 1*H*-benzotriazoles.

¹H-NMR Spectra of vinyl sulfides showed two *singlets* corresponding to the (*Z*)- and (*E*)-vinylic protons (ca. 6.80 and 6.70 ppm), clearly assignment of configuration from this is not possible. In the cases of vinyl sulfides of types **5** and **7**, the isomeric ratios were close to 1:1, but for compounds **12** the ratios were all ca. 4:1. ¹³C-NMR Spectra of vinyl sulfides also indicated (*Z*)- and (*E*)-isomer formation.

We are grateful to the USAID/AED officials Islamabad and Washington D.C. for financial support to A.S. Afridi and to the Education Department N. W. F. P. Pakistan for his leave.

Experimental Part

General. M.p.: determined on a hot-stage apparatus, uncorrected. ¹H- (300 MHz) and ¹³C- (75 MHz) NMR spectra were recorded on a *Varian-VXR-300-MHz* spectrometer; CDCl_3 as solvent, TMS as the internal standard. THF was distilled from sodium-benzophenone prior to use. Column chromatography was performed with MCB silica gel (230–400 mesh). 1-(Chloromethyl)-1*H*-benzotriazole (**1**) [18] and 1-[(phenylthio)methyl]-1*H*-benzotriazole (**3a**) [16] were prepared by the literature method quoted.

1-[(4-Methylphenyl)thio]methyl-1*H*-benzotriazole (**3b**). To an ice-cold soln. of 4-methylbenzenethiol (12.4 g, 0.1 mol) in MeOH (200 ml) was added Na metal (2.3 g, 0.1 mol). After complete dissolution of the metal, 1-(chloromethyl)-1*H*-benzotriazole (16.7 g, 0.1 mol) was added in small portions over 10 min. The soln. was allowed to warm to r.t. and stirred 6 h. The solvent was removed under reduced pressure to give a solid which was triturated with H_2O (3×50 ml). The resulting solid was collected and washed with 50% EtOH/ H_2O (2×50 ml) to give **3b** (20 g) (*Table 1*).

1-[(Phenyl)(phenylthio)methyl]-1*H*-benzotriazole (**11**). A mixture of 1*H*-benzotriazole (24 g, 0.2 mol), benzaldehyde (21.2 g, 0.2 mol), benzenethiol (20.6 ml, 0.2 mol), and TsOH (0.2 g) in toluene (200 ml) was refluxed for 14 h with azeotropic removal of H_2O . The mixture was stirred with 5% NaOH (500 ml), extracted with Et_2O (3×50 ml), washed with H_2O (3×50 ml), dried (MgSO_4), and the solvents were removed under reduced pressure to give an oily residue which was recrystallized from Et_2O (100 ml) (*Table 1*).

α -Substituted and α,α -Disubstituted Sulfides 4a–d, and 6a–j and 13a–c. Resp. General Procedure. To a soln. of the appropriate sulfide **3**, **4**, or **11**, (0.01 mol) in dry THF (30 ml), BuLi (0.01 mol in hexane) was added slowly via a syringe at -78° under N_2 . The mixture was stirred for 1 h at -78° , and a soln. of the electrophile (0.01 mol) in dry THF (5 ml) was slowly added. The resulting mixture was stirred for 2 h at -78° and for 15 h at r.t. The product was quenched with aq. NH_4Cl , extracted with Et_2O (3×30 ml), the Et_2O layer washed with H_2O dried ($MgSO_4$), and the solvent removed. The product was purified by recrystallization (Table 1).

Vinyl Sulfides 5a–j, 7a–d, and 12a–c. General Procedure. $BF_3 \cdot OEt_2$ (3 mmol) was added to a soln. of the appropriate *tert*-alkyl sulfide **6a–j** or **13a–c** (3 mmol) in dry THF (20 ml) and the product stirred at r.t. under N_2 for 2–5 h. The solvent was evaporated and the residue dissolved in Et_2O and washed with 10% Na_2CO_3 soln., H_2O , and dried ($MgSO_4$). The Et_2O was removed under reduced pressure and the oil purified by column chromatography (hexane) followed by recrystallization from MeOH.

REFERENCES

- [1] 'Organic Compounds of Sulphur, Selenium, and Tellurium', The Chemical Society, Burlington House, London, 1970–79, Vol 1–5.
- [2] D. J. Peterson, *J. Org. Chem.* **1968**, *33*, 780.
- [3] F. A. Carey, A. S. Court, *J. Org. Chem.* **1972**, *37*, 939.
- [4] B. M. Trost, D. E. Keeley, *J. Am. Chem. Soc.* **1976**, *98*, 248.
- [5] B. M. Trost, A. C. Lavoie, *J. Am. Chem. Soc.* **1983**, *105*, 5075.
- [6] R. A. Raphael, 'Acetylenic Compounds in Organic Synthesis', Academic Press, New York, 1955, pp. 43.
- [7] S. V. Amosova, N. I. Ivanova, B. A. Trofimov, *Zh. Org. Khim.* **1977**, *13*, 458.
- [8] R. N. Kudyakova, A. N. Volkov, *Zh. Org. Khim.* **1977**, *13*, 934.
- [9] S. R. Buzilova, I. D. Sadekov, T. V. Lipovich, T. M. Filippova, L. I. Vereschagin, *Zh. Obshch. Khim.* **1977**, *47*, 1999.
- [10] G. Wittig, M. Schlosser, *Chem. Ber.* **1961**, *94*, 1373.
- [11] E. J. Corey, J. I. Shulman, *J. Org. Chem.* **1970**, *35*, 777.
- [12] T. Kumamoto, K. Hosoi, T. Mukaiyama, *Bull. Chem. Soc. Jpn.* **1968**, *41*, 2742.
- [13] A. S. Atavin, B. A. Trofimov, A. I. Mikhaleva, I. P. Vasilev, *Izv. Akad. Nauk SSSR, Ser. Khim* **1972**, 2014.
- [14] A. D. Rodriguez, A. Nickon, *Tetrahedron* **1985**, *41*, 4443.
- [15] A. R. Katritzky, S. Perumal, W. Kuzmierkiewicz, P. Lue, J. V. Greenhill, *Helv. Chim. Acta* **1991**, *74*, 1924.
- [16] A. R. Katritzky, S. Rachwal, K. C. Caster, F. Mahni, K. W. Law, O. Rubio, *J. Chem. Soc., Perkin Trans. 1* **1987**, 781.
- [17] E. Campaigne, J. R. Leal, *J. Am. Chem. Soc.* **1954**, *76*, 1272.
- [18] J. H. Burckhalter, V. C. Stephens, L. A. R. Hall, *J. Am. Chem. Soc.* **1952**, *74*, 3868.