


This article was downloaded by: [McMaster University]

On: 14 May 2013, At: 13:37

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/gpss20>

Solvent-Free Acetylation of Thiols Under Catalysis of $\text{MgBr}_2 \cdot \text{OEt}_2$

Mohammad M. Mojtahedi^a, M. Saeed Abaee^a & Mashal Javadpour^a

^a Chemistry and Chemical Engineering Research Center of Iran, Tehran, Iran

Published online: 05 Nov 2010.

To cite this article: Mohammad M. Mojtahedi, M. Saeed Abaee & Mashal Javadpour (2010): Solvent-Free Acetylation of Thiols Under Catalysis of $\text{MgBr}_2 \cdot \text{OEt}_2$, Phosphorus, Sulfur, and Silicon and the Related Elements, 185:11, 2362-2365

To link to this article: <http://dx.doi.org/10.1080/10426501003645878>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.tandfonline.com/page/terms-and-conditions>

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae, and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand, or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

SOLVENT-FREE ACETYLATION OF THIOLS UNDER CATALYSIS OF $\text{MgBr}_2 \cdot \text{OEt}_2$

Mohammad M. Mojtahedi, M. Saeed Abaee,
and Mashal Javadpour

Chemistry and Chemical Engineering Research Center of Iran, Tehran, Iran

Solvent-free protection of aromatic and aliphatic thiols with acetic anhydride was performed at room temperature under trace quantities of magnesium bromide ethyl etherate, affording rapid formation of various thiol esters in excellent yields.

Keywords Acylation; catalysis; magnesium bromide ethyl etherate; solvent-free reaction; thiol esters


Thiol esters constitute an important class of organic compounds due to their bioorganic applications.¹ In addition they are key synthetic precursors in the preparation of heterocyclic compounds,² acyl radicals,³ asymmetric aldol products,⁴ and various ketones.⁵ Among available methods for the preparation of thiol esters,⁶ direct acylation of thiols with acetic anhydride or acetyl chloride is very common. This preparation is traditionally carried out in the presence of amine bases or protic acids. Recent developments include the use of various Lewis acids such as niobium(V) chloride,⁷ pyridine/alumina in combination with microwave irradiation,⁸ acetonitriletriphenylphosphonium bromide,⁹ ruthenium(III) chloride,¹⁰ copper-based reagents,¹¹ and aluminum dodecatungstophosphate.¹² However, many of these procedures usually require the use of environmentally unsafe conditions, demand application of expensive or commercially unavailable reagents, or involve the use of an additional stimulant.

Recently, we released a self-promoted chemoselective acylation of amines and thiols in the presence of no extra catalyst or additive.¹³ However, protection of thiols was limited to aromatic substrates and took relatively longer reaction time periods. Based on our previous experience on the use of magnesium bromide ethyl etherate ($\text{MgBr}_2 \cdot \text{OEt}_2$) as an efficient Lewis acid to catalyze various organic transformations under mild conditions,¹⁴ we envisioned that the acetylation of thiols could be boosted by using trace quantities of the catalyst. As a result, we report in this article a facile and room-temperature transformation of thiols to their corresponding acylated derivatives under very mild conditions (Scheme 1).

Received in final form on 15 January 2010.

















Partial financial support by the Ministry of Science, Research, and Technology of Iran is greatly acknowledged.

Address correspondence to Mohammad M. Mojtahedi and M. Saeed Abaee, Chemistry and Chemical Engineering Research Center of Iran, P.O. Box 14335-186, Tehran, Iran. E-mail: mojtahedi@ccerci.ac.ir; abaee@ccerci.ac.ir



Scheme 1

Table I MgBr₂·OEt₂-catalyzed acetylation of various thiols with Ac₂O

Entry	Thiol	Thiol ester	Mp (°C) found/reported	Yield % ^a
1	 1a	 2a	oil/oil ^{6c}	98
2	 1b	 2b	33–34/38–39 ^{6c}	90
3	 1c	 2c	oil/oil ^{6c}	82
4	 1d	 2d	48–49/50–51 ^{1b}	96
5	 1e	 2e	oil/oil ^{6f}	86
6	 1f	 2f	oil/oil ^{6f}	86
7	 1g	 2g	65–67/68–69 ^{11b}	88 ^b
8	 1h	 2h	oil/oil ^{6f}	86

^aIsolated yields.^bTwo equivalents of Ac₂O.

Results for the acetylation of various aliphatic and aromatic thiols are summarized in Table I. The synthesis of *S*-phenyl ethanethioate (entry 1) represents the typical procedure for solvent-free magnesium bromide ethyl etherate-catalyzed conversion of different thiols to their respective acetyl protected derivatives. An equimolar mixture of acetic anhydride and thiophenol in the presence of 4 mol% of the catalyst yields complete formation of the product **2a** in less than 1 h at room temperature under solventless conditions. The generality of the methodology is demonstrated by reacting other thiols with acetic anhydride under similar conditions. The course of the reactions were monitored by TLC and GC, products were easily separated in good yields from the reaction mixtures by simple aqueous workup, and the identity of the products was confirmed by comparing their spectroscopic data with those of known samples.

TYPICAL PROCEDURE

A mixture of acetic anhydride (5.1 mmol), a thiol (5.0 mmol), and magnesium bromide ethyl etherate (0.2 mmol, 4 mol%) was stirred in a flask at room temperature. The course of the reaction was monitored by TLC, and complete disappearance of the starting thiol was observed within 1 h. The mixture was dissolved in ether (20 mL) and washed twice by water (25 mL portions). The organic phase was dried over sodium sulfate, the volatile portion was removed at reduced pressure, and the product was isolated in good purity. All products are known, and their spectral and physical data are compared with those existing in the literature.

Selected Spectral Data

S-4-Chlorophenyl ethanethioate (2b). ^1H NMR (CDCl_3): δ 2.34 (s, 3H), 7.29 (s, 4H); MS: m/z (%) 186 (60) [M^+], 144 (100), 108 (36); IR (KBr): cm^{-1} 2929, 1713, 1092.

S-Naphthalen-2-yl ethanethioate (2d). ^1H NMR (CDCl_3): δ 2.45 (s, 3H), 7.62–8.00 (m, 7H); MS: m/z (%) 202 (30) [M^+], 160 (100), 115 (24); IR (KBr): cm^{-1} 3050, 1699, 1117.

S-Benzyl ethanethioate (2e). ^1H NMR (CDCl_3): δ 2.29 (s, 3H), 4.08 (s, 2H), 7.24 (s, 5H); MS: m/z (%) 166 (33) [M^+], 123 (26), 91 (100); IR (KBr): cm^{-1} 3030, 1694, 1133.

S,S'-Ethane-1,2-diyl diethanethioate (2g). ^1H NMR (CDCl_3): δ 2.45 (s, 6H), 2.96 (s, 4H); MS: m/z (%) 178 (13) [M^+], 119 (100), 103 (48); IR (KBr): cm^{-1} 1693, 1142.

REFERENCES

1. (a) Y. Hayashi, T. Itoh, and T. Fukuyama, *Org. Lett.*, **5**, 2235 (2003); (b) J. S. Hanker, L. Katzoff, H. R. Rosen, M. L. Seligman, H. Ueno, and A. M. Seligman, *J. Med. Chem.*, **9**, 288 (1966).
2. (a) J. Chen and C. J. Foursyth, *Org. Lett.*, **5**, 1281 (2003); (b) C. Brule, J.-P. Bouillon, E. Nicolai, and C. Portella, *Synthesis*, 436 (2003).
3. S. Ozaki, M. Adachi, S. Sekiya, and R. Kamikawa, *J. Org. Chem.*, **68**, 4586 (2003).
4. (a) J. M. Yost, G. Zhou, and D. M. Coltart, *Org. Lett.*, **8**, 1503 (2006); (b) G. Zhou, J. M. Yost, and D. M. Coltart, *Synthesis*, 478 (2007).
5. T. Shimizu and M. Seki, *Tetrahedron Lett.*, **43**, 1039 (2002).

6. (a) C. Lai and B. J. Backes, *Tetrahedron. Lett.*, **48**, 3033 (2007); (b) A. R. Katritzky, A. A. Shestopalov, and K. Suzuki, *Synthesis*, 1806 (2004); (c) M. Barbero, I. Degani, S. Dughera, and R. Fochi, *Synthesis*, 1225 (2003); (d) S. Kato and T. Murai, Thio derivatives, In *The Chemistry of Functional Groups, Supplement B, The Chemistry of Acid Derivatives*, S. Patai, ed. (Wiley, Chichester, UK, 1992), vol. 2, part 1, p. 803, 810ff; (e) W. Bauer and K. Kühlein, Thiocarbonsäure-S-Derivate, In *Methoden der Organischen Chemie (Houben-Weyl), Carbonsäuren und Carbonsäure-Derivate*, J., Falbe, ed. (Georg Thieme Verlag, Stuttgart, Germany, 1985), vol. E5, p. 832, 849ff; (f) Y.-Q. Cao, Y.-F. Du, B.-H. Chen, and Z. Dai, *J. Chem. Res.*, 762 (2004).
7. J. S. Yadav, A. V. Narsaiah, A. K. Basak, P. R. Goud, D. Sreenu, and K. Nagaiah, *J. Mol. Catal. A. Chem.*, **255**, 78 (2006).
8. S. Paul, P. Nanda, R. Gupta, and A. Loupy, *Tetrahedron Lett.*, **43**, 4261 (2002).
9. A. T. Khan, L. H. Choudhury, and S. Ghosh, *Eur. J. Org. Chem.*, 2782 (2005).
10. S. K. De, *Tetrahedron Lett.*, **45**, 2919 (2004).
11. (a) K. Jeyakumar and D. K. Chand, *J. Mol. Catal. A. Chem.*, **255**, 275 (2006); (b) K. Maekawa, K. Narasaka, and T. Mukaiyama, *Bull. Chem. Soc. Jpn.*, **46**, 3478 (1973).
12. H. Firouzabadi, N. Iranpoor, F. Nowrouzi, and K. Amani, *Chem. Commun.*, 764 (2003).
13. M. M. Mojtahedi, M. S. Abaee, M. M. Heravi, and F. K. Behbahani, *Monatsh. für Chem.*, **138**, 95 (2007).
14. (a) M. S. Abaee, R. Sharifi, and M. M. Mojtahedi, *Org. Lett.*, **7**, 5893 (2005); (b) M. S. Abaee, M. M. Mojtahedi, M. M. Zahedi, and R. Sharifi, *Heteroatom Chem.*, **18**, 44 (2006); (c) M. M. Mojtahedi, M. S. Abaee, and H. Abbasi, *Can. J. Chem.*, 429 (2006); (d) M. M. Mojtahedi, H. Abbasi, and M. S. Abaee, *J. Mol. Catal. A: Chem.*, **250**, 6 (2006); (e) M. S. Abaee, M. M. Mojtahedi, M. M. Zahedi, and G. Khanalizadeh, *Arkivoc*, 48 (2006); (f) M. S. Abaee, M. M. Mojtahedi, M. M. Zahedi, A. W. Mesbah, N. Mohaddes Ghandchi, and W. Massa, *Phosphorus, Sulfur, and Silicon*, **182**, 2891 (2007); (g) M. S. Abaee, M. M. Mojtahedi, M. M. Zahedi, R. Sharifi, and H. Khavasi, *Synthesis*, 3339 (2007).