


Substrate-Controlled Selectivity Switch in the Three-Component Coupling Involving Arynes, Aromatic Tertiary Amines, and CO₂

Sachin Suresh Bhojgude,^{†,§} Tony Roy,^{†,§} Rajesh G. Gonnade,[‡] and Akkattu T. Biju^{*,†,§}

[†]Organic Chemistry Division, [‡]Centre for Materials Characterization, CSIR-National Chemical Laboratory (CSIR-NCL), Dr. Homi Bhabha Road, Pune-411008, India

[§]Academy of Scientific and Innovative Research (AcSIR), New Delhi 110020, India

S Supporting Information




ABSTRACT: The transition-metal-free multicomponent coupling involving arynes, aromatic tertiary amines, and CO₂ is reported. The reaction exhibits switchable selectivity depending on the electronic nature of the aromatic amines used. With amines bearing electron-releasing/neutral groups as the nucleophilic trigger, the reaction afforded 2-arylamino benzoates via a nitrogen to oxygen alkyl group migration. Employing electron-deficient amines in the reaction furnished 2-aminoaryl benzoates proceeding via the aryl to aryl amino group migration resembling a Smiles rearrangement.

The efficient utilization of CO₂ as a one-carbon feedstock is one of the safest and most inexpensive methods for the synthesis of value-added products, as CO₂ is a cheap and readily available gas.¹ However, employing CO₂ as a C1 source in organic chemistry is limited mainly because of its high thermodynamic stability and kinetic inertness.² One way to explore the synthetic transformations using CO₂ involves the use of CO₂ along with high energy substrates/reactive intermediates.³ In this regard, employing CO₂ in reactions using highly electrophilic aryne intermediates constitutes a convenient method to install the CO₂ moiety directly to the aromatic ring.⁴ In 2006, Yoshida, Kunai and co-workers demonstrated, for the first time, the incorporation of CO₂ in aryne multicomponent coupling (MCC)⁵ triggered by imines for the straightforward synthesis of benzoxazinones.⁶ Subsequently, the same group uncovered the synthesis of anthranilic acids by the MCC involving arynes, secondary amines, and CO₂.⁷ We have reported the incorporation of CO₂ in aryne MCCs triggered by isocyanides for the practical synthesis of phthalimides.⁸ Moreover, Kobayashi and co-workers disclosed the Cu-catalyzed aryne MCCs involving terminal alkynes and CO₂ for the synthesis of isocoumarin derivatives.⁹ Apart from these works, the incorporation of CO₂ in aryne reactions remains underexplored.


In the context of our interest in developing transition-metal-free MCCs involving arynes,^{4a,10} we have recently reported the aryne MCCs triggered by aromatic tertiary amines using aldehydes/activated ketones as the third component (Scheme 1, eq 1).^{11,12} The reaction afforded *ortho*-functionalized tertiary

Scheme 1. Aryne MCCs Triggered by Aromatic Tertiary Amines

MCCs involving arynes, tertiary amines and aldehydes/activated ketones



Tertiary amine-controlled switchable selectivity in CO₂ incorporated aryne MCCs




amines proceeding via the aryl to aryl amino group migration similar to the Smiles rearrangement.¹³ Inspired by this finding, we envisaged the use of CO₂ in aryne MCCs triggered by aromatic tertiary amines. When the reaction was performed using electron-releasing/neutral tertiary amines as the nucleophilic trigger, 2-arylamino benzoates are formed via a unique nitrogen to oxygen alkyl group migration (eq 2). Interestingly, with electron-deficient tertiary amines, the reaction afforded 2-

Received: September 21, 2016

aminoaryl benzoates via an aryl to aryl amino group migration (mechanistically similar to Smiles rearrangement).¹⁴ Herein, we report this substrate-controlled switchable selectivity observed in the aryne MCCs involving CO₂ and aromatic tertiary amines. Notably, related anthranilic acid derivatives are endowed with interesting biological properties.¹⁵

Our present study commenced with the treatment of an aryne generated in situ from 2-(trimethylsilyl) aryl triflate **1a**¹⁶ (using KF and 18-crown-6) with *N,N*-dimethyl aniline **2a** in THF under an atmosphere of CO₂ (balloon). Under these conditions, we expected the 2-aminoaryl benzoate derivative **4** formed by the aryl to aryl NMe₂ migration.^{11a} But delightfully, the 2-arylamino benzoate **3a** was isolated in 87% yield (Scheme 2). The product **3a** was formed by the *N*-arylation of **2a**

Scheme 2. MCC Involving Aryne, *N,N*-Dimethylaniline, and CO₂




followed by quenching of the aryl anion using CO₂ and a subsequent nitrogen to oxygen methyl group migration.¹⁷ The reactions performed using other common fluoride sources such as CsF and tetrabutylammonium fluoride (TBAF) resulted in a reduced yield of **3a**.

With optimized conditions in hand, we then examined the substrate scope with respect to the aryne and aniline substrates (Scheme 3). A series of *N,N*-dimethyl anilines with differently substituted electron-releasing/neutral/moderately electron-poor groups are tolerated under the reaction conditions furnishing the 2-arylamino benzoate derivatives in good yields (**3b–j**). Moreover, the NMe₂ moiety present in donor–acceptor system **2k** furnished the MCC product in 56% yield (**3k**) and leuco-malachite green underwent efficient twofold MCC with a methyl group transfer leading to the formation of desired product **3l** in 84% yield. Additionally, this alkyl group migration reaction is not limited to *N,N*-dimethyl aniline derivatives; delightfully, *N,N*-diethyl aniline **2m** and *N*-benzyl-*N*-methyl aniline **2n** resulted in the ethyl and benzyl group migration respectively to afford the target products in high yields (**3m**, **3n**) thus demonstrating the versatility of the present method. Furthermore, electronically diverse 4,5-disubstituted symmetrical arynes generated from their precursors readily afforded the 1,2-disubstituted arene derivatives in good yields (**3o–q**). It is noteworthy that both symmetrical and unsymmetrical naphthalene worked well and the corresponding MCC products are formed in good yields (**3r**, **3s**). Finally, the reaction of **2a** and CO₂ with 4-methylbenzyl generated from the triflate precursor resulted in the formation of an inseparable mixture of regioisomers (**3t** and **3t'**) in 64% yield and 1.4:1 selectivity.

Interestingly, treatment of 4-carboethoxy *N,N*-dimethyl aniline with aryne generated from **1a** (using KF and 18-crown-6) under an atmosphere of CO₂ (balloon) –10 °C afforded the *ortho*-functionalized tertiary amine **4a** in 75% yield (Scheme 4). The product formation took place via the aryl to aryl migration of the NMe₂ group and is analogous to the Smiles rearrangement. This rearrangement reaction worked

Scheme 3. Scope of Aryne MCCs for the Synthesis of 2-Arylamino Benzoates^a




^aGeneral conditions: **1** (0.6 mmol), **2** (0.5 mmol), KF (1.2 mmol), 18-crown-6 (1.2 mmol), THF (2.0 mL), under an atmosphere of CO₂ (balloon), –10 °C to rt, 12 h, then stirred at 60 °C for 12 h. Yields of the isolated products are given. ^bReaction stirred at 100 °C for 12 h instead of 60 °C. ^cRun on 0.25 mmol scale. ^dThe regioisomer ratio was determined by ¹H NMR analysis.

well with various electron-poor aromatic tertiary amines, and in all cases, the desired product was formed in moderate to good yields (**4b–e**). In the case of 4-trifluoromethyl derivative **4b**, the structure was confirmed by single-crystal X-ray analysis.¹⁸ In addition, this MCC worked well with electronically different symmetrically substituted aryne precursors, and the expected Smiles rearrangement products were isolated in good yields (**4f–i**). Notably, the unsymmetrical naphthalene generated from the triflate precursor afforded a single regioisomer **4j** in 62% yield. Besides, attempted CO₂ incorporation MCC with 4-methylbenzyl furnished an inseparable mixture of regioisomers (**4k** and **4k'**) in 61% yield with 1.7:1 selectivity.

To probe the mechanism of the alkyl group migration, a series of experiments were performed. The reaction of **2a** with the aryne generated from **1a** and CO₂ followed by quenching the reaction mixture with TfOH resulted in formation of the ammonium salt **5.HOTf** (Scheme 5, eq 3). Treatment of **5.HOTf** with K₂CO₃ afforded the methyl ester **3a** in 25% yield (two steps). This indicates the intermediacy of the ammonium salt in the reaction. To gain insight into the nature of the alkyl group migration, we performed several mechanistic experiments.¹⁷ The reaction of the aryne generated from **1a** and CO₂ with **2a** and **2m** afforded a mixture of four products (**3a**, **3m**, **3u**, and **3v**) in the ratio 26:23:23:28 (eq 4). The products **3a** and **3m** could be formed by the intramolecular alkyl group migration. Interestingly, the formation of **3u** and **3v** clearly


Scheme 4. Scope of Aryne MCCs Involving CO₂ and Electron-Poor Tertiary Amines^a




^aGeneral conditions: **1** (0.6 mmol), **2** (0.5 mmol), KF (1.2 mmol), 18-crown-6 (1.2 mmol), THF (2.0 mL), under an atmosphere of CO₂ (balloon), -10 °C, 24 h. Yields of the isolated products are given. ^bReaction run at -10 °C to rt for 12 h. ^cGiven is ¹H NMR yield. ^dThe regioisomer ratio was determined by ¹H NMR analysis.

Scheme 5. Mechanistic Experiments


Isolation of the intermediate



Cross-over experiment




Experiment using *N*-ethyl-*N*-methylaniline



indicates the possibility of the intermolecular alkyl group migration in the present reaction. Moreover, performing the reaction using *N*-ethyl-*N*-methylaniline **6** under the present reaction conditions furnished four products (**3a**, **3u**, **3v**, and **3m**) in the ratio 23:18:33:26 shedding further light on the possibility of the intermolecular nature of the present alkyl group migration (eq 5).

The mechanism of this substrate-controlled switchable selectivity in aryne MCCs can be elucidated as follows (Scheme 6). The initial addition of aromatic tertiary amine **2** to the aryne


Scheme 6. Tentative Mechanism of the Substrate-Controlled Switchable MCCs



formed from **1** generates the 1,3-zwitterionic intermediate **7**. In the absence of a proton source, the aryl anion intermediate **7** adds to CO₂ generating the key dimethyl phenyl ammonium benzoate intermediate **5**. With electron-rich/-neutral amines, the intermediate **5** could undergo a methyl group transfer to furnish the 2-arylamino benzoate **3**. However, in the case of tertiary amines having an electron-poor group attached, the intermediate **5** could undergo an intramolecular nucleophilic aromatic substitution reaction (S_NAr) resulting in the aryl to aryl NMe₂ group migration leading to the formation of 2-aminoaryl benzoate **4** via the σ -complex **8**.

Furthermore, the synthetic utility of the present alkyl group migration MCC has been demonstrated in the transition-metal-free synthesis of biologically important acridones and benzoxazinones. 2-Arylamino benzoate **3a** was converted to 2-arylamino benzaldehyde **9** by a reduction–oxidation sequence. Direct aryl-aldehyde C^{sp}2–C^{sp}2 bond formation via PhI(OAc)₂-mediated intramolecular cross-dehydrogenative coupling of **9** furnished the acridone **10** in 49% yield (Scheme 7, eq

Scheme 7. Synthetic Utility of 2-Arylamino Benzoates



6).¹⁹ In addition, benzoate **3a** was hydrolyzed to acid **11** under basic conditions followed by intramolecular C^{sp}3–O bond formation has been accomplished in the presence of PhI(OAc)₂/NaN₃ leading to the formation of benzoxazinone **12** in 96% yield (eq 7).²⁰

In conclusion, we have demonstrated switchable selectivity in the aryne MCCs involving aromatic tertiary amines and CO₂. The selectivity depends on the electronic nature of the aromatic amine used. The reaction of aromatic amines bearing electron-releasing/-neutral groups, arynes, and CO₂ furnished 2-arylamino benzoates via a nitrogen to oxygen alkyl group migration. Interestingly, the selectivity was switched to the

formation of 2-aminoaryl benzoates (via an aryl to aryl amino group migration) employing electron-deficient amines as nucleophiles in aryne MCCs using CO₂ as the third component. A reasonable mechanism for the switchable reactions is provided. Further studies on related aryne MCCs using CO₂ are ongoing in our laboratory.

■ ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: [10.1021/acs.orglett.6b02845](https://doi.org/10.1021/acs.orglett.6b02845).

Details on experimental procedures, characterization data of all compounds (PDF)

X-ray data for **4b** (CIF)

■ AUTHOR INFORMATION

Corresponding Author

*E-mail: at.biju@ncl.res.in.

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

Generous financial support by Science and Engineering Research Board (SERB-DST), Government of India (Grant No. SR/S1/OC/12/2012) is kindly acknowledged. S.S.B. thanks CSIR for the fellowship, and T.R. acknowledges DST for the Inspire fellowship. We thank Dr. P. R. Rajamohanan (CSIR-NCL) for the excellent NMR support and Dr. B. Santhakumari (CSIR-NCL) for the HRMS data.

■ REFERENCES

- (1) For selected reviews on the use of CO₂ as a C1 source, see: (a) Liu, Q.; Wu, L.; Jackstell, R.; Beller, M. *Nat. Commun.* **2015**, *6*, 5933. (b) Kondratenko, E. V.; Mul, G.; Baltrusaitis, J.; Larrazábal, G. O.; Pérez-Ramírez, J. *Energy Environ. Sci.* **2013**, *6*, 3112. (c) Zhang, L.; Hou, Z. *Chem. Sci.* **2013**, *4*, 3395. (d) Cokoja, M.; Bruckmeier, C.; Rieger, B.; Herrmann, W. A.; Kühn, F. E. *Angew. Chem., Int. Ed.* **2011**, *50*, 8510. (e) Sakakura, T.; Choi, J.-C.; Yasuda, H. *Chem. Rev.* **2007**, *107*, 2365.
- (2) For the selected recent reports, see: (a) Anaby, A.; Feller, M.; Ben-David, Y.; Leitun, G.; Diskin-Posner, Y.; Shimon, L. J. W.; Milstein, D. *J. Am. Chem. Soc.* **2016**, *138*, 9941. (b) Börjesson, M.; Moragas, T.; Martin, R. *J. Am. Chem. Soc.* **2016**, *138*, 7504. (c) Nogi, K.; Fujihara, T.; Terao, J.; Tsuji, Y. *J. Am. Chem. Soc.* **2016**, *138*, 5547. (d) Shao, P.; Wang, S.; Chen, C.; Xi, C. *Org. Lett.* **2016**, *18*, 2050. (e) Moragas, T.; Gaydou, M.; Martin, R. *Angew. Chem., Int. Ed.* **2016**, *55*, 5053. (f) Fenner, S.; Ackermann, L. *Green Chem.* **2016**, *18*, 3804.
- (3) (a) Yu, D.; Teong, S. P.; Zhang, Y. *Coord. Chem. Rev.* **2015**, 293–294, 279. (b) Tsuji, Y.; Fujihara, T. *Chem. Commun.* **2012**, 48, 9956. (c) Huang, K.; Sun, C.; Shi, Z. *Chem. Soc. Rev.* **2011**, *40*, 2435.
- (4) For recent reviews on aryne chemistry, see: (a) Bhojgude, S. S.; Bhunia, A.; Biju, A. T. *Acc. Chem. Res.* **2016**, *49*, 1658. (b) Goetz, A. E.; Shah, T. K.; Garg, N. K. *Chem. Commun.* **2015**, *51*, 34. (c) Dubrovskiy, A. V.; Markina, N. A.; Larock, R. C. *Org. Biomol. Chem.* **2013**, *11*, 191. (d) Pérez, D.; Peña, D.; Guitián, E. *Eur. J. Org. Chem.* **2013**, 5981. (e) Wu, C.; Shi, F. *Asian J. Org. Chem.* **2013**, *2*, 116. (f) Tadross, P. M.; Stoltz, B. M. *Chem. Rev.* **2012**, *112*, 3550. (g) Gampe, C. M.; Carreira, E. M. *Angew. Chem., Int. Ed.* **2012**, *51*, 3766. (h) Bhunia, A.; Yetra, S. R.; Biju, A. T. *Chem. Soc. Rev.* **2012**, *41*, 3140. (i) Okuma, K. *Heterocycles* **2012**, *85*, 515. (j) Yoshida, H.; Ohshita, J.; Kunai, A. *Bull. Chem. Soc. Jpn.* **2010**, *83*, 199. (k) Sanz, R. *Org. Prep. Proced. Int.* **2008**, *40*, 215.
- (5) (a) Bhojgude, S. S.; Biju, A. T. *Angew. Chem., Int. Ed.* **2012**, *51*, 1520 and references cited therein. See also: (b) Bhunia, A.; Biju, A. T. *Synlett* **2014**, 25, 608.
- (6) Yoshida, H.; Fukushima, H.; Ohshita, J.; Kunai, A. *J. Am. Chem. Soc.* **2006**, *128*, 11040.
- (7) Yoshida, H.; Morishita, T.; Ohshita, J. *Org. Lett.* **2008**, *10*, 3845.
- (8) (a) Kaicharla, T.; Thangaraj, M.; Biju, A. T. *Org. Lett.* **2014**, *16*, 1728. See also: (b) Fang, Y.; Wang, S.-Y.; Ji, S.-J. *Tetrahedron* **2015**, *71*, 2768.
- (9) Yoo, W.-J.; Nguyen, T. V. Q.; Kobayashi, S. *Angew. Chem., Int. Ed.* **2014**, *53*, 10213.
- (10) (a) Roy, T.; Thangaraj, M.; Gonnade, R. G.; Biju, A. T. *Chem. Commun.* **2016**, 52, 9044. (b) Roy, T.; Bhojgude, S. S.; Kaicharla, T.; Thangaraj, M.; Garai, B.; Biju, A. T. *Org. Chem. Front.* **2016**, *3*, 71. (c) Roy, T.; Baviskar, D. R.; Biju, A. T. *J. Org. Chem.* **2015**, *80*, 11131. (d) Bhunia, A.; Kaicharla, T.; Porwal, D.; Gonnade, R. G.; Biju, A. T. *Chem. Commun.* **2014**, 50, 11389. (e) Bhunia, A.; Roy, T.; Gonnade, R. G.; Biju, A. T. *Org. Lett.* **2014**, *16*, 5132. (f) Bhunia, A.; Roy, T.; Pachfule, P.; Rajamohanan, P. R.; Biju, A. T. *Angew. Chem., Int. Ed.* **2013**, *52*, 10040. (g) Bhunia, A.; Porwal, D.; Gonnade, R. G.; Biju, A. T. *Org. Lett.* **2013**, *15*, 4620.
- (11) (a) Bhojgude, S. S.; Baviskar, D. R.; Gonnade, R. G.; Biju, A. T. *Org. Lett.* **2015**, *17*, 6270. For a related *N*-arylation of tertiary amines using arynes, see: (b) Bhojgude, S. S.; Kaicharla, T.; Biju, A. T. *Org. Lett.* **2013**, *15*, 5452.
- (12) For the application of related MCCs for the synthesis of 9- and 10-membered oxaza heterocycles, see: Okuma, K.; Kinoshita, H.; Nagahora, N.; Shioji, K. *Eur. J. Org. Chem.* **2016**, 2264.
- (13) For a related aryne reaction proceeding via the Smiles rearrangement, see: (a) Holden, C. M.; Sohel, S. M. A.; Greaney, M. F. *Angew. Chem., Int. Ed.* **2016**, *55*, 2450. For a related aryne reaction proceeding via migration of the aryl group from sulfur to nitrogen, see: (b) Yoshida, S.; Yano, T.; Misawa, Y.; Sugimura, Y.; Igawa, K.; Shimizu, S.; Tomooka, K.; Hosoya, T. *J. Am. Chem. Soc.* **2015**, *137*, 14071.
- (14) For reviews, see: (a) Snape, J. *Chem. Soc. Rev.* **2008**, *37*, 2452. (b) Truce, W. E.; Kreider, E. M.; Brand, W. W. *Org. React.* **1970**, *18*, 99.
- (15) (a) Merk, D.; Lamers, C.; Weber, J.; Flesch, D.; Gabler, M.; Proschak, E.; Schubert-Zsilavec, M. *Bioorg. Med. Chem.* **2015**, *23*, 499. (b) Walsh, C. T.; Haynes, S. W.; Ames, B. D. *Nat. Prod. Rep.* **2012**, *29*, 37.
- (16) (a) Himeshima, Y.; Sonoda, T.; Kobayashi, H. *Chem. Lett.* **1983**, *12*, 1211. See also: (b) Peña, D.; Cobas, A.; Pérez, D.; Guitián, E. *Synthesis* **2002**, 1454.
- (17) For details, see the Supporting Information.
- (18) CCDC 1503071 (**4b**) contains the supplementary crystallographic data for this paper. These data can be obtained free of charge from Data Centre via www.ccdc.cam.ac.uk/data_request/cif.
- (19) Zheng, Z.; Dian, L.; Yuan, Y.; Zhang-Negrerie, D.; Du, Y.; Zhao, K. *J. Org. Chem.* **2014**, *79*, 7451.
- (20) Zhang, N.; Cheng, R.; Zhang-Negrerie, D.; Du, Y.; Zhao, K. *J. Org. Chem.* **2014**, *79*, 10581.