

One-Pot Three-Component Approach to the Synthesis of Polyfunctional Pyrazoles


Guolin Zhang,* Hangcheng Ni, Wenteng Chen, Jiaan Shao, Huan Liu, Binhui Chen, and Yongping Yu*

Zhejiang Province Key Laboratory of Anti-Cancer Drug Research, College of Pharmaceutical Science, Zhejiang University, Hangzhou 310058, P. R. China

yyu@zju.edu.cn; guolinzhang@zju.edu.cn.

Received September 29, 2013

ABSTRACT



A simple, multicomponent, and straightforward reaction of vinyl azide, aldehyde, and tosylhydrazine affords the construction of 3,4,5-trisubstituted 1H-pyrazoles regioselectively in the presence of base with moderate to excellent yields. A range of functionality could be tolerated in this methodology, and a possible mechanism is proposed.

Pyrazoles are one of the most important heterocyclic compounds that possess a range of biological activities, such as antibacterial,¹ antileukemic,² antitumor,³ antiobesity,⁴ and anti-inflammatory.⁵ Polysubstituted pyrazoles have also been utilized as useful ligands for some cross-coupling reactions.⁶ Due to pyrazoles' important utility, numerous approaches have been developed over the past decades including the reaction of 1,3-diketones with hydrazine and hydrazine derivatives,⁷ the coupling of *N*-arylhydrazones and nitro-olefins,⁸ and 1,3-dipolar cycloaddition of hydrazines to alkynes.⁹ However, it is still challenging to prepare polysubstituted pyrazoles.

Recently, vinyl azides have emerged as important synthons although they were once regarded as “sleepers” of the reactive azido species for a long time. In recent years, our group¹⁰ and others¹¹ have developed numerous nitrogen-containing heterocycles from vinyl azides. Based on the previous result, herein, we demonstrate a simple, multicomponent, and straightforward approach to provide 3,4,5-trisubstituted 1H-pyrazoles regioselectively in the presence of base with good to excellent yields from vinyl azides, benzaldehyde, and tosylhydrazine. The azido group in vinyl azides serves as a leaving group.¹²

(1) Tanitame, A.; Oyamada, Y.; Ofuji, K.; Fujimoto, M.; Iwai, N.; Hiyama, Y.; Suzuki, K.; Ito, H.; Terauchi, H.; Kawasaki, M.; Nagai, K.; Wachi, M.; Yamagishi, J. *J. Med. Chem.* **2004**, *47*, 3693.

(2) Baraldi, P. I. G.; Cozzi, P.; Geroni, C.; Mongelli, N.; Romagnoli, R.; Spalluto, G. *Bioorg. Med. Chem.* **1999**, *7*, 251.

(3) Havrylyuk, D.; Zimenkovsky, B.; Vasilenko, O.; Gzella, A.; Lesyk, R. *J. Med. Chem.* **2012**, *55*, 8630.

(4) Rinaldi-Carmona, M.; Barth, F.; Heaulme, M.; Shire, D.; Calandra, B.; Congy, C.; Martinez, S.; Maruani, J.; Neliat, G.; Caput, D.; Ferrara, P.; Soubrie, P.; Brelriere, J. C.; Fur, G. L. *FEBS Lett.* **1994**, *350*, 240.

(5) El-Sayed, M. A.; Abdel-Aziz, N. I.; Abdel-Aziz, A. M.; El-Azab, A. S.; El-Tahir, K. E. *Bioorg. Med. Chem.* **2012**, *20*, 3306.

(6) (a) Singer, R. A.; Caron, S.; McDermott, R. E.; Arpin, P.; Do, N. M. *Synthesis* **2003**, *11*, 1727. (b) Singer, R. A.; Dore, M.; Sieser, J. E.; Berliner, M. A. *Tetrahedron Lett.* **2006**, *47*, 3727.

(7) (a) Kost, A. N.; Grandberg, I. I. *Adv. Heterocycl. Chem.* **1966**, *6*, 347. (b) Jiang, J. A.; Huang, W. B.; Zhai, J. J.; Liu, H. W.; Cai, Q.; Xu, L. X.; Wang, W.; Ji, Y. F. *Tetrahedron* **2013**, *69*, 627. (c) Fustero, S.; Roman, R.; Sanz-Cervera, J. F.; Simon-Fuentes, A. X.; Cunat, A. C.; Villanova, S.; Murguia, M. *J. Org. Chem.* **2008**, *73*, 3523.

(8) Deng, X.; Mani, N. S. *J. Org. Chem.* **2008**, *73*, 2412.

(9) (a) Kirkham, J. D.; Edeson, S. J.; Stokes, S.; Harrity, J. P. A. *Org. Lett.* **2012**, *14*, 5354. (b) Zora, M.; Kivrak, A. *J. Org. Chem.* **2011**, *76*, 9379. (c) Ahmed, M. S.; Kobayashi, K.; Mori, A. *Org. Lett.* **2005**, *7*, 4487. (e) Alex, K.; Tillack, A.; Schwarz, N.; Beller, M. *Org. Lett.* **2008**, *10*, 2377.

(10) (a) Chen, W.; Hu, M.; Wu, J.; Zou, H.; Yu, Y. *Org. Lett.* **2010**, *12*, 3863. (b) Hu, M.; Wu, J.; Zhang, Y.; Qiu, F.; Yu, Y. *Tetrahedron* **2011**, *67*, 2676. (c) Shao, Z.; Pan, Q.; Chen, J.; Yu, Y.; Zhang, G. *Tetrahedron* **2012**, *68*, 6565. (d) Shao, J.; Yu, W.; Shao, Z.; Yu, Y. *Chem. Commun.* **2012**, *48*, 2785. (e) Yu, W.; Chen, W.; Liu, S.; Shao, J.; Shao, Z.; Lin, H.; Yu, Y. *Tetrahedron* **2013**, *69*, 1953. (f) Zhang, Y.; Liu, S.; Yu, W.; Hu, M.; Zhang, G.; Yu, Y. *Tetrahedron* **2013**, *69*, 2070.

(11) (a) For a review of vinyl azides, see: Jung, N.; Brase, S. *Angew. Chem., Int. Ed.* **2012**, *51*, 12169. (b) Xie, H.; Liu, J.; Wu, L.; Ding, M. *Tetrahedron* **2012**, *68*, 7984. (c) Ng, E. P. J.; Wang, Y.; Hui, B. W. Q.; Lapointe, G.; Chiba, S. *Tetrahedron* **2011**, *67*, 7728. (d) Hu, B.; Wang, Z.; Ai, N.; Zheng, J.; Liu, X.; Shan, S.; Wang, Z. *Org. Lett.* **2011**, *13*, 6362.

(12) For a vinyl azide serving as a leaving group, see: (a) Zou, H.; Zhu, H.; Shao, J.; Wu, J.; Chen, W.; Giulianotti, M. A.; Yu, Y. *Tetrahedron* **2011**, *67*, 4887. (b) Chen, W.; Shao, J.; Li, Z.; Giulianotti, M. A.; Yu, Y. *Can. J. Chem.* **2012**, *90*, 214.

Initially, vinyl azide **1a**, benzaldehyde **2a**, and tosylhydrazine **3** were selected as model reagents to optimize the reaction conditions (Table 1). No corresponding product was observed in the absence of base (Table 1, entry 1). However, the expected transformation occurred in the presence of 1.5 equiv of NaOH at 60 °C. The desired compound **4a** was obtained as the major product with 49% yield (Table 1, entry 2). Gratifyingly, the yield was notably increased when more equivalents of base were added into the reaction mixture (Table 1, entries 2–5). A perfect isolated yield was obtained when the reaction was performed in the presence of 5 equiv of NaOH in 2 h (Table 1, entry 4), while continuously increasing the amount of base (7 equiv) did not give a significant change (Table 1, entry 5).

Further, other bases were also tested for the reaction, which revealed that NaOH was the most efficient one (Table 1, entry 8–13), while DMF was considered as the most preponderant solvent (Table 1, entries 14–17). Temperature screening showed that 60 °C was optimal to obtain the maximum yield of the product. On the basis of the above studies, the most favorable reaction conditions for the

Table 1. Optimization of Reaction Conditions^a


entry	solvent	base (equiv)	t (h)	conversion (%) ^b
1	DMF	—	12	n.r
2	DMF	NaOH(1.5)	12	49
3	DMF	NaOH(3)	2	74
4	DMF	NaOH(5)	2	89 (83)^c
5	DMF	NaOH(7)	2	88
6	DMF	NaOH(5)	5	71 ^d
7	DMF	NaOH(5)	1	85 ^e
8	DMF	KOH(5)	2	85
9	DMF	^t BuOK(5)	2	54
10	DMF	Cs ₂ CO ₃ (5)	2	75
11	DMF	NaOMe(5)	12	49
12	DMF	DABCO(5)	12	trace
13	DMF	DBU(5)	2	n.r
14	dioxane	NaOH(5)	2	46
15	toluene	NaOH(5)	2	74
16	ethanol	NaOH(5)	2	25
17	acetonitrile	NaOH(5)	2	86

^a Reaction conditions: vinyl azide (0.25 mmol, 1.0 equiv), benzaldehyde (0.25 mmol, 1.0 equiv), tosylhydrazine (0.25 mmol, 1.0 equiv), 2 mL of solvent, 60 °C. n.r. = no reaction. ^b Determined by HPLC, based on the disappearance of the starting vinyl azide. The most successful entry is highlighted in bold. ^c Isolated yield. ^d The reaction temperature is 40 °C. ^e The reaction temperature is 80 °C.

formation of **4a** were established. The vinyl azides were readily prepared according to literature¹³ (Scheme 1).


Scheme 1. Synthesis of Vinyl Azides

Scheme 1A



reaction condition: a) Br₂, Et₂O, 0 °C, 2 h; b) NaN₃, DMF, rt, 3 h

Scheme 1B



reaction condition: a) EtONa, EtOH-THF, -15 °C

Table 2. Reactions Scope of Vinyl Azides, Aldehyde, and Tosylhydrazine^a

entry	R ₁	R ₂	R ₃	product	yield (%) ^b
1	Ph	Ph-CO	Ph	4a	83
2	Ph	Ph-CO	4-MePh	4b	78
3	Ph	Ph-CO	4-BrPh	4c	89
4	Ph	Ph-CO	3-BrPh	4d	85
5	Ph	Ph-CO	4-NO ₂ Ph	4e	92
6	Ph	Ph-CO	4-pyridinyl	4f	82
7	Ph	Ph-CO	piperonyl	4g	63
8	Ph	4-MePh-CO	4-MePh	4h	74
9	Ph	4-MePh-CO	Ph	4i	74
10	Ph	4-ClPh-CO	Ph	4j	87
11	4-BrPh	Ph-CO	Ph	4k	88
12	Ph	4-ClPh-CO	4-NO ₂ Ph	4l	95
13	Ph	4-MePh-CO	3-BrPh	4m	91
14	Ph	4-MePh-CO	4-pyridinyl	4n	71
15	Ph	COOEt	4-NO ₂ Ph	4o	75 ^c
16	4-OMePh	COOEt	4-NO ₂ Ph	4p	59 ^c
17	4-BrPh	COOEt	4-NO ₂ Ph	4q	69 ^c
18	4-NO ₂ Ph	COOEt	4-NO ₂ Ph	4r	77 ^c
19	<i>n</i> -propyl	COOEt	4-NO ₂ Ph	4s	54 ^c
20	H	Ph	Ph	4t	56 ^{c,d}
21	H	Ph	4-MePh	4u	63 ^{c,d}
22	Ph	4-ClPh-CO	<i>n</i> -propyl	—	—

^a Reactions were carried out in DMF (2.0 mL) with **1** (0.25 mmol, 1.0 equiv), **2** (0.25 mmol, 1.0 equiv), **3** (0.25 mmol, 1.0 equiv), NaOH (1.25 mmol, 1.0 equiv), 60 °C. ^b Isolated yield. ^c Reactions were carried out with 5.0 equiv of Cs₂CO₃. ^d Reaction temperature was 80 °C for 12 h.

Based on the optimal conditions, the scope of the reaction was studied using a series of vinyl azides **1** and aldehydes.

(13) (a) Gilchrist, T. L.; Mendonca, R. *ARKIVOC* **2000**, 769. (b) Chiba, S.; Wang, Y.; Lapointe, G.; Narasaka, K. *Org. Lett.* **2008**, 10, 313.

As shown in Table 2, various substituted vinyl azides worked well with aldehydes and hydrazide to provide the corresponding 3,4,5-trisubstituted 1*H*-pyrazoles with moderate to excellent yields. The reaction could tolerate vinyl azides with aromatic substitution and aldehydes with various steric and electronic properties.

At the R₃ position, various groups on the aryl ring were compatible with the reaction. Notably, aldehydes bearing an electron-withdrawing group at the aryl ring afforded the desired products with excellent efficiency (Table 2, entry 5). We also found that the reaction could tolerate heteroaryl motifs (Table 2, entry 6). However, no desired product was observed when butyraldehyde was employed (Table 2, entry 22). It was most likely due to the instability of the hydrazone intermediate generated from butyraldehyde.

Instead of arylcarbonyl substituted vinyl azides, the reaction of vinyl azide (Table 2, entries 15–19) bearing an ester group at the α-position gave the desired products with moderate yields. Meanwhile, vinyl azide with an alkyl group at the R₁ position can also tolerate the reaction (Table 2, entry 19). The reaction of α-azidostyrene (Table 2, entries 20 and 21) was sluggish, requiring a higher temperature and longer reaction time.

The structures of the 3,4,5-trisubstituted 1*H*-pyrazoles were characterized by ¹H NMR, ¹³C NMR, and HRMS (ESI). In order to explore in more detail the structure of our product and the mechanism, we performed an X-ray study of **4n** (shown in Figure 1). More details related to the X-ray crystal structure can be found in the Supporting Information.

To gain information on the mechanistic aspects of the reaction, control reactions were examined (Scheme 2). The





Figure 1. X-ray crystal structure of **4n**.


reaction of benzaldehyde with 1 equiv of tosylhydrazine in DMF at room temperature gave the known hydrazone. Further reaction of hydrazone with 1 equiv of vinyl azide **1a** afforded the corresponding **4a** in 83% yield.

(14) Wu, L.; Shi, M. *J. Org. Chem.* **2010**, 75, 2296.

Scheme 2. Control Reaction



Scheme 3. Proposed Reaction Mechanism



On the basis of the above experimental results, the following possible mechanism for this cycloaddition reaction is proposed. As outlined in Scheme 3, the initial condensation of aromatic aldehyde **2** and tosylhydrazine **3** gave the corresponding hydrazone intermediate **I**, which subsequently generated diazomethane **II** upon heating in the presence of base.¹⁴ Intermediate **II** was then reacted with vinyl azide **1** via a [3 + 2] cycloaddition to afford the intermediate **V** with elimination of the azido group. Finally, the desired product **4** was obtained via the proton transfer of intermediate **VI**.

In conclusion, we have developed a simple and straightforward one-pot reaction from vinyl azide, aryl aldehyde, and tosylhydrazine to provide trisubstituted-1*H*-pyrazoles in the presence of base with good to excellent yields. The azido group serves as a leaving group in this approach.

Acknowledgment. This research was supported by the National Natural Science Foundation of China (Nos. 81072516 and 81273356), the Project of Science Technology Department of Zhejiang Province (2012C33065), the Project of Education Department of Zhejiang Province (Y201223193), Program for Zhejiang Leading Team of S&T Innovation, the Diabetes Research Group, USA.

Supporting Information Available. Experimental procedures, characterization data, and copies of ¹H and ¹³C NMR spectra for all products. This material is available free of charge via the Internet at <http://pubs.acs.org>.

The authors declare no competing financial interest.