This article was downloaded by: [McGill University Library] On: 27 September 2012, At: 06:06 Publisher: Taylor & Francis Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



# Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information: http://www.tandfonline.com/loi/gpss20

### AI(HSO<sub>4</sub>)<sub>3</sub> and AI<sub>2</sub>O<sub>3</sub>-SO<sub>3</sub>H as Efficient Catalysts for Modified Preparation of 3,4-Dihydropyrimidin-2 (1H)-ones/ thiones

Hamid Reza Shaterian<sup>a</sup>, Asghar Hosseinian<sup>a</sup>, Majid Ghashang<sup>a</sup>, Fahimeh Khorami<sup>a</sup> & Neda Karimpoor<sup>a</sup> <sup>a</sup> Department of Chemistry, Faculty of Sciences,

University of Sistan and Baluchestan, Zahedan, Iran

Version of record first published: 04 Sep 2009.

To cite this article: Hamid Reza Shaterian, Asghar Hosseinian, Majid Ghashang, Fahimeh Khorami & Neda Karimpoor (2009):  $AI(HSO_4)_3$  and  $AI_2O_3$ -SO<sub>3</sub>H as Efficient Catalysts for Modified Preparation of 3,4-Dihydropyrimidin-2 (1H)-ones/thiones, Phosphorus, Sulfur, and Silicon and the Related Elements, 184:9, 2333-2338

To link to this article: http://dx.doi.org/10.1080/10426500802454144

#### PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <u>http://www.tandfonline.com/page/terms-and-conditions</u>

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae, and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand, or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.



## AI(HSO<sub>4</sub>)<sub>3</sub> and AI<sub>2</sub>O<sub>3</sub>-SO<sub>3</sub>H as Efficient Catalysts for Modified Preparation of 3,4-Dihydropyrimidin-2 (1*H*)-ones/thiones

Hamid Reza Shaterian, Asghar Hosseinian, Majid Ghashang, Fahimeh Khorami, and Neda Karimpoor Department of Chemistry, Faculty of Sciences, University of Sistan and Baluchestan, Zahedan, Iran

An environmentally friendly procedure for the preparation of dihydropyrimidinone derivatives or their sulfur analogues under thermal solvent-free conditions in the presence of aluminium hydrogen sulfate  $[Al(HSO_4)_3]$  and alumina sulfuric acid  $(Al_2O_3 \cdot SO_3H)$  as heterogeneous catalysts was developed.

**Keywords** Aluminium hydrogen sulfate; alumina sulfuric acid; Biginelli reaction; 3,4-dihydropyrimidin-2(1*H*)-ones/thiones

#### INTRODUCTION

Dihydropyrimidinones (DHPMs) and their derivatives have attracted interest in medicinal chemistry because they exhibit a wide range of biological, pharmacological, and therapeutic properties.<sup>1</sup> DHPMs can serve as the integral backbones of several calcium channel blockers.<sup>2</sup> They are also reported to have antibacterial, antioxidant, antiviral, antitumor, anti-inflammatory,  $\alpha$ -1a-antagonist, and neuropeptide Y (NPY) antagonist activities.<sup>3–6</sup>

The classical Biginelli reaction of an aldehyde, 1,3-dicarbonyl, and urea or thiourea requires strongly acidic conditions with relatively low yields, high reaction times, and harsh conditions.<sup>7</sup> In order to improve the efficiency of Biginelli reaction, a number of catalysts have been developed. These include ammonium chloride,<sup>8</sup> ZrCl<sub>4</sub> or ZrOCl<sub>2</sub>,<sup>9</sup>

Address correspondence to Hamid Reza Shaterian, Department of Chemistry, Faculty of Sciences, University of Sistan and Baluchestan, PO Box 98135-674, Zahedan, Iran. E-mail: hrshaterian@hamoon.usb.ac.ir

Received 16 April 2008; accepted 12 August 2008.

We are thankful to the Sistan and Baluchestan University Research Council for partial support of this work.

vanadium(III) chloride,<sup>10</sup> chloroacetic acid,<sup>11</sup> graphite-supported lanthanum chloride,<sup>12</sup> alkylammonium and alkylimidazolium perhaloborates, phosphates, and aluminates,<sup>13</sup> 1,1,3,3-tetrametylguanidinium trifluoroacetate,<sup>14</sup> ferric chloride and boric acid,<sup>15</sup> sulphamic acid,<sup>16</sup> tetrabutyl ammonium hydrogensulfate,<sup>17</sup> and polyphosphate ester (PPE).<sup>18</sup> However, many of these methods suffer from the drawbacks inconsistent with green chemistry, such as the use of toxic chemicals and heavy metals. Therefore utilization of clean processes as well as eco-friendly and green catalysts has been a subject of intense research. The demand for environmentally benign procedures with heterogeneous catalysts prompted us to develop a safe alternate method for the synthesis of dihydropyrimidinones (DH-PMs) as biologically important compounds in the presence Al(HSO<sub>4</sub>)<sub>3</sub> and alumina sulfuric acid (Al<sub>2</sub>O<sub>3</sub>-SO<sub>3</sub>H) as heterogeneous catalysts (Scheme 1).



Catalyst = AI(HSO<sub>4</sub>)<sub>3</sub> and Al<sub>2</sub>O<sub>3</sub>-SO<sub>3</sub>H

**SCHEME 1** The synthesis of dihydropyrimidinones (DHPMs) in the presence of aluminum hydrogen sulfate and alumina sulfuric acid as catalysts.

In this research, Al(HSO<sub>4</sub>)<sub>3</sub> and alumina sulfuric acid (Al<sub>2</sub>O<sub>3</sub>-SO<sub>3</sub>H) as heterogeneous solid acid catalysts deserve special mention. These catalysts are safe, easy to handle, environmentally benign, and present fewer disposal problems. Al(HSO<sub>4</sub>)<sup>19</sup> and alumina sulfuric acid (0.2 g of alumina sulfuric acid equal to 0.6 mmol H<sup>+</sup>)<sup>20</sup> were prepared according to the reported procedures.

#### **RESULTS AND DISCUSSION**

In order to be able to carry out Biginelli condensation in a more efficient way and minimizing the time, temperature, and amount of the catalyst, the reaction of benzaldehyde, ethyl acetoacetate, and urea was selected as a model system to investigate the effects of the catalyst at different reaction temperatures (50, 60, 100, and 120°C) and different amounts of catalyst (5, 10, 15, 20, 25, and 30 mol%). The best result was obtained by carrying out the reaction with 1:1.2:1.5 molar ratios of aldehyde,

1,3-dicarbonyl compound, urea and 10 mol% of  $Al(HSO_4)_3$  at 100°C under thermal solvent-free conditions. We also used alumina sulfuric acid ( $Al_2O_3$ -SO<sub>3</sub>H) as catalyst in the model system to find optimization conditions at the same molar ratios. The best result was obtained by using 15 mol% of  $Al_2O_3$ -SO<sub>3</sub>H at 120°C.

Using these optimized reaction conditions, the scope and efficiency of these procedures were explored for the synthesis of a wide variety of substituted 3,4-dihydropyrimidin-2(1H)-ones or their sulfur analogues. The results are summarized in Table I.

As shown in Table I, aromatic aldehydes with both electronwithdrawing and electron-donating substituents reacted efficiently with urea/thiourea and methyl/ethyl acetoacetate or acetylacetone in the presence of a catalytic amount of  $Al(HSO_4)_3$  (10 mol %) or  $Al_2O_3$ - $SO_3H$  (15 mol%), forming the corresponding 3,4-dihydropyrimidin-2(1H)-ones/thiones without the formation of any side products, in good to high yields (Table I).

Entry	$\mathbb{R}^1$	$\mathbb{R}^2$	X	$\begin{array}{l} \mathrm{Al}(\mathrm{HSO}_4)_3\\ \mathrm{Time}\ (\mathrm{min}) \\ \mathrm{Yield}\ (\%)^a \end{array}$	$\begin{array}{l} Al_2O_3\text{-}SO_3H\\ Time (h)\!/\\ Yield (\%)^a \end{array}$	$mp^{\circ}C$ (lit $mp^{\circ}C$ )
1	$C_6H_5$	$C_2H_5O$	0	35/79	2.1/88	203-205 (200-202)8
2	$4-ClC_6H_4$	$C_2H_5O$	0	55/90	1.6/94	$210212\ (209211)^8$
3	$(CH_3)_2CH$	$C_2H_5O$	0	150/55	3.3/69	195-197 (196-197)11
4	$C_6H_5$	$C_2H_5O$	$\mathbf{S}$	30/80	2.7/81	$205 - 207 (205 - 206)^8$
5	$3-ClC_6H_4$	$C_2H_5O$	$\mathbf{S}$	110/83	1.5/81	$196 - 198 (192 - 196)^9$
6	$4-NO_2C_6H_4$	$C_2H_5O$	$\mathbf{S}$	90/79	2.5/83	108–110 (107–108) <sup>14</sup>
7	$3-OMeC_6H_4$	$C_2H_5O$	$\mathbf{S}$	100/76	1.7/73	$150-152 \ (150-152)^{12}$
8	$4-HOC_6H_4$	$C_2H_5O$	$\mathbf{S}$	50/60	2.1/75	194–196 $(193–195)^{11}$
9	$2,4$ - $Cl_2C_6H_3$	$CH_3O$	0	20/92	1.7/85	$252 - 254 \ (252 - 253)^{11}$
10	$3-NO_2C_6H_4$	$CH_3O$	0	15/74	2.7/74	$276-278(279-280)^8$
11	$C_6H_5$	$CH_3O$	$\mathbf{S}$	10/87	1.8/84	$221 - 223(221 - 222)^{11}$
12	$3-NO_2C_6H_4$	$CH_3O$	$\mathbf{S}$	30/86	2.7/68	$237 - 239 (237)^{16}$
13	$4-HOC_6H_4$	$CH_3O$	$\mathbf{S}$	10/77	1.1/81	$226 - 228 (227)^{16}$
14	$4-(Me)_2NC_6H_4$	$CH_3O$	$\mathbf{S}$	110/67	1.5/70	$152 - 154 (152 - 153)^{12}$
15	$C_6H_5$	$CH_3$	0	20/90	0.6/93	$234 – 236 \ (232 – 235)^8$
16	$C_6H_5$	$CH_3$	$\mathbf{S}$	15/91	1.1/93	$216 - 217 (214 - 215)^9$
17	$2\text{-ClC}_6\text{H}_4$	$CH_3$	$\mathbf{S}$	40/82	1.6/88	$173 – 174 \ (173 – 174)^9$
18	$3-ClC_6H_4$	$CH_3$	$\mathbf{S}$	15/80	1.1/87	$244 - 246 (243 - 245)^9$
19	$4-FC_6H_4$	$CH_3$	$\mathbf{S}$	25/92	0.7/93	$209-211 (209-212)^{13}$
20	$4\text{-}\mathrm{NO}_2\mathrm{C}_6\mathrm{H}_4$	$CH_3$	$\mathbf{S}$	25/74	1.6/81	$207 - 209 (207 - 209)^{13}$

TABLE I Synthesis of 3,4-Dihydropyrimidin- 2(1H)-ones/thiones

<sup>*a*</sup>Yields refer to the pure isolated products. All known products have been reported previously in the literature and were characterized by comparison of melting points, and IR and NMR spectra with authentic samples.<sup>8–18</sup>

Entry	Catalyst (mol %)	Conditions (molar ratio) <sup><math>a</math></sup>	Time/Yield (%)	
1	$Al(HSO_4)_3$ (10)	Solvent-free/100°C (1/1.2/1.5)	10–150 min/55–92	
2	$Al_2O_3-SO_3H(15)$	Solvent-free/120 °C (1/1.2/1.5)	0.6–3.3 h/68–94	
3	$NH_4Cl(40)$	Solvent-free/100°C (1/1/1.5)	3 h/42–92	
4	ZrOCl <sub>2</sub> .8H <sub>2</sub> O (15)	Solvent-free/90-100°C (1/1/1.3)	30–180 min/40–99	
5	$VCl_3$ (10)	CH <sub>3</sub> CN/Reflux (1/1/1.5)	2 h/65–96	
6	Chloroacetic acid (10)	Solvent-free/90 °C (1/1.1/1.5)	3 h/47–98	

TABLE II Comparison of the Results of  $Al(HSO_4)_3$  and  $Al_2O_3$ -SO<sub>3</sub>H with Other Catalysts Reported in the Literature

<sup>a</sup>The molar ratio of aldehyde/1,3-dicarbonyl compounds/urea or thiourea.

To show the merit of the present work in comparison with reported results in the literature, we compared results of aluminium hydrogen sulfate and alumina sulfuric acid with ammonium chloride,<sup>8</sup> ZrOCl<sub>2</sub>,<sup>9</sup> vanadium(III) chloride,<sup>10</sup> and chloroacetic acid<sup>11</sup> in the synthesis of Biginelli compounds. As shown in Table II, Al(HSO<sub>4</sub>)<sub>3</sub> and Al<sub>2</sub>O<sub>3</sub>-SO<sub>3</sub>H can act as effective catalysts with respect to reaction times, yields, and the obtained products.

In conclusion, we have developed a simple, cost-effective, and green procedure for the synthesis of 3,4-dihydropyrimidinones/thiones using aluminium hydrogen sulfate and alumina sulfuric acid as heterogeneous catalysts under thermal solvent-free conditions.

#### EXPERIMENTAL

All reagents were purchased from Merck and Aldrich and used without further purification. All yields refer to isolated products after purification. Products were characterized by comparison with authentic samples and by spectroscopy data (IR and <sup>1</sup>H NMR spectra). The NMR spectra were recorded on a Bruker Avance DPX 300 MHz instrument. Mass spectra were recorded on an Agilent Technologies 5973 network mass selective detector (MSD) operating at an ionization potential of 70 eV. Melting points were determined in open capillaries with a BUCHI 510 melting point apparatus. TLC was performed on silica-gel polygram SIL G/UV 254 plates.

### Typical Experimental Procedure for the One-Pot Preparation of 5-(Ethoxycarbonyl)-6-methyl-4-phenyl-3,4-dihydropyrimidin-2(1*H*)-one

A mixture of benzaldehyde (4 mmol), ethyl acetoacetate (4.8 mmol), urea (6 mmol), and  $Al(HSO_4)_3$  (0.4 mmol) at 100°C or  $Al_2O_3$ -SO<sub>3</sub>H

 $(0.2 \text{ g}, 0.6 \text{ mmol H}^+)$  at 120 °C was stirred in an oil bath for the appropriate time (Table I). After completion of the reaction as indicated by TLC, the resulting solidified mixture was diluted with ethyl acetate (5 mL), and the catalyst was separated by simple filtration and washed with ethyl acetate (2 × 5 mL). The filtrate obtained was washed with water (2 × 10 mL) and dried over anhydrous MgSO<sub>4</sub>. Evaporation of the solvent under reduced pressure yielded crude product, which was purified by recrystallization with ethanol to afford pure 5-(Ethoxycarbonyl)-6methyl-4-phenyl-3,4-dihydropyrimidin-2(1*H*)-one (Table I, Entry 1).

The desired pure products were characterized by comparison of their physical data with those of known 3,4-dihydropyrimidin-2(1H)-ones/thiones, which are reported in the litrature.<sup>8-18</sup>

#### REFERENCES

- [1] C. O. Kappe, Acc. Chem. Res., 33, 879 (2000).
- [2] K. S. Atwal, B. N. Swanson, S. E. Unger, D. M. Floyd, S. Moreland, A. Hedberg, and B. C. O'Reilly, *J. Med. Chem.*, **34**, 806 (1991).
- [3] C. O. Kappe, Eur. J. Med. Chem., 35, 1043 (2000).
- [4] S. J. Haggarty, T. U. Mayer, D. T. Miyamoto, R. Fathi, R. W. King, T. J. Mitchison, and S. L. Schreiber, *Chem. Biol.*, 7, 275 (2000).
- [5] K. S. Atwal, G. C. Rovnyak, B. C. O'Reilly, and J. Schewartz, J. Org. Chem., 54, 5898 (1989).
- [6] H. A. Stefani, C. B. Oliveira, R. B. Almeida, C. M. P. Pereira, R. C. Braga, R. Cella, V. C. Borges, L. Savegnago, and C. W. *Eur. J. Med. Chem.*, **41**, 513 (2006).
- [7] P. Biginelli, Gazz. Chim. Ital., 23, 360 (1893).
- [8] A. Shaabani, A. Bazgir, and F. Teimouri, Tetrahedron Lett., 44, 857 (2003).
- [9] J. C. Rodriguez-Dominguez, D. Bernardi, and G. Kirsch, *Tetrahedron Lett.*, 48, 5777 (2007).
- [10] G. Sabitha, G. S. K. K. Reddy, K. B. Reddy, and J. S. Yadav, *Tetrahedron Lett.*, 44, 6497 (2003).
- [11] Y. Yu, D. Liu, C. Liu, and G. Luo, Bioorg. Med. Chem. Lett., 17, 3508 (2007).
- [12] H. Khabazzadeh, K. Saidi, and H. Sheibani, *Bioorg. Med. Chem. Lett.*, 18, 278 (2008).
- [13] E. S. Putilova, G. V. Kryshtal, G. M. Zhdankina, N. A. Troitskii, and S. G. Zlotin, *Russ. J. Org. Chem.*, **41**, 512 (2005).
- [14] A. Shaabani and A. Rahmati, Catal. Lett., 100, 177 (2005).
- [15] I. S. Zorkun, S. Sarac, S. Celebib, and K. Erol, Bioorg. Med. Chem., 14, 8582 (2006).
- [16] S. A. Kotharkar, M. R. Jadhav, R. R. Nagawade, S. S. Bahekar, and D. B. Shinde, *Lett. Org. Chem.*, 2, 662 (2005).
- [17] A. Shaabani, A. Bazgir, and S. Arab-Ameri, *Phosphorus, Sulfur, and Silicon*, 179, 2169 (2004).
- [18] N. Foroughifar, A. Mobinikhaledi, and H. F. Jirandehi, *Phosphorus, Sulfur, and Silicon*, **178**, 1241 (2003).
- [19] For the preparation and some applications of Al(HSO<sub>4</sub>)<sub>3</sub> see: (a) H. R. Shaterian, M. Ghashang, and N. Mir, Arkivoc, xv, 1 (2007); (b) M. A. Zolfigol, A. G. Choghamarani, A. Taqian-Nasab, H. Keypour, and S. Salehzadeh, Bull. Kor. Chem. Soc., 24, 638

(2003); (c) F. Shirini, M. A. Zolfigol, and M. Abedini, *Bull. Chem. Soc. Jpn.*, **78**, 1982 (2005).

[20] For the preparation and some applications of Al<sub>2</sub>O<sub>3</sub>-SO<sub>3</sub>H see: (a) M. A. Zolfigol, *Tetrahedron*, **57**, 9509 (2001); (b) H. Sharghi, M. Hosseini-Sarvari, and R. Eskandari, J. Chem. Res., 488 (2005); (c) H. R. Shaterian, A. Hosseinian, H. Yarahmadi, and M. Ghashang, Lett. Org. Chem., **5**, 290 (2008).