


## Communications to the Editor

### Facile Synthesis of Substituted Brendanes via Spirocyclopropane Participation


Waldemar Adam,<sup>\*1</sup> Nêstor Carballeira,<sup>2</sup> Eva-Maria Peters,<sup>3</sup>  
Karl Peters,<sup>3</sup> and Hans Georg von Schnering<sup>3</sup>

*Institut für Organische Chemie, Universität Würzburg  
Am Hubland, D-8700 Würzburg, FRG  
Departamento de Química, Universidad de Puerto Rico  
Rio Piedras, Puerto Rico 00931  
and Max-Planck-Institut für Festkörperforschung  
D-7000 Stuttgart 80, FRG  
Received March 23, 1983*


Ring expansion (1 → 2) and cyclopropane migration (1 → 1a)



are possible structural transformations of  $\beta$ -cyclopropylethyl cations (1, eq 1). For the parent cation ring expansion has been documented as a significant pathway, but C-14 labeling rigorously excluded the cyclopropane migration route.<sup>4</sup> In the present communication we show that given the appropriate molecular skeleton, e.g., the norbornene moiety, spiroannulation as in 4




promotes ring expansion into the brendane derivatives 5 and 6 (eq 2), while endo fusion as in 8 involves cyclopropane migration (eq 3) during attack by *p*-toluenesulfonyl chloride.




Treatment of equimolar amounts of norbornene 4 and of *p*-toluenesulfonyl chloride in  $\text{CH}_2\text{Cl}_2$  at 20 °C afforded a mixture of the three products 5–7. These were separated by PTLC ( $\text{SiO}_2$ -petroleum ether), and analytical samples were obtained by means of Kugelrohr distillation.<sup>5</sup> The first fraction was the addition product 7 (ca. 10% yield), while the second and third fractions (15% and 26% yields, respectively), were identified as the brendane derivatives 6 and 5.

Although the  $^1\text{H}$  and  $^{13}\text{C}$  NMR data were compatible with these structure assignments, in view of the unusual ring-expanding rearrangement involved in this electrophilic addition, rigorous chemical proof was needed. Indeed, either the mixture or pure samples of 5 and 6 were converted in over 90% yield to the parent brendane 12 on treatment with excess sodium in liquid ammonia (eq 4). The spectral data of 12 matched perfectly those reported for authentic brendane.<sup>6</sup> Furthermore, *m*-chloroperbenzoic acid




(*m*-CPBA) oxidation of 5 afforded the sulfone 13 (eq 4) in 81% yield as crystalline solid, whose structure was confirmed by X-ray analysis.

Electrophilic attack by the arenesulfonyl chloride on the double bond of norbornene 4 induces ring expansion of the spirocyclopropane moiety leading to the brendane derivative 5. In view of the established three-center attack in the electrophilic addition of arenesulfonyl chlorides,<sup>7</sup> it is unlikely that free carbocationic intermediates as depicted in eq 1 intervene in this rearrangement. Instead, the initially formed cationic species is presumably the bridged episulfonium ion 14. Even then, several features are



remarkable about its subsequent ring expansion through spirocyclopropane participation. First of all, regioselective attack of the chloride ion at the 3-position in cation 14 gives the electrophilic addition product 7. Migration of the 1,6-bond produces the rearranged cation 16. Such neighboring group participation has recently been convincingly demonstrated in norbornenes.<sup>8</sup> From a synthetic standpoint the fact that the  $\beta$ -cyclopropylethyl cations 14 and 16 undergo ring expansion to give the respective brendyl cations 15 and 17 is most interesting and valuable.

As an extension of this unusual ring-expansion process it was hoped that the norbornene 8 with an endo-fused cyclopropane ring would provide synthetic access into the brexane skeleton.<sup>6</sup> However, treatment of equimolar amounts of norbornene 8 with *p*-toluenesulfonyl chloride in  $\text{CH}_2\text{Cl}_2$  at 20 °C gave a mixture of the three cyclopropane migrated products 9–11. Chromatographic separation (PTLC ( $\text{SiO}_2$ -petroleum ether)) and final purification by Kugelrohr distillation provided analytical samples.<sup>5</sup> Tricyclooctane 10 (8% yield) eluted first, followed by the tricycloheptane 9 as the major product (29% yield). The latter was oxidized in 90% yield to the sulfone 18, whose structure was rigorously established by X-ray analysis. The third fraction (5% yield) was identified as tricyclooctane 11.



The desired ring expansion of the episulfonium ion 19 into the brexyl cation 20 did not take place. Instead, the cyclopropane ring migrated to C-6 either via 3,4- or 2,4-bond breaking to afford the rearranged  $\beta$ -cyclopropylethyl cations 21 and 22, respectively, which are captured by chloride ion to give tricycloheptane 9 as major and tricyclooctane 10 as minor products. Of course, the free carbocations are intended here not as bona fide intermediates

(1) To whom all correspondence should be directed at the University of Würzburg.

(2) N.C. thanks the Thyssen Stiftung for a travel grant.

(3) Max-Planck-Institut für Festkörperforschung.

(4) Cartier, G. E.; Bunce, S. C. *J. Am. Chem. Soc.* **1963**, *85*, 932.

(5) Combustion analyses and spectral and X-ray data available on microfilm.

(6) Nickon, A.; Kwasnik, H. R.; Mathew, C. T.; Schwartz, T. D.; Williams, R. O.; DiGiorgio, J. B. *J. Org. Chem.* **1978**, *43*, 3904.

(7) Smit, W. A.; Zefirov, N. S.; Bodrikov, I. V.; Krimer, M. Z. *Acc. Chem. Res.* **1979**, *12*, 282.

(8) Carrupt, P.-A.; Vogel, P. *Tetrahedron Lett.* **1982**, *23*, 2563.

but rather as mechanistic constructs. The chemistry observed here for the episulfonium cation **19** matches perfectly that reported for the 3-nortricycylcarbanyl cation.<sup>9</sup> The tricyclooctane **11** is the regular skeletal rearrangement product expected in arenesulfenyl chloride addition to norbornenes.<sup>7,10</sup>

Our results display an interesting mechanistic dichotomy about cyclopropylethyl cationic fragments that are embodied in the norbornyl skeleton. Thus, spiroannulated cyclopropanes as in norbornene **4** (eq 2) prefer ring expansion on arenesulfenyl chloride addition, thereby providing a facile and efficient entry into structurally complex molecules such as brendanes. Including the preparation of the norbornene **4**, the parent brendane **12** was prepared in three steps in an overall yield of 15% starting from cheap, commercial compounds. On the other hand, endo-fused cyclopropanes as in norbornene **8** prefer cyclopropane migration on arenesulfenyl chloride addition.

**Acknowledgments** are made to the Fonds der Chemischen Industrie and Deutsche Forschungsgemeinschaft for financial support. We thank Dr. P. Binger (Mülheim) for a generous sample of methylenecyclopropane. We are grateful to Dr. D. Scheutzw for assisting us in the <sup>1</sup>H and <sup>13</sup>C NMR spectra.

**Registry No.** **4**, 6572-50-5; **5**, 86146-93-2; **6**, 86146-94-3; **7**, 86146-95-4; **8**, 3635-94-7; **9**, 86146-96-5; **10**, 86146-97-6; **11**, 86162-06-3; **12**, 1521-75-1; **13**, 86146-98-7; **18**, 86146-99-8; *p*-toluenesulfenyl chloride, 933-00-6.

**Supplementary Material Available:** Experimental, physical, and X-ray data and structures of **13** and **18** (16 pages). Ordering information is given on any current masthead page.

(9) (a) Sauers, R. R.; Beisler, J. A.; Feilich, H. *J. Org. Chem.* **1967**, *32*, 569. (b) Wiberg, K. B.; Wenzinger, G. R. *Ibid.* **1965**, *30*, 2278.

(10) Of mechanistic interest is to mention that the exo-fused norbornene **8** gave exclusively 6-endo-chloro-7-exo-thio-*p*-toluenesulfonylcyclo[3.2.1.0<sup>2,4</sup>]-octane in 80% yield as the regular trans-addition product with arenesulfenyl chloride.

### Proton-Transfer Spectroscopy of 3-Hydroxychromones. Extreme Sensitivity to Hydrogen-Bonding Perturbations<sup>†,‡</sup>

Dale McMorro and Michael Kasha\*


*Institute of Molecular Biophysics  
and Department of Chemistry  
Florida State University  
Tallahassee, Florida 32306  
Received April 7, 1983*

The discovery of excited-state proton transfer in *o*-hydroxychromones<sup>1</sup> (structure I, R = phenyl in 3-hydroxyflavone and R = methyl in 2-methyl-3-hydroxychromone) has generated considerable interest in laser kinetic<sup>2-4</sup> and piezospectroscopic<sup>5,6</sup> study of the mechanism of the phototautomerization. Our extension of the original work now indicates that the presence of stoichiometric and substoichiometric traces of water, or other H-bonding impurities, in the supposedly dry hydrocarbon solvents controls and competes with the excited-state proton-transfer process.

<sup>†</sup> This work was supported by Contract No. DE-AS-5-78EV05855 between the division of Biomedical and Environmental Research of the U.S. Department of Energy and Florida State University and by Contract No. INT 7808503 between the National Science Foundation and Florida State University.


<sup>‡</sup> This work was presented at the Gordon Conference on Molecular Electronic Spectroscopy, Wolfeboro, NH, Aug 1982.

- (1) Sengupta, P. K.; Kasha, M. *Chem. Phys. Lett.* **1979**, *68*, 382.
- (2) Woolfe, G. J.; Thistlethwaite, P. J. *J. Am. Chem. Soc.* **1981**, *103*, 6916.
- (3) Itoh, M.; Tokumura, K.; Tanimoto, Y.; Okada, Y.; Takeuchi, H.; Obi, K.; Tanaka, I. *J. Am. Chem. Soc.* **1982**, *104*, 4146.
- (4) Itoh, M.; Kurokawa, H. *Chem. Phys. Lett.* **1982**, *91*, 487.
- (5) Salaman, O. A.; Drickamer, H. G. *J. Chem. Phys.* **1981**, *75*, 572.
- (6) Salaman, O. A.; Drickamer, H. G. *J. Chem. Phys.* **1982**, *77*, 3329.



**Figure 1.** Fluorescence emission spectra of  $2.00 \times 10^{-5}$  M 3-hydroxyflavone in quick-frozen methylcyclohexane glass at 77 K as a function of addition of substoichiometric traces of water. Lowest curve shows unique tautomer fluorescence in the anhydrous solvent; highest curve, for a moderately wet solvent, shows tautomer fluorescence (I), perturbed-tautomer fluorescence (II), and "normal"-molecule fluorescence (III). The curves are normalized at 523 nm for clarity.

At temperatures greater than 200 K the roles of water solvates (structure II and III) are disguised, whereas at lower temperatures the properties of the solvated molecules dominate the excitation processes. In rigorously dry dilute hydrocarbon solutions, including both glass-forming and multicrystalline Shpol'skii matrices, we observe only the green tautomer fluorescence at all temperatures between 293 and 77 K.



The luminescence behavior at 77 K of 3-hydroxyflavone (2-phenyl-3-hydroxychromone) is reported in Figure 1. The solute concentration is  $2.00 \times 10^{-5}$  M in methylcyclohexane solvent. Because the effects that are to be described here are found to be strongly temperature and cooling-rate dependent, all spectra are reported for quick-frozen (i.e., sample tube plunged into liquid nitrogen) rigid-glass solutions at 77 K.

The lowest curve gives the luminescence spectrum of an extremely dry 3-hydroxyflavone solution.<sup>7</sup> The yellow-green fluorescence is completely analogous to the room-temperature tautomer emission previously reported<sup>1</sup> and thus is associated with the unsolvated proton-transferred species (structure I, tautomer). Upon the addition of substoichiometric traces of water to the anhydrous hydrocarbon solution, a fluorescence shoulder, desig-

(7) For drying procedures refer to the full paper, ref 10.