

Article

Subscriber access provided by Stanford University Libraries

An Efficient Synthesis of Functionalized Dihydro-1H-indol-4(5H)-ones via One-Pot Three-Component Reaction under Catalyst-Free Conditions

Hui-Yan Wang, and Da-Qing Shi

ACS Comb. Sci., Just Accepted Manuscript • DOI: 10.1021/co4000198 • Publication Date (Web): 18 Apr 2013 Downloaded from http://pubs.acs.org on April 29, 2013

Just Accepted

"Just Accepted" manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides "Just Accepted" as a free service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. "Just Accepted" manuscripts appear in full in PDF format accompanied by an HTML abstract. "Just Accepted" manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are accessible to all readers and citable by the Digital Object Identifier (DOI®). "Just Accepted" is an optional service offered to authors. Therefore, the "Just Accepted" Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the "Just Accepted" Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these "Just Accepted" manuscripts.



ACS Combinatorial Science is published by the American Chemical Society. 1155 Sixteenth Street N.W., Washington, DC 20036

Published by American Chemical Society. Copyright © American Chemical Society. However, no copyright claim is made to original U.S. Government works, or works produced by employees of any Commonwealth realm Crown government in the course of their duties.

An Efficient Synthesis of Functionalized Dihydro-1*H*-indol-4(5*H*)-ones via One-Pot Three-Component Reaction under Catalyst-Free Conditions

Hui-Yan Wang, Daqing Shi



An Efficient Synthesis of Functionalized Dihydro-1*H*-indol-4(5*H*)-ones via One-Pot Three-Component Reaction under Catalyst-Free Conditions

Hui-Yan Wang^{1,2}, Da-Qing Shi^{1,*}

¹Key Laboratory of Organic Synthesis of Jiangsu Province, College of Chemistry, Chemical Engineering and Materials Science, Soochow University, Suzhou 215123, P. R. China

²Department of Chemical Engineering, Huaihai Institute of Technology, Lianyungang 222005, P. R. China

ABSTRACT: A facile and efficient one-pot procedure for the preparation of functionalized dihydro-1*H*-indol-4(5*H*)-ones by a catalyst-free, three-component reaction of 1,3-dicarbonyl compounds, arylglyoxal monohydrate and enaminones under mild conditions in excellent yield is reported. This synthesis was confirmed to follow the group-assisted-purification (GAP) chemistry process, which can avoid traditional purifications, chromatography and recrystllization.

KEYWORDS: functionalized dihydro-1*H*-indol-4(5*H*)-ones, multi-component reactions, group-assisted-purification (GAP) chemistry process.

Introduction

Multi-component reactions (MCRs) are chemical transformations in which three or more different starting materials combine together via a one-pot procedure to give a final complex product. Such reactions have emerged as powerful and bond-forming efficient tools in organic, combinatorial and medicinal chemistry for their facileness and efficiency as well as their economy and ecology in organic synthesis.¹ In the past decade, there have been tremendous developments in three- and four-component reaction and significant efforts continue to be made to develop new MCRs.²

The indole nucleus is probably the most well-known heterocyclic compound, a common and important feature of a variety of natural products and medicinal agents.³ Compounds carrying the indole moiety exhibit antibacterial and antifungal activities.⁴

ACS Combinatorial Science

It is used as an important skeleton in organic synthesis⁵ and is also utilized in other important fields, such as medicinal chemistry.⁶ As a consequence, a number of methods have been reported for the construction of indoles.⁷ Recently, some functionalized indoles have been synthesized by using different starting materials.⁸

4-Hydroxycoumarins are a very important class of biologically active substances in nature and in medicine. They exhibit mainly anticoagulant activity and there are some drugs which are widely used as anticoagulants.⁹ They exhibit cytotoxic¹⁰ and antioxidant activities¹¹ as well as activity against HIV.¹² However, to the best of our knowledge, there have been few reports about the synthesis of coumarin-fused indole derivatives by multi-component reactions. Our interest in the synthesis of indole derivatives¹³ guided by the observation that the presences of two or more different heterocyclic moieties in a single molecule often enhances remarkably the biological activity, we herein described a facile synthesis of 3-(4-hydroxycoumarin-3-yl)dihydro-1*H*-indol-4(5*H*)-ones derivatives by a three-component reaction of 4-hydroxycoumarin, arylglyoxal monohydrates and enaminones in ethanol without any catalyst.

Results and discussion

We initially evaluated the three-component reaction of the 4-hydroxycoumarin $1\{I\}$, phenylglyoxal monohydrate $2\{I\}$, and enaminone $3\{I\}$ (Scheme 1). The reaction mixture, which was composed of a 1:1:1 mixture of $1\{1\}$, $2\{1\}$ and $3\{3\}$, was tested under a variety of different conditions. The effects of solvent and temperature were evaluated for this reaction, and the results are summarized in Table 1. It was found that when the reaction was carried out in water without any catalyst the yield of product was very low (Table 1, Entry 1). Chloroform and ethanol provided higher yields than those using other organic solvents (Table 1, Entry 5-6 vs Entries 2-4). Considering the volatility and toxicity of the chloroform, ethanol was chosen as the solvent for all further reactions. To identify the optimum reaction temperature, the reaction was carried out at r.t., 40°C, 60°C and reflux temperature, providing the product $4\{1,1,1\}$ in yields of 34%, 52%, 80% and 94% (Table 1, Entries 7-9 and 6), respectively. The use of additives, such as *L*-proline, resulted in no significant improvement of the yield (Table 1, Entry 10). Thus, the optimum conditions tested were refluxing ethanol without any catalyst.

Scheme 1. Model Reaction



Table 1. Optimizing the Reaction Conditions for the Synthesis of 4{1,1,1}

Entry	Solvent	Temperature (°C)	Time (h)	Yield ^a (%)
1	H ₂ O	80	1.0	34
2	DMF	80	1.0	69
3	Acetone	reflux	1.0	78
4	CH ₃ CN	reflux	1.0	80
5	CHCl ₃	reflux	1.0	92
6	EtOH	reflux	1.0	94
7	EtOH	r. t.	1.0	34
8	EtOH	40	1.0	52
9	EtOH	60	1.0	80
10	EtOH+ <i>L</i> -proline (10 mol %)	reflux	1.0	90

^a Yield was determined by HPLC-MS, if the precipitated solids were occurred during the reaction, acetonitrile should be added to dissolve the solids before determining the yield.

The optimized reaction conditions were then tested for library construction with eight 1,3-dicarbonyl compounds $1\{1-8\}$, five phenylglyoxal monohydrates $2\{1-5\}$, and sixteen enaminones $3\{1-16\}$ (Figure 1). The corresponding functionalized indole derivatives **4** were obtained in good yields in catalyst free, refluxing ethanol. The results are summarized in Table 2. This protocol was efficient with 1,3-dicarbonyl compounds with either 1,3-diketones or β -keto esters. It was also found that *n*-butyl and phenyl groups, bearing either electron-withdrawing or electron-donating groups on the enaminone ring, were tolerated under the reaction conditions, leading to the final products in satisfactory yields (up to 94%). Moreover, this synthesis followed the GAP chemistry (group-assisted-purification chemistry)¹⁴ process, which can avoid

ACS Combinatorial Science

traditional chromatography and recrystallization purifications. Pure products can be obtained by washing the solid, crude products with cold ethanol.

The structures of all products **4** were characterized by IR, ¹H NMR, ¹³C NMR, and HRMS analysis. The structure of compound $4\{1,1,1\}$ was further confirmed by X-ray diffraction analysis. The molecular structure of compound $4\{1,1,1\}$ is shown in Figure 2.

1,3-Dicarbonyl compounds 1:



Figure 1. Diversity of reagents.

1	
2	
3 ⊿	
5	
6	
7	
8	
9 10	
11	
12	
13	
14	
15	
17	
18	
19	
20	
22	
23	
24	
25	
20	
28	
29	
30	
32	
33	
34	
35	
30 37	
38	
39	
40	
41 42	
43	
44	
45	
46 47	
48	
49	
50	
51 52	
53	
54	
55	
56 57	
58	
59	
60	

Table 2.	Synthesis	of funct	ionalized	dihvdro	-1 <i>H</i> -indol	-4(5H)-one	derivative	s 4
	•			•		•	,		

OH t	- OOH Ar OH +	R^1 NH R^2	Ar ethanol OH reflux	R^2 R^1 R^1
1 { <i>1</i> -8}	2 {1-5}	3 {1-16}	4 { 1 { <i>1</i> -8}, 2 {	1-5}, 3 {1-16}}
Entry	Products	Time (h)	Isolated Yield (%)	M.P. (°C)
1	4 { <i>l</i> , <i>l</i> , <i>l</i> }	1.0	89	162-163
2	4 { <i>1</i> , <i>1</i> , <i>2</i> }	1.0	91	308-310
3	4 { <i>1</i> , <i>1</i> , 3 }	2.0	88	170-172
4	4 { <i>1</i> , <i>1</i> , <i>4</i> }	1.5	90	176-178
5	4 { <i>1</i> , <i>1</i> , 5 }	4.0	77	185-186
6	4 { <i>1</i> , <i>1</i> , <i>6</i> }	0.7	90	270-271
7	4 { <i>1</i> , <i>1</i> , <i>7</i> }	1.0	94	152-154
8	4 { <i>1</i> , <i>1</i> , <i>8</i> }	1.0	87	294-295
9	4 { <i>1,1,9</i> }	1.0	88	268-269
10	4 { <i>1</i> , <i>1</i> , <i>10</i> }	1.5	82	278-280
11	4 { <i>1</i> , <i>1</i> , <i>1</i> 2}	1.0	81	114-116
12	4 { <i>1</i> , <i>1</i> , <i>1</i> 3}	1.0	87	282-284
13	4 { <i>1</i> , <i>1</i> , <i>1</i> 4}	1.5	90	172-174
14	4 { <i>1</i> , <i>1</i> , <i>1</i> 5}	1.0	94	278-279
15	4 { <i>1,1,16</i> }	5.0	72	167-170
16	4 { <i>1,2,3</i> }	1.5	83	250-251
17	4 { <i>1,2,11</i> }	1.0	90	172-174
18	4 { <i>1</i> , <i>3</i> , <i>1</i> }	1.5	85	264-265
19	4 { <i>1</i> , 3 , 3 }	1.5	92	172-173
20	4 { <i>1</i> , <i>4</i> , <i>1</i> }	1.5	89	> 300
21	4 { <i>1</i> , <i>4</i> , <i>3</i> }	1.5	94	> 300
22	4 { <i>1</i> ,5,3}	1.5	92	> 300
23	4 { <i>1,5,11</i> }	1.0	93	> 300

Page 7 of 16

_	24	4 {2,1,1}	1.0	81	268-270
	25	4 { <i>2</i> , <i>1</i> , <i>3</i> }	3.0	88	282-283
	26	4{2,1,6}	0.7	85	142-144
	27	4 { <i>2</i> , <i>1</i> , <i>12</i> }	1.0	80	279-281
	28	4 { <i>3</i> , <i>1</i> , <i>1</i> }	1.0	80	196-198
	29	4{3,1,6}	0.7	82	183-184
	30	4 { <i>4</i> , <i>1</i> , <i>1</i> }	1.0	75	174-175
	31	4 { <i>5</i> , <i>1</i> , <i>1</i> }	1.0	77	137-138
	32	4{5,1,3}	1.5	88	154-156
	33	4 { <i>6</i> , <i>1</i> , <i>1</i> }	1.0	84	293-295
	34	4 { <i>6</i> , <i>1</i> , <i>3</i> }	1.5	89	290-292
	35	4 {7,1,1}	1.0	76	142-144
	36	4 { <i>8</i> , <i>1</i> , <i>1</i> }	1.5	82	239-240
	37	4{8,1,3}	1.0	85	224-226



Figure 2. Crystal structure of compound 4{1,1,1}

Although the detailed mechanism of this reaction remains to be fully clarified, the formation of compound **4** could be explained by the reaction sequence in Scheme 2. First, a Knoevenagel condensation of 1,3-dicarbonyl compounds **1** with phenylglyoxal **2** is proposed to give intermediate **A**. Michael addition of enaminone **3** to intermediate **A** then occurs to provide intermediate **B**, which undergoes intramolecular cyclization and dehydration to form the desired product **4**.





Conclusion

The new GAP synthesis of functionalized dihydro-1*H*-indol-4(5*H*)-one derivatives has been achieved by three-component reaction of 1,3-dicarbonyl compounds, arylglyoxal monohydrates and enaminones under mild conditions without any catalyst. Good chemical yields have been achieved without the use of the traditional purifications, chromatography and recrystallization.

Experimental Section

Melting points are uncorrected. IR spectra were recorded on Varian F-1000 spectrometer in KBr with absorptions in cm⁻¹. ¹H NMR and ¹³C NMR were determined on Varian Invoa-400 MHz or Invoa-300 MHz spectrometer in DMSO- d_6 solution. *J* values are in Hz. Chemical shifts are expressed in ppm downfield from internal standard TMS. HRMS analyses were carried out using Bruker micrOTOF-Q instrument or TOF-MS instrument.

General Procedure for the Synthesis of 4. A dry 25 mL flask was charged with 1,3-dicarbonyl compounds 1 (1 mmol), phenylglyoxal monohydrate 2 (1 mmol),

ACS Combinatorial Science

enaminones **3** (1 mmol) and ethanol (5 mL). The mixture was stirred at refluxing temperature for 0.7-5 h. After completion of the reaction (confirmed by TLC), the reaction mixture was cooled to room temperature. The crystalline solids were collected and washed with little cold ethanol to give the pure products **4**.

Compound 4{*1,1,1*}: White powder; IR (KBr, v, cm⁻¹): 3430, 3280, 3058, 2956, 2869, 1729, 1694, 1665, 1624, 1515, 1492, 1451, 1408, 1366, 1270, 1194, 1103, 1044, 1020, 961, 897, 837, 757, 699; ¹H NMR (DMSO-*d*₆, 400 MHz) δ (ppm): 10.81 (s, 1H, OH), 7.77 (d, J = 8.0 Hz, 1H, ArH), 7.54 (t, J = 7.6 Hz, 1H, ArH), 7.31-7.24 (m, 2H, ArH), 7.19 (d, J = 7.6 Hz, 2H, ArH), 7.09-6.99 (m, 7H, ArH), 2.56 (d, J = 16.4 Hz, 1H, CH₂), 2.47-2.45 (m, 1H, CH₂), 2.30-2.26 (m, 4H, CH₂+CH₃), 2.16 (d, J = 15.6 Hz, 1H, CH₂), 1.04 (s, 3H, CH₃), 1.01 (s, 3H, CH₃); ¹³C NMR (DMSO-*d*₆, 75 MHz) δ (ppm): 192.80, 162.17, 161.17, 152.78, 144.37, 138.06, 135.37, 134.90, 132.36, 131.41, 130.19, 130.02, 129.77, 128.26, 127.96, 127.65, 124.25, 123.86, 118.27, 116.45, 109.36, 100.34, 52.62, 36.85, 35.38, 28.91, 28.38, 21.07; HRMS calcd for C₃₂H₂₇NNaO₄ [M+Na]⁺: 512.1838, found: 512.1842.

Compound 4{2,1,1}: Light red powder; IR (KBr, v, cm⁻¹): 3427, 3063, 3035, 2938, 2867, 2735, 2691, 1662, 1637, 1576, 1515, 1492, 1447, 1404, 1367, 1279, 1233, 1209, 1107, 1125, 1081, 1050, 1000, 934, 923, 828, 804, 766, 726, 699, 650; ¹H NMR (DMSO-*d*₆, 400 MHz) δ (ppm): 10.64 (s, 1H, OH), 7.17-6.97 (m, 9H, ArH), 5.91 (s, 1H, =CH), 2.50-2.45 (m, 2H, CH₂), 2.25-2.18 (m, 5H, CH₂+CH₃), 2.10 (s, 3H, CH₃), 1.00 (s, 6H, 2×CH₃); ¹³C NMR (DMSO-*d*₆, 75 MHz) δ (ppm): 192.46, 166.34, 164.14, 160.89, 143.64, 137.88, 134.96, 134.27, 131.88, 130.18, 129.67, 128.19, 127.91, 127.33, 118.14, 110.74, 100.51, 97.07, 52.75, 36.89, 35.19, 28.67, 28.59, 21.05, 19.75; HRMS calcd forC₂₉H₂₇NO₄ [M]⁺: 453.1940, found: 453.1929.

Compound 4{3,1,6}: White powder ; IR (KBr, v, cm⁻¹): 3588, 3425, 3056, 3030, 2957, 2892, 2644, 1670, 1585, 1562, 1513, 1486, 1451, 1367, 1299, 1257, 1211, 1166, 1148, 1046, 1028, 1013, 921, 845, 807, 775, 718, 702; ¹H NMR (DMSO- d_6 , 400 MHz) δ (ppm): 9.63 (s, 1H, OH), 7.08-7.03 (m, 5H, ArH), 6.92-6.89 (m, 4H, ArH), 3.71 (s, 3H, OCH₃), 2.46-1.96 (m, 8H, 4×CH₂), 1.09-0.87 (m, 12H, 4×CH₃); ¹³C NMR (DMSO- d_6 , 75 MHz) δ (ppm): 196.51, 192.26, 170.40, 158.83, 143.57, 133.94,

132.17, 130.50, 129.71, 129.32, 127.92, 126.99, 118.32, 114.69, 112.19, 109.66, 56.49, 55.72, 52.88, 36.93, 35.17, 32.06, 28.77, 28.29, 19.02; HRMS calcd forC₃₁H₃₃NO₄[M]⁺: 483.2410, found: 483.2398.

Compound 4{*4,1,1*}: White powder ; IR (KBr, *v*, cm⁻¹): 3431, 3059, 3033, 2952, 2875, 2620, 1664, 1578, 1515, 1491, 1448, 1422, 1381, 1334, 1280, 1209, 1183, 1160, 1117, 1071, 1049, 985, 923, 833, 811, 769, 730, 697, 649; ¹H NMR (DMSO-*d*₆, 400 MHz) δ (ppm): 9.76 (s, 1H, OH), 7.15-6.90 (m, 9H, ArH), 2.47-1.70 (m, 13H, CH₃+5×CH₂), 0.98 (s, 6H, 2×CH₃); ¹³C NMR (DMSO-*d*₆, 75 MHz) δ (ppm): 192.36, 143.35, 137.63, 135.27, 133.55, 132.12, 130.10, 129.61, 129.53, 128.07, 127.95, 127.89, 126.92, 118.40, 112.67, 110.82, 52.88, 36.98, 36.95, 35.16, 28.67, 21.04, 20.91, 19.01; HRMS calcd forC₂₉H₂₈NO₃ [M-H]⁺: 438.2069, found: 438.2073.

Compound 4{*5*,*1*,*3*}: White powder; IR (KBr, *v*, cm⁻¹): 3432, 3089, 3028, 2955, 2892, 1049, 1620, 1578, 1448, 1417, 1383, 1363, 1244, 1116, 1091, 1069, 1050, 1014, 923, 888, 843, 760, 724, 649; ¹H NMR (DMSO-*d*₆, 400 MHz) δ (ppm): 10.04 (s, 1H, OH), 7.04-6.93 (m, 14H, ArH), 3.35-3.10 (m, 1H, CH), 2.61-2.19 (m, 8H, 4×CH₂), 0.99 (s, 6H, 2×CH₃); ¹³C NMR (DMSO-*d*₆, 75 MHz) δ (ppm): 192.57, 192.44, 144.09, 143.95, 143.47, 143.42, 136.71, 133.81, 133.56, 132.83, 131.86, 131.74, 130.00, 129.70, 129.65, 129.02, 128.96, 128.22, 128.19, 127.33, 127.15, 118.59, 112.84, 112.79, 110.51, 110.42, 52.90, 52.84, 38.87, 38.53, 36.86, 35.22; 31.14, 28.67; HRMS calcd forC₃₄H₃₀ClNO₃ [M]⁺: 535.1914, found: 535.1907.

Compound 4{*6*,*1*,*1*}: White powder; IR (KBr, *v*, cm⁻¹): 3427, 3031, 2955, 2932, 2869, 2590, 1665, 1576, 1515, 1487, 1410, 1313, 1208, 1116, 1045, 1018, 924, 840, 802, 771, 721, 699, 665, 649; ¹H NMR (DMSO-*d*₆, 400 MHz) δ (ppm): 11.36 (s, 1H, OH), 7.15-6.95 (m, 9H, ArH), 2.47-1.93 (m, 11H, CH₃+4×CH₂), 0.97 (s, 6H, 2×CH₃); ¹³C NMR (DMSO-*d*₆, 75 MHz) δ (ppm): 192.42, 143.61, 137.88, 135.07, 134.15, 133.95, 132.03, 131.87, 130.21, 129.73, 128.08, 127.92, 127.18, 118.32, 112.52, 110.16, 52.88, 36.95, 36.87, 35.12, 28.66, 28.58, 21.06; HRMS calcd forC₂₈H₂₆NO₃ [M-H]⁺: 424.1913, found: 424.1940.

Compound 4{*7*,*1*,*1*}: Grey powder; IR (KBr, *v*, cm⁻¹): 3427, 3001, 2958, 2869, 1755, 1663, 1578, 1515, 1490, 1423, 1281, 1117, 1045, 1014, 924, 831, 768, 728, 698, 649;

ACS Combinatorial Science

¹H NMR (DMSO-*d*₆, 400 MHz) δ (ppm): 11.82 (s, 1H, OH), 7.14-7.03 (m, 9H, ArH), 4.57-4.00 (m, 2H, CH₂), 2.49-2.23 (m, 7H, CH₃+2×CH₂), 0.98 (s, 6H, 2×CH₃); ¹³C NMR (DMSO-*d*₆, 75 MHz) δ (ppm): 192.58, 174.58, 173.84, 143.84, 138.07, 134.88, 134.81, 131.43, 130.25, 129.89, 128.25, 127.92, 127.52, 118.34, 107.80, 95.79, 67.08, 52.73, 36.86, 35.17, 28.63, 28.41, 21.05; HRMS calcd forC₂₇H₂₄NO₄ [M-H]⁺: 426.1705, found: 426.1714.

Compound 4{*8*,*1*,*1*}: Yellow powder; IR (KBr, *v*, cm⁻¹): 3433, 3001, 2955, 2869, 1751, 1651, 1516, 1490, 1465, 1445, 1417, 1367, 1321, 1246, 1206, 1154, 1114, 1040, 1014, 924, 886, 843, 778, 764, 746, 701, 649; ¹H NMR (DMSO-*d*₆, 400 MHz) δ (ppm): 7.88 (s, 4H, ArH), 7.20-7.15 (m, 7H, ArH), 7.07 (d, *J* = 7.6 Hz, 2H, ArH), 4.45 (s, 1H, CH), 2.47 (s, 2H, CH₂), 2.25 (s, 3H, CH₃), 2.05 (s, 2H, CH₂), 0.90 (s, 6H, 2×CH₃); ¹³C NMR (DMSO-*d*₆, 75 MHz) δ (ppm): 198.95, 193.07, 144.01, 142.27, 138.21, 137.54, 135.54, 134.56, 130.50, 130.27, 129.98, 128.83, 128.26, 127.87, 122.98, 116.47, 110.78, 53.86, 51.35, 36.46, 35.53, 28.47, 21.04; HRMS calcd forC₃₂H₂₆NO₃ [M-H]⁺: 472.1913, found: 472.1923.

Acknowledgement. This work was partially supported by the Major Basic Research Project of the Natural Science Foundation of the Jiangsu Higher Education Institutions (No. 10KJA150049), A Project Funded by the Priority Academic Project Development of Jiangsu Higher Education Institutions and the Foundation of Key Laboratory of Organic Synthesis of Jiangsu Province (No. JSK0812) and the Jiangsu College Graduate Research and Innovation Project of the Jiangsu Province Department of Education (CXZZ11 0090).

Supporting Information Available. Experimental details and spectroscopic characterization for compounds **4**, and the X-ray crystallographic information for compound $4\{1,1,1\}$. This information is available free of charge via the Internet at http://pubs.acs.org/.

References and Notes

(1) (a) Thompson, B. B.; Montgomery, J. Enone-Alkyne Reductive Coupling: A versatile entry to substituted pyrroles. *Org. Lett.* **2011**, *13*, 3289-3291. (b) Trost, B.

M.; Frontier, A. J. Atom economical syntheses of oxygen heterocycles via tandem palladium-catalyzed reactions. J. Am. Chem. Soc. 2000, 122, 11727-11728. (c) Trost,
B. M.; Gutierrez, A. C.; Livingston, R. C. Tandem ruthenium-catalyzed redox isomerization-O-conjugate addition: An atom-economic synthesis of cyclic ethers. Org. Lett. 2009, 11, 2539-2542.

(2) (a) Nair, V.; Vinod, A. U.; Rajesh, C. A novel synthesis of 2-aminopyrroles using a three-component reaction. J. Org. Chem. 2001, 66, 4427-4429. (b) Bertozzi, F.; Gustafsson, M.; Olsson, R. A novel metal iodide promoted three-component synthesis of substituted pyrrolidines. Org. Lett. 2002, 4, 3147-3150. (c) Bora, U.; Saikia, A.; Boruah, R. C. A novel microwave-mediated one-pot synthesis of indolizines via a three-component reaction. Org. Lett. 2003, 5, 435-438. (d) Umkehrer, M.; Kalinski, C.; Kolb, J.; Burdack, C. A new and versatile one-pot synthesis of indol-2-ones by a novel Ugi-four-component-Heck reaction. Tetrahedron Lett. 2006, 47, 2391-2393. (e) Jiang, B.; Li, Y.; Tu, M. S.; Wang, S. L.; Tu, S. J.; Li, G. G. Allylic amination and N-arylation-based domino reactions providing rapid three-component strategies to fused pyrroles with different substituted patterns. J. Org. Chem. 2012, 77, 7497-7505. (f) Thirumurugan, P.; Perumal, P. T. A simple one-pot synthesis of functionalised 6-(indol-3-yl)-2,2'-bipyridine derivatives via multi-component reaction under neat condition. Tetrahedron Lett. 2009, 50, 4145-4150. (g) Ganguly, N. C.; Roy, S.; Mondal, P.; Saha, R. An efficient one-pot organocatalytic synthesis of 9-(1H-indol-3-yl)-xanthen-4-(9H)-ones under mild aqueous micellar conditions. Tetrahedron Lett. 2012, 53, 7067-7071. (h) Jiang, B.; Tu, S. J.; Kaur, P.; Wever, W.; Li, G. Four-component domino reaction leading to multifunctionalized quinazolines. J. Am. Chem. Soc. 2009, 131, 11660-11661. (i) Jiang, B.; Yi, M. S.; Shi, F.; Tu, S. J.; Pindi, S.; McDowell, P.; Li, G. A multi-component domino reaction for the direct access to polyfunctionalized indoles via intermolecular allylic esterification and indolation. Chem. Commun. 2012, 48, 808-810. (j) Sun, J.; Sun, Y.; Xie, Y. J.; Yan, C. G. Facile synthesis of dispirooxindole-fused heterocycles via domino 1,4-dipolar addition and Diels-Alder reaction of in situ generated Huisgen 1,4-dipoles. Org. Lett. 2012, 14, 5172-5175. (k) Li, M.; Cao, H.; Wang, Y.; Lv, X. L.; Wen, L. R. One-pot

ACS Combinatorial Science

multicomponent cascade reaction of *N*,*S*-ketene acetal: Solvent-free synthesis of imidazo[1,2-*a*]thiochromeno[3,2-*e*]pyridines. *Org. Lett.* **2012**, *14*, 3470-3473. (l) Wen, L. R.; Shi, Y. J.; Liu, G. Y.; Li, M. Modulating the reactivity of functionalized *N*,*S*-ketene acetal in MCR: Selective synthesis of tetrahydropyridines and thiochromeno[2,3-*b*]pyridines via DABCO-catalyzed tandem annulation. *J. Org. Chem.* **2012**, *77*, 4252-4260.

(3) Houlihan, W. J.; Remers, W. A.; Brown, R. K. *Indoles: Part I*; Wiley: New York, 1992.

(4) Sundberg, R. J. The Chemistry of Indoles; Academic: New York, 1996.

(5) (a) Fuchs, J. R.; Funk, R. L. Indol-2-one Intermediates: ☐ Mechanistic Evidence and Synthetic Utility. Total syntheses of (±)-flustramines A and C. *Org. Lett.* **2005**, *7*, 677-680. (b) Chen, W. L.; Cai, Y. F.; Fu, X.; Liu, X. H.; Lin, L. L.; Feng, X. M. Enantioselective one-pot synthesis of 2-amino-4-(indol-3-yl)-4*H*-chromenes. *Org. Lett.* **2011**, *13*, 4910-4913. (c) Qiu, Y. A.; Kong, W. Q.; Fu, C. L.; Ma, S. M. Carbazoles via AuCl₃-catalyzed cyclization of 1-(indol-2-yl)-3-alkyn-1-ols. *Org. Lett.* **2012**, *14*, 6198-6201. (d) Thirumurugan, P.; Perumal, P. T. InCl₃ mediated one-pot synthesis of indol-3-yl pyridine and 2,2'-bipyridine derivatives through multi-component reaction. *Tetrahedron* **2009**, *65*, 7620-7629.

(6) (a) Wu, S.; Fluxe, A.; Janusz, J. M.; Sheffer, J. B.; Browning, G.; Blass, B.; Cobum, K.; Hedges, R.; Murawsky, M.; Fang, B.; Fadayel, G. M.; Hare, M.; Djandjighian, L. Discovery and synthesis of tetrahydroindolone derived semicarbazones as selective Kv1.5 blockers. *Bioorg. Med. Chem. Lett.* 2006, *16*, 5859-5863. (b) Barta, T. E.; Barabasz, A. F.; Foley, B. E.; Geng, L.; Hall, S. E.; Hanson, G. J.; Jenks, M.; Ma, W.; Rice, J. W.; Veal, J. Novel carbazole and acyl-indole antimitotics. *Bioorg. Med. Chem. Lett.* 2009, *19*, 3078-3080. (c) Pita, B.; Masaguer, C. F.; Raviña, E. New synthetic approaches to CNS drugs. A straightforward, efficient synthesis of tetrahydroindol-4-ones and tetrahydroquinolin-5-ones via palladium-catalyzed oxidation of hydroxyenaminones. *Tetrahedron Lett.* 2002, *43*, 7929-7932. (d) Barraja, P.; Diana, P.; Montalbano, A.; Carbone, A.; Cirrincione, G.; Viola, Giampietro.; Salvador, A.; Vedaldi, D.; Dall'Acqua, F.

Thiopyrano[2,3-*e*]indol-2-ones: Angelicin heteroanalogues with potentphotoantiproliferative activity. *Bioorg. Med. Chem.* **2008**, *16*, 9668-9683. (e) Olson, G. L.; Cheung, H. C.; Morgan, K. D.; Blount, J. F.; Todaro, L.; Berger, Leo. A dopamine receptor model and its application in the design of a new class of rigid pyrrolo[2,3-*g*]isoquinoline antipsychotics. *J. Med. Chem.* **1981**, *24*, 1026-1034. (f) Chiang, C. C.; Lin, Y. H.; Lin, S. F.; Lai, C. L.; Liu, C. W.; Wei, W. Y.; Yang, S. C.; Wang, R. W.; Teng, L. W.; Chuang, S. H.; Chang, J. M.; Yuan, T. T.; Lee, Y. S.; Chen, P.; Chi, W. K.; Yang, J. Y.; Huang, H. J.; Liao, C. B.; Huang, J. J. Discovery of pyrrole-indoline-2-ones as aurora kinase inhibitors with a different inhibition profile. *J. Med. Chem.* **2010**, (16), 5929-5941.

(7) (a) England, D. B.; Merey, G.; Padwa, A. Substitution and cyclization reactions involving the quasi-antiaromatic 2*H*-indol-2-one ring system. *Org. Lett.* **2007**, *9*, 3805-3807. (b) Ban, X.; Pan, Y.; Lin, Y. F.; Wang, S. Q.; Du, Y. F.; Zhao, K. Synthesis of carbazolones and 3-acetylindoles via oxidative C-N bond formation through PIFA-mediated annulation of 2-aryl enaminones. *Org. Biomol. Chem.* **2012**, *10*, 3606-3609. (c) Kuethe, J. T.; Wong, A.; Davies, I. W. Rapid and efficient synthesis of 1*H*-indol-2-yl-1*H*-quinolin-2-ones. *Org. Lett.* **2003**, *5*, 3975-3978. (d) Caron, S.; Vazquez, E. Efficient synthesis of [6-chloro-2-(4-chlorobenzoyl)-1*H*-indol-3-yl]acetic acid, a novel COX-2 inhibitor. *J. Org. Chem.* **2003**, *68*, 4104-4107.

(8) (a) Chen, Z. Y.; Zhang, D. Q.; Wu, J. A facile route to polysubstituted indoles via three-component reaction of 2-ethynylaniline, sulfonyl azide, and nitroolefin, *Org. Lett.* **2011**, *13*, 848-851. (b) Jiang, B.; Li, Y.; Tu, M. S.; Wang, S. L.; Tu, S. J.; Li, G. Allylic amination and *N*-arylation-based domino reactions providing rapid three-components strategies to fused pyrroles with different substituted patterns, *J. Org. Chem.* **2012**, *77*, 7497-7505. (c) Jiang, B.; Yi, M. S.; Shi, F.; Tu, S. J.; Pindi, S.; McDowell, P.; Li, G. A multi-component domino reaction for the direct access to polyfunctionalized indoles via intermolecular allylic esterification and indolation, *Chem. Commun.* **2012**, *48*, 808-810. (d) Fu, L. P.; Shi, Q. Q.; Shi, Y.; Jiang, B.; Tu, S. J. Three-component domino reactions for regioselective formation of bis-indole derivatives, *ACS Comb. Sci.* **2013**, *15*, 135-140.

ACS Combinatorial Science

(9) (a) Abdelhafez, O. M.; Amin, K. M.; Batran, R. Z.; Maher, T. J.; Nada, S. A.; Sethumadhavan, S. Synthesis, anticoagulant and PIVKA-II induced by new 4-hydroxycoumarin derivatives. *Bioorg. Med. Chem.* **2010**, *18*, 3371-3378. (b) Gumńska, M.; Eckstein, M. The anticoagulant action of some derivatives of 3,3'-(arylidene)-bis-4-hydroxycoumarin. *J. Med. Pharm. Chem.* **1961**, *3*, 583-595.

(10) (a) Stanchev, S.; Momekov, G.; Jensen, F.; Manolov, Ilia. Synthesis, computational study and cytotoxic activity of new 4-hydroxycoumarin derivatives. *Eur. J. Med. Chem.* **2008**, *43*, 694-706. (b) Kostova, I.; Trendafilova, N.; Momekov, G. Theoretical and spectroscopic evidence for coordination ability of 3,3'-benzylidenedi-4-hydroxycoumarin, New neodymium (III) complex and its cytotoxic effect. *J. Inorg. Biochem.* **2005**, *99*, 477-487.

(11) (a) Pérez-Cruz, F.; Serra, S.; Delogu, G.; Lapier, M.; Maya, J. D.; Olea-Azar, C.; Santana, L.; Uriarte, E. Antitrypanosomal and antioxidant properties of 4-hydroxycoumarins derivatives. *Bioorg. Med. Chem. Lett.* **2012**, *22*, 5569-5573. (b) Rodrígueza, S. A.; Nazareno, M. A.; Baumgartner, M. T. Effect of different C3-aryl substituents on the antioxidant activity of 4-hydroxycoumarin derivatives. *Bioorg. Med. Chem.* **2011**, *19*, 6233-6238. (c) Stanchev, S.; Hadjimitova, V.; Traykov, T.; Boyanov, T.; Manolova, I. Investigation of the antioxidant properties of some new 4-hydroxycoumarin derivatives. *Eur. J. Med. Chem.* **2009**, *44*, 3077-3082.

(12) (a) Thaisrivongs, S.; Watenpaugh, K. D.; Howe, W. J.; Tomich, P. K.; Dolak, L. A.; Chong, K. T.; Tomich, C. S. C.; Tomasselli, A. G.; Turner, S. R.; Strohbach, J. W.; Mulichak, A. M.; Janakiraman, M. N.; Moon, J. B.; Lynn, J. C.; Horng, M. M.; Hinshaw, R. R.; Curry, K. A.; Rothrock, D. J. Structure-based design of novel HIV protease inhibitors: carboxamide-containing 4-hydroxycoumarins and 4-hydroxy-2-pyrones as potent nonpeptidic inhibitors. *J. Med. Chem.* 1995, *38*, 3624-3637. (b) Thaisrivongs, S.; Tomich, P. K.; Watenpaugh, K. D.; Chong, K. T.; Howe, W. J.; Yang, C. P.; Strohbach, J. W.; Turner, S. R.; McGrath, J. P.; Bohanon, M. J.; Lynn, J. C.; Mulichak, A. M.; Spinelli, P. A.; Hinshaw, R. R.; Pagano, P. J.; Moon, J. B.; Ruwart, M. J.; Wilkinson, K. F.; Rush, B. D.; Zipp, G. L.; Dalga, R. J.; Schwinde, F. J.; Howard, G. M.; Padbury, G. E.; Toth, L. N.; Zhao, Z.; Koeplinger, K. A.; Kakuk,

T. J.; Cole, S. L.; Zaya, R. M.; Piper, R. C.; Jeffrey, P. Structure-based design of HIV protease inhibitors: 4-hydroxycoumarins and 4-hydroxy-2-pyrones as non-peptidic inhibitors. *J. Med. Chem.* **1994**, *37*, 3200-3204.

(13) (a) Liu, H; Dou, G. L.; Shi, D. Q. Region- and stereoselective synthesis of novel dispiropyrrolidine bisoxindole derivatives via multicomponent reactions. J. Comb. Chem. 2010, 12, 292-294. (b) Liu, H.; Dou, G. L.; Shi, D. Q. Regioselective synthesis of novel spiropyrrolidines and spirothiapyrrolizidines through multi-component 1,3-dipolar cycloaddition reaction of azomethine ylides. J. Comb. Chem. 2010, 12, 633-637. (c) Li, Y. L.; Chen, H.; Shi, C. L.; Shi, D. Q.; Ji, S. J. Efficient one-pot synthesis of spirooxindole derivatives catalyzed by L-proline in aqueous medium. J. Comb. Chem. 2010, 12, 231-237. (d) Chen, H.; Shi, D. Q. Efficient one-pot synthesis of novel spirooxindole derivatives via three-component reaction in aqueous medium. J. Comb. Chem. 2010, 12, 571-576.

(14) (a) Kaur, P.; Pindi, S.; Wever, W.; Rajale, T.; Li, G. Asymmetric catalytic Strecker reaction of *N*-phosphonyl imines with Et₂AlCN using amino alcohols and BINOLs as catalysts. *Chem. Commun.* **2010**, *46*, 4330-4332. (b) Kaur, P.; Pindi, S.; Wever, W.; Rajale, T.; Li, G. Asymmetric catalytic *N*-phosphonyl imine chemistry: The use of primary free amino acids and Et₂AlCN for asymmetric Strecker reaction. *J. Org. Chem.* **2010**, *75*, 5144-5150. (c) Kaur, P.; Wever, W.; Pindi, S.; Milles, R.; Gu, P.; Shi, M.; Li, G. The GAP chemistry for chiral N-phosphonyl imine-based Strecker reaction. *Green Chem.* **2011**, *13*, 1288-1292. (d) Zou, Y.; Hu, Y.; Liu, H.; Shi, D. Q. Rapid and efficient ultrasound-assisted method for the combinatorial synthesis of spiro[indoline-3,4'-pyrano[2,3-c]pyrazole] derivatives. *ACS Comb. Sci.* **2012**, *14*, 38-43.