absence of cyclopropenecarbaldehyde (18) in the direct irradiation of the diene 13. Additionally this experiment shows that cyclopropenecarbaldehyde (18) is not the precursor to furan (19) produced in the photolysis of 13, although this is a well-documented photorearrangement of substituted acylcyclopropenes.<sup>21</sup> As no furan is produced upon irradiation of the diene 13 in the presence of isobenzofuran, either Dewar furan (4) or some intermediate species involved in its thermal rearrangement to cvclopropenecarbaldehyde (18) must be photoreactive and act as the precursor to furan. On the basis of these results we can summarize the photolysis of the diene 13 as shown in Scheme IV, in which Dewar furan (4) plays a key role. Experiments on elucidating the nature of such intermediate(s) are under active investigation and will be reported in due course.

Registry No. 4, 74496-19-8; 6, 3005-27-4; 7, 12078-17-0; 8, 98652-86-9; 9, 98652-87-0; 10, 98652-88-1; 11, 26307-17-5; 12, 98652-89-2; 13, 98652-90-5; 14, 95641-37-5; 15, 270-75-7; 16, 98652-91-6; 17, 98717-52-3; 18, 36998-21-7; 20, 98652-92-7; 21, 98717-53-4; 23, 2957-95-1; 24, 98652-93-8; 25, 98717-54-5; 26, 98717-55-6; 27, 98652-94-9; 28, 98717-56-7; 29, 84636-45-3.

(21) Inter alia: Padwa, A.; Akiba, M.; Chou, C. S.; Cohen, L. J. Org. Chem. 1982, 47, 183-191.

## Phenyliodonium Bis(phenylsulfonyl)methylide: A New Hypervalent Iodonium Ylide

Lazaros Hadjiarapoglou, Spyros Spyroudis, and Anastasios Varvoglis\*

> Laboratory of Organic Chemistry University of Thessaloniki GR 54006 Thessaloniki, Greece Received July 15, 1985

Compounds with a sulfonyl group in general<sup>1</sup> and gem-disulfones in particular<sup>2</sup> are increasingly attracting attention for their potential in organic synthesis. At the same time, iodonium vlides are being established as a class of ylides of synthetic and mechanistic interest.<sup>3</sup> We report the synthesis and chemistry of a new type of iodonium ylide, the carbanion moiety of which is a  $\beta$ -disulfonyl group, i.e., phenyliodonium bis(phenylsulfonyl)methylide (3). This compound, which could not be formed from the disulfone 2 and iodosylbenzene,<sup>4</sup> has now been prepared from diacetoxyiodobenzene (1) and 2 in 90% yield by the general method of Schank and Lick.5

$$\frac{\text{PhI}(\text{OAc})_2 + \text{CH}_2(\text{SO}_2\text{Ph})_2 \xrightarrow{\text{KOH, MeOH}} \text{PhI}^+ \text{C}(\text{SO}_2\text{Ph})_2}{2} \xrightarrow{\text{KOH, MeOH}} \text{PhI}^+ \text{C}(\text{SO}_2\text{Ph})_2$$

It is mentioned that the synthesis of the related ylides  $RC_6H_4I^+-C(SO_2F)_2$  has been reported,<sup>6</sup> without any reactions.

(5) Schank, K.; Lick, C. Synthesis 1983, 392.





<sup>a</sup> Reactants, conditions, and products (with % yields). (i) cyclohexene, MeCN-hv, 4 h, 4 (31%); (ii) PhH, hv, 6 h, 5 (65%); (iii)  $\label{eq:constraint} \begin{array}{l} Z = \text{pyridine, reflux, } Cu(acac)_2, 4 \text{ h}, \textbf{ 6a} (75\%); \ Z = \text{Ph}_3\text{P}, \text{CHCl}_3, \\ \text{reflux, } Cu(acac)_2, 3 \text{ h}, \textbf{ 6b} (92\%); \ Z = (\text{Me}_2\text{N})_2\text{CS}, \text{CHCl}_3, \text{reflux}, \end{array}$  $Cu(acac)_2$ , 7 h, 6c (78%); Z = Me<sub>2</sub>S,  $h\nu$ , 2.5 h, 6d (84%); Z = PhSMe, reflux, Cu(acac), 1 h, 6e (29%).

Scheme II



The stability of 3 is limited, and it decomposes on attempted recrystallization. However, it can be stored for at least 2 weeks at -10 °C without change. Its spectral characteristics are in agreement with the proposed structure.<sup>7</sup> Ylide 3 enters into several types of reactions, three of which are shown in Scheme I and involve transfer of the bis(phenylsulfonyl)methylene moiety to several nucleophiles to give gem-disulfones which are otherwise not easily accessible. Thus, cycloaddition occurs with cyclohexene to yield the bicyclic disulfone 4; C-H bond insertion is observed with benzene to give the phenylated disulfone 5, and transylidation occurs with various S, N, and P nucleophiles which are converted into their ylides 6. With few exceptions yields are good. These reactions have been carried out under two sets of conditions, either photolytically<sup>8</sup> or thermally in the presence of catalytic amounts of Cu(acac)<sub>2</sub>, and optimal conditions are shown in Scheme I. It is noted that ylides 6b,<sup>9</sup> 6c,<sup>10</sup> and 6d<sup>11</sup> have been prepared previously in low yields.

When 3 was heated in carbon disulfide under reflux precipitation of the tetrathianic tetrasulfone 7 occurred in 30% yield. The formation of 7 probably proceeds as suggested for an analogous reaction of diphenyldiazomethane with CS2.12 Heating 7 in diglyme with Cu powder leads to partial desulfurization yielding the dithietanic tetrasulfone 8 (Scheme II).

The dihalobis(phenylsulfonyl)methanes 9a-c have been obtained from 3 and N-halosuccinimides without any added catalyst. The

- (11) Dieckman, J. J. Org. Chem. 1965, 30, 2272
- (12) Schonberg, A.; Frese, E.; Brosowski, K. H. Chem. Ber. 1962, 95,

<sup>(1)</sup> Magnus, P. D. Tetrahedron 1977, 33, 2019. Durst, T. In 'Comprehensive Organic Chemistry"; Pergamon: Oxford, 1979; Vol. 3, Chapter 11.8, pp 171-213.

<sup>(2)</sup> De Lucchi, O.; Pasquato, L.; Modena, G. Tetrahedron Lett. 1984, 25, 3643, 3647. Griffiths, G.; Hughes, S.; Stirling, C. J. M. J. Chem. Soc., Chem. Commun. 1982, 236, 237. Hendrickson, J. B; Boudreaux, G. J.; Palumbo, D. S. Tetrahedron Lett. 1984, 25, 4617. Trost, B. M.; Cossy, J.; Burks, J. J. Am. Chem. Soc. 1983, 105, 1052. Lazukina, L. A.; Mushalo, I. L.; Neplyuev, V. M.; Kukhar, V. P. Zh. Org. Khim. 1983, 19, 2417. Rao, Y. K.; Nagarajan, M. Synthesis 1984, 757. Messinger, P.; Kusuma, K. Synthesis 1980, 565.
 Usalam, B. L. Samara, M. B. J. Chem. Par. Sunon, 1985, 44 Harlow, R. L.; Sammes, M. P. J. Chem. Res., Synop. 1985, 44.

<sup>(3)</sup> Koser, G. F. In "The Chemistry of Functional Groups, Supplement D", Wiley: New York, 1983; Chapter 18, pp 774-806. Varvoglis A. Synthesis 1984, 709. Moriarty R. M.; Prakash, I.; Prakash, O.; Freeman, W. A. J. Am. Chem. Soc. 1984, 106, 6082. Moriarty, R. M.; Bailey, B. R.; Prakash, O.; Prakash, I. J. Am. Chem. Soc. 1985, 107, 1375. Mansuy, D.; Battioni, J. P.; Akhrem, I.; Dupré, D.; Fischer, J.; Weiss, R. Morgenstern-Badaran, I. J. Am. Chem. Soc. 1984, 106, 6112.
 (4) Hayasi, Y.; Okada, T.; Kawanisi, M. Bull. Chem. Soc. Jpn. 1970, 43,

<sup>2506</sup> 

<sup>(6)</sup> Maletina, I. I.; Mironova, A. A.; Savina, T. I. Yagupolskii, Y. L. Zh. Org. Khim. 1979, 15, 2416.

<sup>7)</sup> All new compounds have elemental analysis and spectral data (IR, <sup>1</sup>H

<sup>(1)</sup> All five composite with the assigned structures. (8) A Philips 400-W low-pressure mercury lamp has been used. Irradiation was performed in Pyrex tubes under N<sub>2</sub> ( $\lambda_{max}$  (3) 277 nm, in MeOH). (9) Horner, L.; Oedinger, H. Chem. Ber. 1958, 91, 437. (10) Lloyd, D.; Millar, R. W. Tetrahedron 1980, 36, 2675.

<sup>3077</sup> 



formation of 9c is noteworthy since, unlike 9a and 9b, 9c cannot be prepared by direct iodination of  $2.^{13}$ 

 $CH_2Cl_2$ 3 + N-X succinimide room temperature  $(PhSO_2)_2CX_2$ 9a, X = Cl (54%) 9b, X = Br (81%) 9c, X = I (79%)

Compounds 9b and 9c were also formed from 3 and the corresponding elemental halogen, while 9a was obtained from 3 with  $SO_2Cl_2$ . The ylide 3 has oxidizing properties: thus 3 reacts with aniline to give trans-azobenzene, with anthracene to give 9,10anthraquinone, and with diphenylacetylene to give benzil, all in low yields. Furthermore, phenylalkenes such as styrene, transstilbene, and ethyl cinnamate undergo cleavage of the double bond with 3 at room temperature to afford benzaldehyde as the main product. Acrylonitrile was polymerized by 3 exothermically.

When 3 was heated in such diverse H-containing solvents as AcOH, CH<sub>2</sub>Cl<sub>2</sub>, MeCN, etc., the formation of iodobenzene and the disulfone 2 was observed. However, when 3 was heated under reflux in t-BuOH in the presence of  $Cu(acac)_2$  and under N<sub>2</sub>, CO<sub>2</sub> was evolved, and the unexpected product phenyl benzenethiosulfonate (14) was obtained, in 80% yield. The same ester was formed in varying amounts in most of the reactions of 3. We propose the mechanism shown in Scheme III for the formation of 14. Apparently the solvent plays a passive role since  $CO_2$ evolution has been detected during the thermolysis of solid 3. Support for this mechanism is provided by an attempted synthesis of ylide 16 from (phenylsulfonyl)(phenylthio)methane (15) and 1, where the decomposition product of the expected ylide diphenyl disulfide (17) was obtained. The formation of 17 probably proceeds analogously to the pathway in Scheme III.

PhSO<sub>2</sub>CH<sub>2</sub>SPh 
$$\xrightarrow{M_{eOH, KOH, -10 \circ C}}$$
  
15  $[PhI^{+-}C(SO_2Ph)(SPh)] \xrightarrow{-PhI} PhSSPh + CO_2$   
16  $17$ 

The reactions of 3 with nucleophiles as well as its thermolysis probably involve dissociation of the ylide into iodobenzene and bis(phenylsulfonyl)carbene (10).<sup>14</sup> Carbene formation from iodonium ylides has previously been suggested in two cases.<sup>4,15</sup> The carbene 10 is apparently in equilibrium with 11, as oxirenes are with ketocarbenes.<sup>16</sup> Rearrangement of **11** into **12** followed by intramolecular nucleophilic collapse of 12 to 13 and decarboxylation of 13 would give 14. This stage  $(13 \rightarrow 14)$  is possibly responsible for the production of free radicals and the polymerization of acrylonitrile.

There is precedent<sup>17</sup> for O-sulfonyl attack at highly electrophilic C. The enhanced nucleophilic character of the sulfonyl oxygen

Scheme IV



of 3 or 10 must be responsible for the oxidations effected by 3 under mild conditions. Both 3 and 10 are probably stabilized by resonance so that considerable negative charge may be acquired by the sulfonyl oxygens. However, carbene formation here and in reactions of 3 with electrophiles seems unlikely. The formation of transient iodonium salts is more justifiable. We note, for example, that phenyliodonium dinitromethylide gives isolable iodonium salts<sup>18</sup> with FSO<sub>3</sub>H.

Iodobenzene is a byproduct in all the above reactions of 3. A different type of reactivity was observed when 3 was allowed to react with dimedone (18) in non-hydroxylic solvents. In this case, "reversed" transylidation occurred, and the phenyliodonio dimedonate (19) was obtained in 50% yield (Scheme IV). This reaction may involve proton transfer from 18 ( $pK_a = 5$ ) to 3 (estimated<sup>19</sup> p $K_a$  of protonated 3 is ~4) and subsequent attack of the dimedonate ion on protonated 3.

Registry No. 1, 3240-34-4; 2, 3406-02-8; 3, 98858-34-5; 4, 98858-35-6; 5, 38564-68-0; 6 (Z = pyridine), 98858-36-7; 6 (Z = Ph<sub>3</sub>P) (P(V) entry), 96415-47-3; 6 (Z = Ph<sub>3</sub>P) (ylide entry), 25809-68-1; 6 (Z =  $(Me_2N)_2CS)$ , 77134-48-6; 6 (Z = Me\_2S), 2292-72-0; 6 (Z = PhSMe), 53799-65-8; 7, 98858-37-8; 8, 98858-38-9; 9a, 603-35-0; 9b, 2782-91-4; 9c, 75-18-3; 14, 1212-08-4; 15, 15296-86-3; 17, 882-33-7; CS<sub>2</sub>, 75-15-0; N-chlorosuccinimide, 128-09-6; N-bromosuccinimide, 128-08-5; N-iodosuccinimide, 516-12-1; aniline, 62-53-3; azobenzene, 103-33-3; 9,10anthraquinone, 84-65-1; diphenylacetylene, 501-65-5; benzyl, 2154-56-5; styrene, 100-42-5; trans-stilbene, 103-30-0; ethyl cinnamate, 103-36-6; benzaldehyde, 100-52-7; acrylonitrile, 107-13-1; polyacrylonitrile, 25014-41-9; anthracene, 120-12-7.

(18) Semenov, V. V.; Shevelev, S. A.; Fainzilberg, A. A. Izv. Akad. Nauk. SSSR, Ser. Khim. 1978, 2348. (19) Neiland, O.; Karele, B. Zh. Org. Khim. 1971, 7, 1611.

## **Oxidative Nucleophilic Addition of Organovanadium Reagents to Aldehydes with Formation of Ketones**

Toshikazu Hirao,\* Daisuke Misu, and Toshio Agawa

Department of Applied Chemistry Faculty of Engineering, Osaka University 2-1 Yamadaoka, Suita, Osaka 565, Japan Received June 17, 1985

A variety of organometallic reagents have been developed for selective molecular elaboration.<sup>1,2</sup> In particular, alcohol formation via carbonyl addition reactions constitutes one of the important C-C bond construction methods. Herein we describe a new methodology for carbonyl alkylation which involves an organovanadium compound as a key reagent for oxidative nucleophilic addition.

The organovanadium reagents employed here were generated in situ in dichloromethane from equimolar amounts of vanadium trichloride and organolithium or magnesium compounds. The reactions of the reagents thus obtained with aldehydes resulted in oxidative C-C bond formation leading to the corresponding ketones (Scheme I). For example, vanadium trichloride was

<sup>(13)</sup> Jarvis, B.; Fried, H. E. J. Org. Chem. 1975, 40, 1278

<sup>(14)</sup> The formation of 10 has been proposed in reactions of the unusually stable bis(phenylsulfonyl)diazomethane,<sup>11</sup> but no reaction occurred with benzene and alkenes, whereas triphenylphosphine did reaction occurred with benzene and alkenes, whereas triphenylphosphine did react to give the phos-phazene. Therefore formation of 10 from this precursor is unlikely. (15) Hood, J. N. C.; Lloyd, D.; MacDonald, W. A.; Shepherd, T. M. Tetrahedron 1982, 38, 3355.

<sup>(16)</sup> Csizmadia, I. G.; Font, J.; Strauss, O. P. J. Am. Chem. Soc. 1968, 90, 7360.

<sup>(17)</sup> Chalkey, G. R.; Snodin, D. J.; Stevens, G.; Whiting, M. C. J. Chem Soc. C 1970, 682. Braverman, S.; Reisman, D. Tetrahedron Lett. 1977, 1753.

<sup>(1)</sup> For example: Schwartz, J.; Labinger, J. A. Angew. Chem., Int. Ed. Engl. 1976, 15, 333. Mukaiyama, T. Ibid. 1977, 16, 817. Reetz, M. T. Topics *Curr. Chem.* **1982**, *106*, 1. Imamoto, T.; Tawarayama, Y.; Kusumoto, T.; Yokoyama, M. *J. Synth. Org. Chem. Jpn.* **1984**, *42*, 143. (2) Seebach, D. In "Modern Synthetic Methods"; Scheffold, R., Ed.; Wiley: New York, 1983; Vol. 3, p 217.