# Cycloadditions of 3,4-Dihydro-2*H*-pyrrole *N*-Oxide with Thioketones and a Selenoketone

## Kosei Shioji,<sup>\*1</sup> Akio Matsumoto,<sup>1</sup> Masahiko Takao,<sup>1</sup> Yoshimitsu Kurauchi,<sup>1</sup> Toshiyuki Shigetomi,<sup>1</sup> Yoshinobu Yokomori,<sup>2</sup> and Kentaro Okuma<sup>1</sup>

<sup>1</sup>Department of Chemistry, Faculty of Science, Fukuoka University, Jonan-ku, Fukuoka 814-0180

<sup>2</sup>Department of Chemistry, National Defense Academy, Hashirimizu, Yokosuka 239-8686

Received July 18, 2006; E-mail: shioji@fukuoka-u.ac.jp

Cycloadditions of 2-diphenylphosphinoyl-2-methyl-3,4-dihydro-2*H*-pyrrole *N*-oxide (DPhPMPO), 3,4-dihydro-2*H*pyrrole *N*-oxide having a diphenylphosphinoyl group at the C2 position with thioketones afforded the corresponding 1,4,2-oxathiazolidines. Dissociation constants of these 5-membered ring products were determined. The cycloadducts were stabilized by the diphenylphosphinoyl group. The reaction of DPhPMPO with di-*tert*-butyl selenoketone gave the corresponding selenoamide under microwave irradiation. The formation of the selenoamide indicated that the cycloaddition of DPhPMPO with the selenoketone analogue also proceeded through the formation of the corresponding 5-membered ring product.

The pathways of 1,3-dipolar cycloadditions of nitrones to carbon-carbon multiple bonds, such as alkenes, or alkynes to form isoxazolidines or 2,3-dihydroisoxazoles are involved in many synthetic strategies.<sup>1</sup> Studies on the chemical properties of N-oxide as a 1.3-dipole are of substantial interest in this regard. It is also well known that compounds having heteroatom-carbon multiple bonds react with electron-rich 1,3-dipoles.<sup>2</sup> Black and Watson have described 1,3-cycloaddition reactions of nitrones including 2,2-dimethyl-3,4-dihydro-2Hpyrrole N-oxide (DMPO) (1a) with thicketones, such as adamantane-2-thione (2a) and 2,2,4,4-tetramethyl-3-thioxocyclobutanone (2b), to afford 1,4,2-oxathiazolidine derivatives.<sup>3</sup> Huisgen, Sustmann, et al. have reported kinetic studies on these cycloadditions and evaluated the high reactivity of the C=S double bond toward nitrones by using MO calculations.<sup>4</sup> Mloston, Heimgartner, and co-worker have reported the cycloaddition of azole N-oxide with several thiocarbonyl compounds.<sup>5</sup> However, to our knowledge, there are no other reports on the 1,3-dipolar cycloaddition reactions of thioketones with nitrones due to the instability and inaccessibility of nitrones. Furthermore, to our knowledge, cycloaddition of 3,4-dihydro-2H-pyrrole N-oxide with a C=Se bond has not been reported. The difference in reactivity between thioketones and selenoketones toward nitrones is also of interest. We have recently investigated the synthesis of 3,4-dihydro-2H-pyrrole N-oxide containing a phosphinoyl group as an electron-withdrawing group at the 2-position.<sup>6</sup> In this paper, we report that the cycloaddition reactions of 3,4-dihydro-2H-pyrrole N-oxide, 2-diphenylphosphinoyl-2-methyl-3,4-dihydro-2H-pyrrole N-oxide (DPhPMPO) (1b), with several dipolarophiles, such as thicketone and selenoketone, behave in a variety of ways.

### **Results and Discussion**

**Synthesis of DPhPMPO (1b).** Nitrone **1b** was synthesized by the following three-step reactions described in Scheme 1. Diphenylphosphine oxide (**3**) was prepared from commercially



available chlorodiphenylphosphine (**4**) in 97% yield. 2-(Diphenylphosphinoyl)-2-methyl-1-pyrrolidine (**5**) was obtained by the cyclization of 5-chloro-2-pentanone (**6**) and ammonia with **3** (63% yield). Oxidation of **5** was carried out by using Oxone<sup>TM</sup> in acetone to give the desired nitrone **1b** in 47% yield without purification by column chromatography. Purifications of the overall synthesis were carried out by recrystallization from water containing acetonitrile. The obtained crystals could be stored at room temperature.

The structure of **1b** was confirmed by <sup>1</sup>H, <sup>13</sup>C, <sup>31</sup>PNMR, elemental analysis, and X-ray crystallographic analysis. The X-ray data are reported in Table 1. A single crystal had one stereoisomer, and the crystal lattice had two water molecules. No interaction was observed between the two oxygen atoms on *N*-oxide and the phosphinoyl moiety. The P–C(1) bond is moderately longer than the other P–C bonds. It appered that the position of the diphenylphosphinoyl group caused steric hindrance at the nitrone face in the crystalline state (Fig. 1).

**Thermal Reaction of DPhPMPO (1b) with Thioketones.** We reacted DPhPMPO with adamantane-2-thione (2a). The reaction proceeded quickly in CHCl<sub>3</sub> at room temperature to give the corresponding oxathiazolidine **7a** in 90% yield. Although the relative configuration was not elucidated, the adduct was obtained as a single diastereomer. The reaction of **1b** with thioxocyclobutanone **2b** gave cycloadduct **7b** as colorless crystals in 60% yield (Scheme 2). These structures were identified by <sup>1</sup>H, <sup>13</sup>C NMR and elemental analysis. These results were similar to those obtained for the reaction of **1a**. On the other hand, no reaction occurred between **1b** and bornane-6-thione (**2c**) or 1,1,3,3-tetramethylindane-2-thione (**2d**)

Table 1. Selected Bond Lengths (Å) and Angles (°) of 1b

P(1)-O(1)	1.481(4)	O(2)–N(1)	1.298(6)	C(2)–C(3)	1.507(9)
P(1)–C(6)	1.800(6)	N(1)-C(4)	1.298(6)	C(3)–C(4)	1.480(10)
P(1)–C(12)	1.802(6)	N(1)-C(1)	1.513(7)		
P(1)-C(1)	1.850(6)	C(1)-C(2)	1.521(8)		
N(1)-C(1)-1	P(1)	112.7(4)	C(4)-N(	(1)-C(1)	111.0(5)
C(2)-C(1)-I	P(1)	109.7(4)	C(3) - C(3)	(2)-C(1)	106.2(6)
C(5)-C(1)-I	P(1)	109.4(4)			



Fig. 1. ORTEP of **1b**.

at room temperature. It has been reported that the steric encumbrance of these thicketones 2a-2d increases in the sequence of 2a < 2b < 2c < 2d.<sup>7</sup> The poor reactivity of 2c and 2d was caused by steric hindrance of thiccarbonyl compounds.

The reported cycloadducts of nitrones with thioketones dissociate to a large extent in dilute solution. Huisgen et al. have reported that the dissociation constant of the oxathiazolidine 9 generated from linear N-(diphenylmethylene)methylamine *N*-oxide (1c) with 2b in CDCl<sub>3</sub> was  $1.2 \times 10^{-3}$  mol L<sup>-1</sup> when the initial concentrations of the nitrone and 2b were 0.94 and 0.37 mol L<sup>-1</sup>, respectively.<sup>4</sup> The dissociation constants of adducts 7a and 7b at 25 °C in CDCl<sub>3</sub> were determined by <sup>1</sup>H NMR. The dissociation constants of cycloadducts 8 derived from 1a and 2 were also determined. These results are summarized in Table 2. Cycloadducts 7a and 8a had smaller dissociation constants than those of cycloadducts 7b and 8b formed from 2b. This may be because the steric congestion around O,S-acetal moiety destabilizes the latter cycloadducts. Although the initial concentrations of 1b and 2b for the reaction of 1b and 2b were lower than those of the linear nitrone 1c and 2b for the corresponding reaction, the dissociation constant of 7b was three times smaller than that of the adduct of 1c with **2b**. Compared with these values, our results indicated that the oxathiazolidines 7 generated from 1b and 2a or 2b are more stable than 9 generated from the reaction of the linear nitrone

Table 2. Dissociation Constants of 1,3-Cycloadducts 7 and 8

<b>1</b> <sup>a)</sup>	<b>2</b> <sup>b)</sup>	Adduct	$10^3 K_{\rm diss}/{\rm mol}{\rm L}^{-1{\rm c})}$
а	a	<b>8</b> a	$6.96 \pm 0.39$
а	b	8b	$19.5 \pm 0.39$
b	a	7a	$0.114 \pm 0.025$
b	b	7b	$0.4107 \pm 0.041$
c	b	9	$1.2^{d}$

a) Initial concentration of  $[1] = 0.049 \text{ mol } L^{-1}$ . b) Initial concentration of  $[2] = 0.049 \text{ mol } L^{-1}$ . c) Determined by <sup>1</sup>H NMR at 25 °C. d) Ref. 4.



Scheme 2.



Scheme 3.

1c with 2b. The dissociation constant of 8b was higher than that of **7b**, because adamantane-2-thione (**2a**) is less sterically crowded than thioxocyclobutanone 2b. These observations suggested that the diphenylphosphinoyl moiety contributes stability to the adduct 7a and 7b. The bulky diphenylphosphinoyl moiety did not influence cycloaddition of 1b with 2a and 2b.

The cycloadditions of 1b with 2c or 2d did not proceed at room temperature. When a microwave reaction between 1b with 2c was carried out in chloroform, the pale orange solution originating from the thicketone 2c gradually became colorless. Using TLC analysis, a product ( $R_f = 0.65$ , EtOAc:MeOH = 9:1) was detected along with 1b ( $R_f = 0.1$ ) and 2c ( $R_f =$ 0.95). From the reaction of 1b with 2d, a trace amount of a product ( $R_f = 0.45$ ) was also detected. However, these products were not observed by <sup>1</sup>HNMR of the reaction mixture.

Reaction of DPhPMPO with Di-tert-butyl Selenoketone. Since selenoketones are also useful 1,3-dipolarophiles, several selenoketones were employed instead of 2.8 The reaction of 1b with di-tert-butyl selenoketone (10) did not proceed at room temperature. However, under microwave irradiation, the cycloaddition of 1b with selenoketone 10 afforded the corresponding selenolactam 11 in 92% yield along with the ditert-butyl ketone (12) (Scheme 3), suggesting that the initial adduct of the reaction was oxaselenazolidine 13 or betaine 14. Since the stability of 13 or 14 is lower than that of 7, rearrangement took place under these conditions.

#### Conclusion

3.4-Dihydro-2H-pyrrole N-oxide 1b reacted like a 1,3-dipole with thiocarbonyl compounds. The dissociation constants indicated that the formation of 5-membered ring products was not decelerated by a bulky group attached to the carbon atom adjacent to the 1,3-dipole. Selenium atom transfer through the oxaselenazolidine intermediate took place with microwave irradiation.

#### Experimental

General Methods. Solvents were distilled under a nitrogen atmosphere. All chemicals were obtained from a commercial supplier and used without further purification. Adamantane-2-thione (2a),<sup>7</sup> 2,2,4,4-tetramethyl-3-thioxocyclobutanone (2b),<sup>9</sup> bornane-6-thione (2c),<sup>10</sup> 1,1,3,3-tertamethylindane-2-thione (2d),<sup>11</sup> and di-*tert*-butyl selenoketone  $(10)^8$  were prepared according to literature procedures. Analytical TLC was carried out on precoated plates (Merck, silica gel 60, F254) and flash column chromatography was performed with silica (Merck, 70-230 mesh). NMR spectra (<sup>1</sup>H at 400 MHz; <sup>13</sup>C at 100 MHz; <sup>31</sup>P at 161 MHz) were recorded in CDCl<sub>3</sub>, and the chemical shifts are expressed in ppm relative to the internal standard TMS. <sup>31</sup>PNMR spectra were obtained in CDCl<sub>3</sub> using 85% H<sub>3</sub>PO<sub>4</sub> as an internal standard with broadband <sup>1</sup>H decoupling. The melting points are uncorrected.

Synthesis of 2-Diphenylphosphinoyl-2-methyl-3,4-dihydro-2H-pyrrole N-Oxide (DPhPMPO) (1b). Water (4.5 mL) was added to a solution of chlorodiphenylphosphine (4) (4.6 mL, 25 mmol) in acetonitrile (38 mL) and the mixture was refluxed for 6 h. The reaction mixture was then concentrated under reduced pressure. The residue was extracted with  $CH_2Cl_2$  (20 mL  $\times$  3) and dried over MgSO<sub>4</sub>. The solvent was removed in vacuo to give diphenylphosphine oxide (3) as a colorless oil (97% yield). <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  7.98–8.02 (m, 10H), 8.11 (d, 2H,  $J_{P-H} =$ 468 Hz). <sup>31</sup>P NMR (CDCl<sub>3</sub>)  $\delta$  23.1.

5-Chloro-2-pentanone (6) (2.75 mL, 24 mmol) was added to a solution of diphenylphosphine oxide (4.04 g, 20 mmol) in ethanol (16 mL), and the mixture was stirred at 0 °C for 1 h. After stirring at room temperature, the reaction mixture was warmed up to 60 °C, and then gaseous NH<sub>3</sub> was bubbled into the solution for 1 h. After stirring for 3 h, the solvent was removed under reduced pressure. The residue was dissolved in  $2 \mod L^{-1} \operatorname{HCl}(50 \operatorname{mL})$  and extracted with  $CH_2Cl_2$  (100 mL  $\times$  3). The organic layer was dried over MgSO<sub>4</sub>, and the solvent was removed. The residue was crystallized from ethanol-hexane to give 2-(diphenylphosphinoyl)-2methyl-1-pyrrolidine (5) as colorless crystals (63%). <sup>1</sup>HNMR  $(\text{CDCl}_3) \delta 1.35 \text{ (d, 3H, } J = 14 \text{ Hz}), 1.48-1.50 \text{ (m, 1H)}, 1.62-1.65$ (m, 1H), 1.76–1.80 (m, 1H), 2.11–2.19 (m, 1H), 2.57–2.59 (m, 1H), 3.07-3.09 (m, 1H), 7.41-7.51 (m, 6H), 8.14-8.20 (m, 4H). <sup>13</sup>C NMR (CDCl<sub>3</sub>)  $\delta$  24.8 (*J*<sub>P-C</sub> = 9.1 Hz), 25.9 (*J*<sub>P-C</sub> = 4.6 Hz), 35.1  $(J_{P-C} = 5.1 \text{ Hz})$ , 47.8  $(J_{P-C} = 6.9 \text{ Hz})$ , 62.6  $(J_{P-C} = 94.7 \text{ Hz})$ Hz), 128.2, 128.3, 128.4, 128.5, 130.8, 131.1, 131.5, 131.6, 131.7, 131.8, 132.1, 132.5, 132.6, 132.7, 132.8, 132.9, 133.1, 133.2.  $^{31}$ P NMR (CDCl<sub>3</sub>)  $\delta$  33.1.

Aqueous NaHCO<sub>3</sub> (50 mL, 4.0 g, 48 mmol) and oxone (50 mL, 7.4 g, 12 mmol) were added dropwise to a solution of pyrrolidine 5 in acetone (2.85 g, 10 mmol). After stirring for 1 h at room temperature, the reaction mixture was quenched with 50 mL of 10% sodium thiosulfate and acetone was removed under reduced pressure. The residue was extracted with  $CH_2Cl_2$  (150 mL  $\times$  3) and dried over MgSO<sub>4</sub>. The organic layer was concentrated to give a crude product as a pale yellow oil. Purification was carried out via recrystallization from water-acetonitrile to give 3,4-dihydro-2H-pyrrole N-oxide 1b as colorless crystals (47%). mp 137138 °C (decomp.). <sup>1</sup>H NMR (CDCl<sub>3</sub>), δ 1.72 (d, 3H, J = 6.8 Hz), 1.86–1.97 (m, 1H), 2.12–2.23 (m, 1H), 2.37–2.46 (m, 1H), 3.18– 3.28 (m, 1H), 6.66 (dd, 1H, J = 5.2, 2.4 Hz), 7.45–7.56 (m, 6H), 8.08–8.12 (m, 2H), 8.39–8.44 (m, 2H). <sup>13</sup>C NMR (CDCl<sub>3</sub>) δ 21.3 ( $J_{P-C} = 4.6$  Hz), 25.8, 30.2, 78.7 ( $J_{P-C} = 71.0$  Hz), 128.4, 128.6, 128.7, 128.8, 128.9, 129.3, 129.9, 132.1, 132.2, 132.4, 132.6, 133.2, 133.3, 136.3. <sup>31</sup>P NMR (CDCl<sub>3</sub>) δ 34.3. Found: C, 64.49; H, 6.51; N, 4.40%. Calcd for C<sub>17</sub>H<sub>18</sub>NO<sub>2</sub>P•H<sub>2</sub>O: C, 64.35; H, 6.35; N, 4.41%.

X-ray Crystallographic Data of DPhPMPO (1b). X-ray crystallographic data were collected using a DIP Image Plate with Cu K $\alpha$  radiation. The reflections were collected at room temperature (298 K). An orthorhombic lattice was determined from systematic absence of lattice (hkl, h + k = 2n). A  $P2_12_12_1$  space group was determined from systematic absence (hol, l = 2n). A good model was found by direct methods (SIR97) using the  $P2_12_12_1$ space group.<sup>12</sup> Crystal data for 1b: FW 299.31. Orthorhombic, space group  $P2_12_12_1$ , a = 11.4480(5) Å, b = 10.3370(4) Å, c = 14.3840(6) Å, V = 1702.17(12) Å<sup>3</sup>, Z = 4,  $D_{calcd} = 1.168$  $g \text{ cm}^{-3}$ ,  $\mu = 1.457 \text{ mm}^{-1}$ , R = 0.0778,  $R_w = 0.2131$ . Crystallographic data have been deposited with the Cambridge Crystallographic Data Centre: Deposit No. CCDC-629371 for Compound 1b. Copies of the data can be obtained free of charge via http:// www.ccdc.cam.ac.uk/conts/retrieving.html (or from the Cambridge Crystallographic Data Centre, 12, Union Road, Cambridge, CB2 1EZ, UK: Fax: +44 1223 336033; or e-mail: deposit@ccdc. cam.ac.uk).

**Reaction of 1b with Adamantane-2-thione (2a).** Adamantane-2-thione (**2a**) (16.6 mg, 0.1 mmol) was added to a solution of **1b** (30 mg, 0.1 mmol) in CHCl<sub>3</sub> (1.0 mL), and the mixture was stirred at room temperature for 15 min. After stirring, the reaction mixture was concentrated under reduced pressure to give a white solid, which was recrystallized from hexane–ether to afford colorless crystals of oxathiazolidine **7a** (41.9 mg, 0.09 mmol). **7a**: mp 128–129 °C. <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  1.63 (d, 3H, *J*<sub>P-H</sub> = 15.6 Hz), 1.68–2.19 (m, 18H), 3.96 (dd, 1H, *J* = 6.0, 8.4 Hz), 7.42–7.56 (m, 6H), 7.98–8.02 (t, 2H, *J* = 8.4 Hz), 8.26–8.30 (t, 2H, *J* = 8.4 Hz). <sup>13</sup>C NMR (CDCl<sub>3</sub>)  $\delta$  19.2, 25.3, 25.9, 26.4, 30.2, 32.9, 34.0, 34.7, 34.8, 36.2, 37.3, 38.3, 38.7, 69.64 (*J*<sub>P-C</sub> = 94.0 Hz), 72.3 (*J*<sub>P-C</sub> = 6.3 Hz), 100.9, 127.2, 127.3, 130.7, 130.8, 131.6, 131.7. <sup>31</sup>P NMR (CDCl<sub>3</sub>)  $\delta$  31.7. Found: C, 69.43; H, 6.96; N, 2.93%. Calcd for C<sub>27</sub>H<sub>32</sub>NO<sub>2</sub>PS: C, 69.65; H, 6.93; N, 3.01%.

Reaction of 1b with 2,2,4,4-Tetramethyl-3-thioxocyclobutanone (2b). Thione 2b (15.6 mg, 0.1 mmol) was added to a solution of 1b (30 mg, 0.1 mmol) in CHCl<sub>3</sub> (1.0 mL), and the mixture was stirred at room temperature for 15 min. After stirring, the reaction mixture was concentrated under reduced pressure to give a white solid, which was recrystallized from hexane-ether to afford colorless crystals of oxathiazolidine 7b (27.4 mg, 0.06 mmol). 7b: mp 102–104 °C. <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 1.1 (s, 3H), 1.29 (s, 3H), 1.31 (s, 3H), 1.38–1.46 (m, 1H), 1.41 (s, 3H), 1.61 (d, 3H,  $J_{P-H} = 15$ Hz), 1.94-1.99 (m, 1H), 2.23-2.36 (m, 1H), 2.54-2.64 (m, 1H), 4.71 (brd, 1H, J = 4.8 Hz), 7.36–7.55 (m, 6H), 8.08 (t, 2H, J =8.4 Hz), 8.21 (t, 2H, J = 8.4 Hz). <sup>13</sup>C NMR (CDCl<sub>3</sub>)  $\delta$  18.6, 20.6, 21.0  $(J_{P-C} = 4.4 \text{ Hz})$ , 22.6, 24.5, 27.0, 33.9  $(J_{P-C} = 4.9)$ , 63.4, 68.1, 71.7, 72.7, 75.4 ( $J_{P-C} = 11 \text{ Hz}$ ), 103.6, 128.3, 128.4, 128.5, 132.0, 132.1, 132.8, 132.9, 133.0, 219.4. <sup>31</sup>PNMR (CDCl<sub>3</sub>)  $\delta$ 30.0. Found: C, 65.77; H, 6.65; N, 2.90%. Calcd for C<sub>25</sub>H<sub>30</sub>-NO<sub>3</sub>PS: C, 65.91; H, 6.64; N, 3.07%.

**Dissociation Constants of Oxathiazolidine 7.** DPhPMPO **1b** (0.246 mmol), thioketone (0.246 mmol), and tetrachloroethane (0.246 mmol) in CDCl<sub>3</sub> (5.0 mL) reached equilibrium with oxa-

thiazolidine **7** in 24 h. The integral was compared with TCE (s, 5.96). **7b**: 0.224 mmol by the signal at  $\delta$  4.71 (brd, C-2). **1b**: 0.022 mmol by  $\delta$  6.60 (d, N=C*H*-),  $K_{\text{diss}} = 0.41 \pm 0.041 \times 10^{-3}$  mol L<sup>-1</sup> at 25 °C.

General Procedure of Microwave Irradiation Reaction. The appropriate thioketone 2 or selenoketone 10 was added to a solution of 1b in CHCl<sub>3</sub>, and the mixture was stirred for 1 min. It was then irradiated with a microwave by Electrolux EMC-M20Y (TOSHIBA). The reaction was monitored by TLC. After the irradiation, the reaction mixture was dissolved in CDCl<sub>3</sub> (0.5 mL) for NMR analysis.

**Reaction of 1b with Di-***tert***-butyl Selenoketone 10.** Di-*tert*butyl selenoketone (**10**) (41.0 mg, 0.2 mmol) was added to a solution of **1b** (38.9 mg, 0.13 mmol) in CHCl<sub>3</sub> (2 mL) and irradiated with a microwave for 5 min. The reaction mixture was concentrated under reduced pressure to give a white solid, which was recrystallized from benzene to afford colorless crystals of selenoamide **11** (44.0 mg, 0.12 mmol). **11**: mp 209–211 °C. <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 1.60 (d, 3H, *J* = 15.6 Hz), 1.99–2.13 (m, 2H), 2.59–2.64 (m, 1H), 2.65–2.86 (m, 1H), 7.50–7.67 (m, 6H), 7.83–7.87 (m, 2H), 7.97–8.02 (m, 2H), 8.84 (brs, 1H). <sup>13</sup>C NMR (CDCl<sub>3</sub>) δ 23.0 (*J*<sub>P-C</sub> = 7.4 Hz), 33.4 (*J*<sub>P-C</sub> = 3.4 Hz), 48.0, 71.4 (*J*<sub>P-C</sub> = 75.8 Hz), 127.5, 128.5, 128.6, 129.2, 129.3, 129.4, 129.5, 132.1, 132.3, 132.4, 133.0, 133.1, 207.7 (*J*<sub>P-C</sub> = 3.5 Hz). <sup>31</sup>P NMR (CDCl<sub>3</sub>) δ 31.2. Found: C, 56.37; H, 5.04; N, 3.87%. Calcd for C<sub>17</sub>H<sub>18</sub>NO-PSe: C, 56.36; H, 5.01; N, 3.87%.

#### References

1 For a review see: P. Caramella, P. Grünanger, *1,3-Dipolar Cycloaddition Chemistry*, ed. by A. Padwa, New York, **1984**, Vol. 1, pp. 291–392.

2 a) For a review see: R. Huisgen, *1,3-Dipolar Cycloaddition* Chemistry, ed. by A. Padwa, New York, **1984**, Vol. 1, pp. 1–176. b) L. Fisera, R. Huisgen, I. Kalwinsch, E. Langhals, X. Li, G. Mloston, K. Polborn, J. Rapp, W. Sicking, R. Sustmann, *Pure* Appl. Chem. **1996**, 68, 789. c) R. Huisgen, X. Li, *Tetrahedron* Lett. **1983**, 24, 4185. d) R. Huisgen, E. Langhals, *Tetrahedron* Lett. **1989**, 30, 5369. e) H. Giera, R. Huisgen, *Liebigs Ann./Recl.* **1997**, 1685.

3 D. S. C. Black, K. G. Watson, Aust. J. Chem. 1973, 26, 2491.

4 a) R. Huisgen, L. Fisera, H. Giera, R. Sustmann, J. Am. Chem. Soc. **1995**, 117, 9671. b) R. Sustmann, W. Sicking, R. Huisgen, J. Am. Chem. Soc. **1995**, 117, 9679.

5 G. Mlostoń, T. Gendek, H. Heimgartner, *Helv. Chim. Acta* **1998**, *81*, 1585.

6 K. Shioji, S. Tsukimoto, H. Tanaka, K. Okuma, *Chem. Lett.* **2003**, *32*, 604.

7 J. W. Greidanus, W. J. Schwalm, *Can. J. Chem.* **1969**, *47*, 3715.

8 T. G. Back, D. H. R. Barton, M. R. Britten-Kelly, F. S. Guziec, Jr., J. Chem. Soc., Perkin Trans. 1 1976, 2079.

9 E. U. Elam, H. E. Davis, J. Org. Chem. 1967, 32, 1562.

10 J. W. Scheeren, P. H. J. Ooms, R. J. F. Nivard, *Synthesis* 1973, 149.

11 A. Ishii, R. Okazaki, N. Inamoto, Bull. Chem. Soc. Jpn. **1988**, 61, 861.

12 SIR97, Program for the Solution of Crystal Structures: A. Altomare, M. C. Burla, M. Camalli, G. L. Cascarano, C. Giacovazzo, A. Guagliardi, A. G. G. Moliterni, G. Polidori, R. Spagna, *J. Appl. Crystallogr.* **1999**, *32*, 115.