

New Stable Dication of 1,3,4,6-Tetrakis(isopropylthio)-2 λ^4 , δ^2 -thieno[3,4-*c*]thiophene generated by its Oxidation using Nitrosonium Tetrafluoroborate


Akira Tsubouchi, Noboru Matsumura and Hiroo Inoue*

Department of Applied Chemistry, College of Engineering, University of Osaka Prefecture, Sakai, Osaka 591, Japan


The oxidation of 1,3,4,6-tetrakis(isopropylthio)-2 λ^4 , δ^2 -thieno[3,4-*c*]thiophene **1** by NOBF₄ (2 equiv.) gives the dication **2**, which reacts with H₂O, MeOH and aniline to give the thiophene derivatives **3–5** by fission of the C–S bond of the thieno[3,4-*c*]thiophene ring; **2** also reacts with 1,2-dimercaptobenzene and NaBH₄ to give the adducts **6** and **7** respectively.

1,3,4,6-Tetrakis(alkylthio)-2 λ^4 , δ^2 -thieno[3,4-*c*]thiophenes¹ are interesting compounds which have a nonclassical structure with 10 π -electrons. Previously we have disclosed that the cyclic voltammogram of 1,3,4,6-tetrakis(isopropylthio)-2 λ^4 , δ^2 -thieno[3,4-*c*]thiophene **1** in MeCN shows two redox waves with oxidation potentials at $E_1 = +0.19$ V and $E_2 = +0.54$ V vs. saturated calomel electrode (SCE) and **1** is more electron-donating than tetrathiafulvalene.² These findings led us to

oxidise **1** chemically to its dication which is an unknown sulphur analogue of pentalene with 8 π -electrons. We now report a new stable dication, 1,3,4,6-tetrakis(isopropylthio)-thieno[3,4-*c*]thiophenium bis(tetrafluoroborate) **2**, which is prepared from **1** by a two-electron oxidation using nitrosonium tetrafluoroborate, NOBF₄, as the oxidant (Scheme 1). Furthermore, we describe that the reaction of **2** with H₂O, MeOH and aniline gives the thiophene derivatives **3–5** by the



Scheme 1



Scheme 2 Reagents and conditions: i, H₂O (large excess), room temp., 1 h; ii, MeOH (10 equiv.), -30 °C, 2 h; iii, PhNH₂ (10 equiv.), -30 °C, 0.5 h; iv, 1,2-dimercaptobenzene (1.2 equiv.), -30 °C, 1 h; v, NaBH₄ (10 equiv.), -30 °C, 2 h

fission of the C–S bond of the thieno[3,4-*c*]thiophene ring and that reaction with 1,2-dimercaptobenzene and NaBH₄ gives the adducts **6** and **7** respectively.

† Selected spectroscopic data: Compound **2**: ¹H NMR (CD₃CN, -30 °C) δ 1.71 (24H, d), 4.26 (4H, br sept.); ¹³C NMR (CD₃CN, -30 °C) δ 22.4, 51.2, 139.5, 176.2; UV-VIS (MeCN) λ_{max} 274 (log ε 4.34), 295 (4.45), 418 (4.24), 579 nm (4.51); satisfactory elemental analysis was obtained.

Compound **3**: ¹H NMR (CDCl₃) δ 1.28 (6H, d), 1.33 (6H, d), 1.34 (6H, d), 1.44 (6H, d), 3.32 (1H, sept.), 3.38 (1H, sept.), 3.78 (1H, sept.), 4.07 (1H, sept.); IR ν_{max}/cm⁻¹ (neat) 1690 (C=O); *m/z* 452 (M⁺).

Compound **4**: ¹H NMR (CDCl₃) δ 1.27 (6H, d), 1.41 (6H, d), 1.45 (6H, d), 3.28 (1H, sept.), 3.48 (1H, sept.), 3.73 (3H, s), 4.10 (1H, sept.); IR ν_{max}/cm⁻¹ (neat) 1700 (C=O); *m/z* 408 (M⁺).


Compound **5**: ¹H NMR [(CD₃)₂SO, 100 °C] δ 1.23 (6H, d), 1.25 (6H, d), 1.42 (6H, d), 3.32 (1H, sept.), 3.4–3.6 (1H, br sept.), 4.01 (1H, sept.), 6.84 (2H, d), 6.99 (1H, t), 7.20 (2H, t); IR ν_{max}/cm⁻¹ (neat) 1600 (C=N); *m/z* 527 (M⁺).

Compound **6**: ¹H NMR (CDCl₃) δ 1.33 (6H, d), 1.34 (12H, d), 1.45 (6H, d), 3.33 (2H, sept.), 3.39 (2H, sept.), 7.17 (2H, dd), 7.51 (2H, dd); *m/z* 500 (M⁺ - 76).

Compound **7**: ¹H NMR (CDCl₃) δ 1.30 (6H, d), 1.32 (12H, d), 1.36 (6H, d), 3.35 (2H, sept.), 3.37 (2H, sept.), 5.46 (2H, s); *m/z* 438 (M⁺). The *cis* configuration of **7** was assigned on the basis of the fact that the ¹H NMR spectrum of **7** agreed with that of the product prepared by the hydrogenation of **1** over Pd/C catalyst.³

The reaction of **1** with NOBF₄ was carried out as follows. A solution of NOBF₄ (177 mg, 1.51 mmol) in MeCN (20 ml) was added under nitrogen to a suspended solution of **1** (300 mg, 0.688 mmol) in MeCN (7.5 ml) at -30 °C. A blue–purple colour appeared immediately and the mixture became homogeneous. After stirring for 1 h at -30 °C, dry Et₂O (200 ml) was added to the reaction mixture. The precipitate was filtered off, washed with ether and dried under vacuum to give the dication **2** (350 mg, 84%) as a reddish-purple solid [m.p. 147–148 °C (decomp.)].† The formation of the dication **2** indicates that the addition of NO⁺ to the 1-position of **1** does not occur. The dication **2** was quite stable under nitrogen at room temperature. The ¹H NMR spectrum of **2** at -30 °C exhibited a doublet for methyl protons and a septet for methine protons at lower field than those of **1** by 0.42 and 0.82 ppm respectively. Furthermore, the peaks of the ring carbons of **2** in its ¹³C NMR spectrum at -30 °C also appeared at lower field than those of **1** by 23.1 and 25.0 ppm respectively. These downfield shifts are attributable to the positive charge delocalised in a π-electron system of **2**. In the UV-VIS spectrum of **2** in MeCN, the absorption peak at the longest wavelength was shifted 84 nm to a longer wavelength than that of **1** (513 nm).

The dication **2** was reduced by *N,N,N',N'*-tetramethyl-*p*-phenylenediamine (TMPD) (2 equiv.) in MeCN to give **1** in 73% yield by the transfer of two electrons, accompanied by the formation of TMPD^{•+} which was confirmed by the measurement of the UV-VIS spectrum [eqn. (1)]. The



reaction of **2** with various nucleophiles was carried out under nitrogen in MeCN (Scheme 2). It was found that the dication **2** reacts readily with H₂O, MeOH and aniline to give the ring-opening products **3**, **4** and **5** in 98, 99 and 88% yields respectively.† This reaction is considered to proceed by the attack of the nucleophiles on the 1-position of **2**, followed by the fission of the C(3)–S bond of the ring. When 1,2-dimercaptobenzene and NaBH₄ were used, the adducts **6** and **7** were obtained in 63 and 95% yields respectively.†

Thus, the present results provide evidence for the formation of the stable dication **2** in the reaction of **1** with NOBF₄.

This work was supported by the Grant-in Aid for Scientific Research on Priority Area No. 02230226 from the Japanese Ministry of Education, Science and Culture.

Received, 23rd January 1991; Com. 1/00348H

References

- 1 S. Yoneda, K. Ozaki, A. Tsubouchi and H. Kojima, *J. Heterocycl. Chem.*, 1988, **25**, 599; A. Tsubouchi, N. Matsumura, H. Inoue, N. Hamasaki, S. Yoneda and K. Yanagi, *J. Chem. Soc., Chem. Commun.*, 1989, 223; A. Tsubouchi, N. Matsumura and H. Inoue, *Chem. Lett.*, 1990, 1853. For other derivatives, see: M. P. Cava and M. V. Lakshmikantham, *Acc. Chem. Res.*, 1975, **8**, 139; C. A. Ramsden in *Comprehensive Heterocyclic Chemistry*, ed. K. T. Potts, Pergamon Press, Oxford, 1984, vol. 6, p. 1027; J. Nakayama, A. Ishii, Y. Kobayashi and M. Hoshino, *J. Chem. Soc., Chem. Commun.*, 1988, 959.
- 2 S. Yoneda, A. Tsubouchi and K. Ozaki, *Nippon Kagaku Kaishi*, 1987, 1328.
- 3 M. P. Cava, M. Behforouz, G. E. M. Husbands and M. Srinivasan, *J. Am. Chem. Soc.*, 1973, **95**, 2561.