

**Reactivity of  $[M(C\wedge P)(S_2C-R)]$  ( $M = Pd, Pt$ ;  $C\wedge P = CH_2-C_6H_4-P(o-tolyl)_2-\kappa C,P$ ;  $R = NMe_2, OEt$ ) toward  $HgX_2$  ( $X = Br, I$ ). X-ray Crystal Structures of  $[Pt\{CH_2-C_6H_4P(o-tolyl)_2-\kappa C,P\}(S_2CNMe_2)HgI(\mu-I)]_2$  and  $[PdBr(S_2COEt)\{\mu-P(o-tolyl)_2-C_6H_4-CH_2-\}HgBr]\cdot 0.5HgBr_2\cdot C_2H_4Cl_2$**

Larry R. Falvello, Juan Forniés,\* Antonio Martín, Rafael Navarro, Violeta Sicilia, and Pablo Villarroya

Departamento de Química Inorgánica, Instituto de Ciencia de Materiales de Aragón, Universidad de Zaragoza-CSIC, Facultad de Ciencias, E-50009 Zaragoza, Spain

Received May 15, 1997<sup>®</sup>

The reaction of the complexes  $[Pt(C\wedge P)(S_2C-R)]$  ( $C\wedge P = CH_2-C_6H_4-P(o-tolyl)_2-\kappa C,P$ ,  $R = NMe_2, OEt$ ) with an equimolar amount of  $HgX_2$  ( $X = Cl, Br$ ) gives the tetranuclear derivatives  $[Pt(C\wedge P)(S_2C-R)HgX(\mu-X)]_2$  [ $R = NMe_2$ ,  $X = Br$  (**3**), **I** (**4**);  $R = OEt$ ,  $X = Br$  (**5**), **I** (**6**)] containing  $Pt \rightarrow Hg$  donor–acceptor bonds. The reaction of  $[Pd(C\wedge P)(S_2CNMe_2)]$  with  $HgI_2$  affords the complex  $[Pd(C\wedge P)(S_2CNMe_2)HgI(\mu-I)]_2$  (**9**) similar to the complexes **3–6**; by contrast the reaction of  $[Pd(C\wedge P)(S_2C-R)]$  ( $R = NMe_2, OEt$ ) with  $HgBr_2$  leads to the corresponding dinuclear complexes  $[PdBr(S_2C-R)(\mu-C\wedge P)HgBr]$  [ $R = NMe_2$  (**10**),  $OEt$  (**11**)] with the didentate  $C\wedge P$  cyclometalating ligand,  $-CH_2-C_6H_4-P(o-tolyl)_2-\kappa C,P$  (resulting from the C–H activation of the  $P(o-tolyl)_3$ ) acting in an unprecedented bridging mode. Compound **4** ( $C_{24}H_{26}HgI_2NPtS_2$ ) crystallizes in the triclinic system, space group  $P1$ :  $a = 9.5755(11)$  Å,  $b = 11.1754(12)$  Å,  $c = 14.504(2)$  Å,  $\alpha = 84.826(5)^\circ$ ,  $\beta = 81.611(7)^\circ$ ,  $\gamma = 68.606(9)^\circ$ ,  $V = 1428.5(3)$  Å<sup>3</sup>, and  $Z = 1$ . Compound **11**·0.5  $HgBr_2\cdot C_2H_4Cl_2$  ( $C_{24}H_{25}Br_2HgOPdS_2\cdot 0.5 HgBr_2\cdot C_2H_4Cl_2$ ) crystallizes in the monoclinic system, space group  $P2_1/c$ :  $a = 15.571(2)$  Å,  $b = 10.7425(10)$  Å,  $c = 19.655(2)$  Å,  $\beta = 94.741(12)^\circ$ ,  $V = 3276.5(5)$  Å<sup>3</sup>, and  $Z = 4$ .

## Introduction

Metal–metal bonding in heteronuclear complexes containing  $d^8$  and  $d^{10}$  metal ions has been known for some years.<sup>1</sup> We have developed chemistry for  $Pt^{II}$  and  $Ag^I$  using mono- or dinuclear anionic pentahalophenyl complexes of  $Pt(II)$  as Lewis bases,<sup>2</sup> and, more recently, heteronuclear  $Pt \rightarrow Hg$  complexes obtained from this kind of  $Pt(II)$  substrates have also been reported.<sup>3</sup>

The ability of mercury to form heterodimetallic complexes with transition metal complex fragments is well-known.<sup>4</sup> Some of these products have been prepared by reaction of  $HgX_2$  ( $X = Cl, Br, I$ ) with  $d^8$  transition metals,<sup>5–13</sup> and the resulting complexes can be divided into those with a covalent metal–metal bond<sup>5–9</sup> (type I) and those with a metal to metal donor–acceptor bond<sup>10–13</sup> (type II). Reactions of  $HgX_2$  ( $X = Cl, Br,$

**I**) with organoplatinum(II) compounds have been found to proceed differently: In some cases, oxidative addition of  $HgX_2$  to  $Pt^{II}$  renders compounds containing a  $Pt^{IV}-Hg$  covalent bond, as seems to occur in the reaction of  $cis-[PtMe_2(Ph_2Me_2phen)]$  with  $HgX_2$  ( $X = Cl, Br, I$ ),<sup>9</sup> and in other cases the oxidative addition process is followed by a reductive elimination so that the platinum substrate acts as an alkylating agent toward the  $Hg$  center, as happens in the reactions of  $cis-[PtMe_2(PPhMe_2)]$  and  $cis-[PtMe_2(bpy)]$  with  $HgCl_2$ .<sup>14</sup> As far as we know, no examples of  $Pt^{II} \rightarrow HgX_2$  adducts have yet been prepared by reacting the  $Pt$  substrate with  $HgX_2$ . In comparison to the situation with platinum, the chemistry of  $Pd^{II}$  derivatives with  $HgX_2$  is much less developed.<sup>15</sup>

We have previously reported the reactivity of  $[Pt\{CH_2-C_6H_4-P(o-tolyl)_2-\kappa C,P\}(S_2C-R)]$  ( $R = NMe_2, OEt$ ) toward halogens ( $Cl_2, Br_2, I_2$ ) to give the  $Pt(IV)$  compounds  $[Pt\{CH_2-C_6H_4-P(o-tolyl)_2-\kappa C,P\}(S_2C-R)X_2]$  ( $R = NMe_2, OEt$ ;  $X = Cl, Br, I$ ) as a result of the oxidative addition of  $X_2$  to the  $Pt(II)$  substrates.<sup>16</sup> Since  $Hg(II)$  salts are able to add oxidatively to  $d^8$  metal fragments affording heteronuclear complexes containing metal–metal bonds, we have extended this work to the study of the reactivity of such mononuclear  $Pt(II)$  complexes  $[Pt\{CH_2-C_6H_4-P(o-tolyl)_2-\kappa C,P\}(S_2C-R)]$  ( $R = NMe_2, OEt$ ) toward  $HgX_2$  ( $X = Br, I$ ) and additionally to a study of similar  $Pd(II)$  complexes. As a result of this work, we report here the synthesis, structural characterization, and NMR studies of new mixed  $Pt-Hg$  and  $Pd-Hg$  complexes.

<sup>®</sup> Abstract published in *Advance ACS Abstracts*, December 1, 1997.

- (1) (a) Coffey, E.; Lewis, J.; Nyholm, R. S. *J. Chem. Soc.* **1964**, 1741. (b) Kuyper, J.; Vrieze, K. *J. Organomet. Chem.* **1976**, 107, 129.
- (2) (a) Usón, R.; Forniés, J. *Adv. Organomet. Chem.* **1988**, 28, 219. (b) Usón, R.; Forniés, J. *Inorg. Chim. Acta* **1992**, 198–200, 165.
- (3) Usón, R.; Forniés, J.; Falvello, L. R.; Ara, I.; Usón, I. *Inorg. Chim. Acta* **1993**, 212, 105.
- (4) Gade, L. H. *Angew. Chem., Int. Ed. Engl.* **1993**, 32, 24.
- (5) Brotherton, P. D.; Raston, C. L.; White, A. H.; Wild, S. B. *J. Chem. Soc., Dalton Trans.* **1976**, 1799.
- (6) Tiripicchio, A.; Lahoz, F. J.; Oro, L. A.; Pinillos, M. T. *J. Chem. Soc., Chem. Commun.* **1984**, 936.
- (7) Einstein, F. W. B.; Yan, X.; Zhang, X.; Sutton, D. *J. Organomet. Chem.* **1992**, 439, 221.
- (8) Brekau, U.; Werner, H. *Organometallics* **1990**, 9, 1067.
- (9) Kuyper, J. *Inorg. Chem.* **1978**, 17, 1458.
- (10) Faraone, F.; Schiavo, S. L.; Bruno, G.; Bombieri, G. *J. Chem. Soc., Chem. Commun.* **1984**, 6.
- (11) Faraone, F.; Tresoldi, G.; Loprete, G. A. *J. Chem. Soc., Dalton Trans.* **1979**, 933.
- (12) Nowell, I. W.; Russell, D. R. *J. Chem. Soc., Dalton Trans.* **1972**, 2393.
- (13) Faraone, F.; Bruno, G.; Tresoldi, G.; Faraone, G.; Bombieri, G. *J. Chem. Soc., Dalton Trans.* **1981**, 1651.

- (14) (a) Cross, R. J.; Wardle, R. *J. Chem. Soc. A* **1970**, 840. (b) Jawad, J. K.; Puddephatt, R. J. *J. Chem. Soc., Chem. Commun.* **1977**, 892. (c) Jawad, J. K.; Puddephatt, R. J. *Inorg. Chim. Acta* **1978**, 31, L391.
- (15) Barr, R. M.; Goldstein, M.; Hairs, N. D.; McPartlin, M.; Markwell, A. J. *J. Chem. Soc., Chem. Commun.* **1974**, 221.
- (16) Forniés, J.; Martín, A.; Navarro, R.; Sicilia, V.; Villarroya, P. *Organometallics* **1996**, 15, 1826.

## Experimental Section

Elemental analyses were performed on a Perkin-Elmer 240-B micro-analyzer. IR spectra were recorded on a Perkin-Elmer 599 spectrophotometer (Nujol mulls between polyethylene plates in the range 200–4000  $cm^{-1}$ ). NMR spectra were recorded on either a Varian XL-200 or a Varian Unity 300 NMR spectrometer using the standard references.  $[Pd(OOCCH_3)_2]_2$ ,<sup>17</sup>  $[Pt\{CH_2-C_6H_4-P(o-tolyl)_2-kC,P\}(S_2CNMe_2)]$  (**1**)<sup>16</sup> and  $[Pt\{CH_2-C_6H_4-P(o-tolyl)_2-kC,P\}(S_2COEt)]$  (**2**)<sup>16</sup> were prepared as described elsewhere.

**Safety Note.** Caution! Perchlorate salts are potentially explosive. Only small amounts of material should be prepared, and these should be handled with great caution.

**$[Pd\{CH_2-C_6H_4-P(o-tolyl)_2-kC,P\}(\mu-OOCCH_3)]_2$ .**<sup>18</sup> A mixture of  $[Pd(OOCCH_3)_2]$  (1.216 g, 5.42 mmol) and  $P(o-tolyl)_3$  (1.649 g, 5.42 mmol) was refluxed in  $Me_2CO$  (45 mL) for 1 h. The dark green precipitate was filtered off and recrystallized from  $CH_2Cl_2/Et_2O$  (yield: 2.012 g, 79%).

**$[Pd\{CH_2-C_6H_4-P(o-tolyl)_2-kC,P\}(\mu-Cl)]_2$ .** To a solution of  $[Pd\{CH_2-C_6H_4-P(o-tolyl)_2-kC,P\}(\mu-OOCCH_3)]_2$  (0.119 g, 2.38 mmol) in  $MeOH/CH_2Cl_2$  (25:25 mL) was added  $LiCl$  (0.202 g, 4.77 mmol), and a dark green solid precipitated immediately. The solid was filtered off, washed with  $Et_2O$ , and dried in vacuo (yield: 1.06 g, 99.7%). IR data ( $cm^{-1}$ ):  $C\wedge P$ , 460 (vs), 470 (vs), 513 (s), 523 (vs), 560 (vs), 581 (vs), 753 (vs, br), 1568 (w), 1583 (m), 1590 (m);  $\nu(Pd-Cl)$ , 247 (s), 289 (s). Anal. Calcd for  $C_{42}Cl_2H_{40}P_2Pd_2$ : C, 56.65; H, 4.53. Found: C, 56.34; H, 4.17.

**$[Pt\{CH_2-C_6H_4-P(o-tolyl)_2-kC,P\}(S_2CNMe_2)HgX(\mu-X)]_2$  ( $X = Br$  (**3**), **I**(**4**)).**  $X = Br$  (**3**). To a pale yellow solution of compound **1** (0.2 g, 0.32 mmol) in  $CH_2Cl_2$  (15 mL) was added  $HgBr_2$  (0.116 g, 0.32 mmol), and the color of the solution turned orange immediately. The mixture was stirred for 10 min; then, the solvent was evaporated to dryness, and  $n$ -hexane (20 mL) was added to the residue (yield: 0.275 g, 87%). IR data ( $cm^{-1}$ ):  $C\wedge P$ , 456 (s), 474 (s), 488 (m), 504 (m), 528 (s), 562 (m), 586 (s), 758 (s), 774 (s), 1584 (w), 1595 (w);  $-S_2CNMe_2$ , 1554 (vs) ( $\nu(C-N)$ ), 968 (m) ( $\nu(C-S)$ ).  $^1H$  NMR:  $C\wedge P$ , 3.94 (s, 2H) ( $^2J_{Pt-H} = 85.30$  Hz, 75.80 Hz) ( $-CH_2-Pt$ ), 2.37 (s), 2.61 (s) ( $CH_3-C_6H_4$ ), 6.8–7.6 (m) ( $C_6H_4$ );  $-S_2CNMe_2$ , 3.33 (s), 3.34 (s). Anal. Calcd for  $Br_4C_{48}H_{52}Hg_2N_2P_2Pt_2S_4$ : C, 29.44; H, 2.68; N, 1.43. Found: C, 29.25; H, 2.64; N, 1.42.

$X = I$  (**4**). Compound **1** (0.151 g, 0.24 mmol) and  $HgI_2$  (0.111 g, 0.24 mmol) in  $CH_2Cl_2$  (25 mL) were reacted for 40 min. The solution was filtered and evaporated to dryness, whereupon addition of  $n$ -pentane (20 mL) gave compound **4** as an orange solid (yield: 0.228 g, 87%). IR data ( $cm^{-1}$ ):  $C\wedge P$ , 456 (s), 473 (s), 486 (s), 503 (m), 528 (s), 562 (s), 585 (s), 756 (s), 762 (s), 773 (s), 1584 (w), 1592 (w);  $-S_2CNMe_2$ , 1552 (vs) ( $\nu(C-N)$ ), 969 (m) ( $\nu(C-S)$ ).  $^1H$  NMR:  $C\wedge P$ , 3.88 (s, 2H) ( $^2J_{Pt-H} = 95.20$  Hz, 70.34 Hz) ( $-CH_2-Pt$ ), 2.37 (s), 2.64 (s) ( $CH_3-C_6H_4$ ), 6.8–7.8 (m) ( $C_6H_4$ );  $-S_2CNMe_2$ , 3.32 (s), 3.37 (s). Anal. Calcd for  $C_{48}H_{52}Hg_2I_4N_2P_2Pt_2S_4$ : C, 26.86; H, 2.44; N, 1.31. Found: C, 26.73; H, 2.39; N, 1.29.

**$[Pt\{CH_2-C_6H_4-P(o-tolyl)_2-kC,P\}(S_2COEt)HgX(\mu-X)]_2$  ( $X = Br$  (**5**), **I**(**6**)).**  $X = Br$  (**5**). To a yellow solution of compound **2** (0.218 g, 0.35 mmol) in  $CH_2Cl_2$  (25 mL) was added  $HgBr_2$  (0.127 g, 0.35 mmol); the color of the solution turned to a brighter yellow. The mixture was stirred for 5 min and the solution evaporated to dryness. Addition of  $n$ -pentane (30 mL) to the residue afforded compound **5** as a yellow solid (yield: 0.29 g, 84%). IR data ( $cm^{-1}$ ):  $C\wedge P$ , 464 (m), 482 (m), 492 (m), 510 (w), 529 (s), 569 (m), 597 (s), 758 (s), 779 (s), 1579 (w), 1597 (w), 1602 (w);  $-S_2COEt$ , 1022 (vs), 1126 (s), 1275 (vs).  $^1H$  NMR:  $C\wedge P$ , 3.98 (s, 2H) ( $^2J_{Pt-H} = 103.44$  Hz, 72.41 Hz) ( $-CH_2-Pt$ ), 2.39 (s), 2.65 (s) ( $CH_3-C_6H_4$ ), 6.8–7.6 (m) ( $C_6H_4$ );  $-S_2COEt$ , 4.70 (q), 1.53 (t) ( $^3J_{H-H} = 7.10$  Hz). Anal. Calcd for  $Br_4C_{48}H_{50}Hg_2O_2P_2Pt_2S_4$ : C, 29.41; H, 2.57. Found: C, 29.12; H, 2.47.

$X = I$  (**6**). Complex **6** was prepared in a similar way. Compound **2** (0.135 g, 0.22 mmol) and  $HgI_2$  (0.099 g, 0.22 mmol) were mixed in  $CH_2Cl_2/OEt_2$  (10/40 mL) for 30 min. After filtration the resulting

orange solution was evaporated to dryness. Addition of  $n$ -pentane (30 mL) to the residue afforded compound **6** as an orange solid (yield: 0.162 g, 70%). IR data ( $cm^{-1}$ ):  $C\wedge P$ , 450 (w), 479 (m), 490 (m), 512 (w), 526 (s), 568 (m), 599 (s), 756 (s), 788 (s), 1574 (m), 1605 (m);  $-S_2COEt$ , 1024 (vs), 1135 (s), 1278 (vs).  $^1H$  NMR:  $C\wedge P$ , 3.71 (s, 2H) ( $^2J_{Pt-H} = 97.30$  Hz, 85.10 Hz) ( $-CH_2-Pt$ ), 2.40 (s), 2.69 (s) ( $CH_3-C_6H_4$ ), 6.8–7.6 (m) ( $C_6H_4$ );  $-S_2COEt$ , 4.64 (q), 1.49 (t) ( $^3J_{H-H} = 7.10$  Hz). Anal. Calcd for  $C_{48}H_{50}I_4Hg_2O_2P_2Pt_2S_4$ : C, 26.84; H, 2.35. Found: C, 26.64; H, 2.28.

**$[Pd\{CH_2-C_6H_4-P(o-tolyl)_2-kC,P\}(S_2CNMe_2)]$  (**7**).** To a stirred suspension of  $[Pd\{CH_2-C_6H_4-P(o-tolyl)_2-kC,P\}(\mu-Cl)]_2$  (0.475 g, 0.53 mmol) in tetrahydrofuran (THF, 30 mL) was added  $AgClO_4$  (0.221 g, 1.07 mmol), and the mixture was stirred at room temperature for 15 min. The  $AgCl$  precipitated was filtered off, and the resulting solution was evaporated almost to dryness. The residue was treated with 30 mL of methanol, and the addition of  $NaS_2CNMe_2 \cdot 2H_2O$  (0.191 g, 1.07 mmol) to the solution gave a yellow solid **7**, which was filtered off and washed with  $MeOH$  and  $n$ -pentane (yield: 0.534 g, 94%). IR data ( $cm^{-1}$ ):  $C\wedge P$ , 445 (s), 461 (s), 468 (s), 480 (s), 509 (s), 525 (s), 558 (s), 578 (s), 754 (vs), 761 (vs), 1581 (m);  $-S_2CNMe_2$ , 1531 (vs) ( $\nu(C-N)$ ), 974 (s) ( $\nu(C-S)$ ).  $^1H$  NMR:  $C\wedge P$ , 3.38 (s) ( $-CH_2-Pd$ ), 2.44 (s), 2.74 (s) ( $CH_3-C_6H_4$ ), 6.7–7.4 (m) ( $C_6H_4$ );  $-S_2CNMe_2$ , 3.29 (s), 3.34 (s). Anal. Calcd for  $C_{24}H_{26}NPPdS_2$ : C, 54.39; H, 4.94; N, 2.64. Found: C, 53.95; H, 4.33; N, 2.88.

**$[Pd\{CH_2-C_6H_4-P(o-tolyl)_2-kC,P\}(S_2COEt)]$  (**8**).** To the solution resulting from the treatment of  $[Pd\{CH_2-C_6H_4-P(o-tolyl)_2-kC,P\}(\mu-Cl)]_2$  (0.469 g, 0.53 mmol) with  $AgClO_4$  (0.218 g, 1.05 mmol), in THF (20 mL), after elimination of the  $AgCl$ , was added  $KS_2COEt$  (0.169 g, 1.05 mmol), and the mixture was stirred for 15 min at room temperature. After filtration the solution was evaporated to dryness,  $n$ -hexane (25 mL) was added, and the solution was concentrated to a small volume (5 mL). The resulting suspension was cooled at 5 °C to afford a yellow solid which was recrystallized from  $CH_2Cl_2/n$ -hexane (−20 °C), compound **8** (yield: 0.4253 g, 76%). IR data ( $cm^{-1}$ ):  $C\wedge P$ , 461 (s), 470 (s), 478 (s), 511 (s), 527 (s), 559 (s), 579 (s), 752 (vs), 761 (vs), 1583 (m), 1591 (w);  $-S_2COEt$ , 1030 (s), 1203 (s).  $^1H$  NMR:  $C\wedge P$ , 3.48 (s) ( $-CH_2-Pd$ ), 2.39 (s), 2.66 (s) ( $CH_3-C_6H_4$ ), 6.6–7.4 (m) ( $C_6H_4$ );  $-S_2COEt$ , 4.56 (q), 1.39 (t) ( $^3J_{H-H} = 7.20$  Hz). Anal. Calcd for  $C_{24}H_{25}OPPdS_2$ : C, 54.20; H, 4.75. Found: C, 54.49; H, 4.76.

**$[Pd\{CH_2-C_6H_4-P(o-tolyl)_2-kC,P\}(S_2CNMe_2)HgI(\mu-I)]$  (**9**).** To a yellow solution of compound **7** (0.216 g, 0.41 mmol) in  $CH_2Cl_2$  (25 mL) was added an equimolar amount of  $HgI_2$  (0.185 g, 0.41 mmol) and  $OEt_2$  (45 mL). The mixture was stirred for 1 h, and the resulting yellow solid **9** was filtered off and dried in vacuo. An additional amount of compound **9** was obtained from the resulting solution by evaporation almost to dryness and addition of  $n$ -pentane (20 mL) to the residue (total yield: 0.36 g, 90%). IR data ( $cm^{-1}$ ):  $C\wedge P$ , 451 (s), 471 (s), 481 (s), 500 (m), 520 (s), 559 (m), 584 (m), 751 (vs), 758 (vs), 1584 (m), 1591 (m);  $-S_2CNMe_2$ , 1549 (vs) ( $\nu(C-N)$ ), 971 (vs) ( $\nu(C-S)$ ).  $^1H$  NMR:  $C\wedge P$ , 3.59 (s) ( $-CH_2-Pd$ ), 2.50 (s), 2.70 (s) ( $CH_3-C_6H_4$ ), 6.8–7.5 (m) ( $C_6H_4$ );  $-S_2CNMe_2$ , 3.44 (s, 6H). Anal. Calcd for  $C_{48}H_{52}Hg_2I_4N_2P_2Pd_2S_4$ : C, 29.28; H, 2.66; N, 1.42. Found: C, 29.33; H, 2.59; N, 1.47.

**$[PdBr(S_2CNMe_2)\{\mu-P(o-tolyl)_2-C_6H_4-CH_2\}HgBr]$  (**10**).** To a yellow solution of compound **7** (0.15 g, 0.28 mmol) in  $CH_2Cl_2$  (25 mL) at −25 °C was added an equimolar amount of  $HgBr_2$  (0.102 g, 0.28 mmol), whereupon the color of the solution turned orange. After 40 min of stirring, the mixture was filtered, and the resulting solution was evaporated to dryness. Addition of  $Et_2O$  (−20 °C) to the residue rendered a yellow solid, **10** (yield: 0.195 g, 77%). IR data ( $cm^{-1}$ ):  $C\wedge P$ , 442 (m), 461 (s), 478 (s), 501 (m), 522 (m), 537 (s), 551 (m), 568 (s), 754 (vs), 1591 (m);  $-S_2CNMe_2$ , 1554 (vs) ( $\nu(C-N)$ ), 970 (vs) ( $\nu(C-S)$ ).  $^1H$  NMR:  $C\wedge P$ , 3.43 (d, 1H), 3.72 (d, 1H,  $^2J_{H-H} = 9.1$  Hz) ( $-CH_2-Pd$ ), 1.52 (s), 1.76 (s) ( $CH_3-C_6H_4$ ), 6.8–7.6 (m, 11H), 9.12 (d, 1H) ( $C_6H_4$ );  $-S_2CNMe_2$ , 3.12 (s), 3.28 (s). Anal. Calcd for  $Br_2C_{24}H_{26}HgNPPdS_2$ : C, 32.37; H, 2.94; N, 1.57. Found: C, 32.17; H, 2.90; N, 1.62.

**$[PdBr(S_2COEt)\{\mu-P(o-tolyl)_2-C_6H_4-CH_2\}HgBr]$  (**11**).** To a solution of compound **8** (0.125 g, 0.24 mmol) in  $CHCl_3$  (30 mL) at −30 °C was added  $HgBr_2$  (0.17 g, 0.47 mmol); the color of the solution turned orange. After 2 h, the mixture was filtered through celite and the filtrate was evaporated almost to dryness. Addition of cold  $n$ -pentane (−20 °C) to the residue afforded compound **11** as a yellow

(17) Heyn, B.; Hipler, B.; Kreisel, G.; Schreer, H.; Walther, D. *Anorganische Syntheseschemie*; Springer-Verlag: Berlin, 1986; p 142.

(18) This complex was prepared recently by Herrmann in a different way: Herrmann, W. A.; Brossmer, Ch.; Öfele, K.; Reisinger, C.-P.; Priermeier, T.; Beller, M.; Fischer, H. *Angew. Chem., Int. Ed. Engl.* 1995, 34, 1844.

**Table 1.** Crystallographic Data for [Pt{CH<sub>2</sub>-C<sub>6</sub>H<sub>4</sub>P(*o*-tolyl)<sub>2</sub>-κC,P}(S<sub>2</sub>CNMe<sub>2</sub>)HgI(μ-I)]<sub>2</sub> (**4**) and [PdBr(S<sub>2</sub>COEt){μ-P(*o*-tolyl)<sub>2</sub>-C<sub>6</sub>H<sub>4</sub>-CH<sub>2</sub>-}HgBr]·0.5 HgBr<sub>2</sub>·C<sub>2</sub>H<sub>4</sub>Cl<sub>2</sub> (**11**·0.5 HgBr<sub>2</sub>·C<sub>2</sub>H<sub>4</sub>Cl<sub>2</sub>)


|                                          | <b>4</b>                                                                                                    | <b>11</b> ·0.5 HgBr <sub>2</sub> ·C <sub>2</sub> H <sub>4</sub> Cl <sub>2</sub>                                                           |
|------------------------------------------|-------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------|
| empirical formula                        | C <sub>48</sub> H <sub>52</sub> Hg <sub>2</sub> I <sub>2</sub> N <sub>2</sub> P <sub>2</sub> S <sub>4</sub> | C <sub>24</sub> H <sub>25</sub> Br <sub>2</sub> HgOPdS <sub>2</sub> ·0.5 HgBr <sub>2</sub> ·C <sub>2</sub> H <sub>4</sub> Cl <sub>2</sub> |
| fw                                       | 2146.13                                                                                                     | 1348.72                                                                                                                                   |
| temp (K)                                 | 273(1)                                                                                                      | 143(1)                                                                                                                                    |
| wavelength (Å)                           |                                                                                                             | 0.710 73                                                                                                                                  |
| space group                              | P1̄                                                                                                         | P2 <sub>1</sub> /c                                                                                                                        |
| unit cell dimensions                     |                                                                                                             |                                                                                                                                           |
| <i>a</i> (Å)                             | 9.5755(11)                                                                                                  | 15.571(2)                                                                                                                                 |
| <i>b</i> (Å)                             | 11.1754(12)                                                                                                 | 10.7425(10)                                                                                                                               |
| <i>c</i> (Å)                             | 14.504(2)                                                                                                   | 19.655(2)                                                                                                                                 |
| α (deg)                                  | 84.826(5)                                                                                                   | 90                                                                                                                                        |
| β (deg)                                  | 81.611(7)                                                                                                   | 94.741(12)                                                                                                                                |
| γ (deg)                                  | 68.606(9)                                                                                                   | 90                                                                                                                                        |
| vol (Å <sup>3</sup> )                    | 1428.5(3)                                                                                                   | 3276.5(5)                                                                                                                                 |
| <i>Z</i>                                 | 1                                                                                                           | 4                                                                                                                                         |
| ρ <sub>calc</sub> (Mg/m <sup>3</sup> )   | 2.50                                                                                                        | 2.37                                                                                                                                      |
| abs coeff (mm <sup>-1</sup> )            | 12.64                                                                                                       | 11.58                                                                                                                                     |
| final <i>R</i> indices <sup>a</sup>      | <i>R</i> = 0.0351,<br>[ <i>I</i> > 2σ( <i>I</i> )] <i>R</i> <sub>w</sub> = 0.0371                           | <i>R</i> = 0.0410,<br><i>R</i> <sub>w</sub> = 0.0848                                                                                      |
| <i>R</i> indices <sup>a</sup> (all data) | <i>R</i> = 0.0839,<br><i>R</i> <sub>w</sub> = 0.0501                                                        | <i>R</i> = 0.0735,<br><i>R</i> <sub>w</sub> = 0.0964                                                                                      |

<sup>a</sup> *R* = Σ||*F*<sub>o</sub>| - |*F*<sub>c</sub>||/Σ|*F*<sub>o</sub>|; *R*<sub>w</sub> = [Σw(|*F*<sub>o</sub>| - |*F*<sub>c</sub>||)<sup>2</sup>/Σw|*F*<sub>o</sub>|<sup>2</sup>]<sup>1/2</sup>; *R*<sub>w</sub><sup>2</sup> = [Σ[w(*F*<sub>o</sub><sup>2</sup> - *F*<sub>c</sub><sup>2</sup>)/Σw(*F*<sub>o</sub><sup>2</sup>)]<sup>1/2</sup>.

solid (yield: 0.159 g, 76%). IR data (cm<sup>-1</sup>): C≡P, 458 (s), 478 (s), 501 (m), 521 (s), 534 (s), 560 (s), 580 (s), 755 (vs, sh), 1563 (m), 1589 (m); <sup>-</sup>S<sub>2</sub>COEt, 1021 (vs, br), 1234 (vs, br). <sup>1</sup>H NMR (218 K): C≡P, 3.38 (d, 1H), 3.76 (d, 1H, <sup>2</sup>J<sub>H-H</sub> = 10.8 Hz) (-CH<sub>2</sub>-Pd), 1.54 (s), 1.76 (s) (CH<sub>3</sub>-C<sub>6</sub>H<sub>4</sub>), 6.7–7.6 (m, 11H), 9.04 (d, 1H) (C<sub>6</sub>H<sub>4</sub>); <sup>-</sup>S<sub>2</sub>COEt, 4.65 (q), 1.42 (t) (<sup>3</sup>J<sub>H-H</sub> = 7.10 Hz). Anal. Calcd for Br<sub>2</sub>C<sub>24</sub>H<sub>25</sub>HgOPdS<sub>2</sub>: C, 32.34; H, 2.83. Found: C, 31.98; H, 2.81.

**Crystal Structure Determination of [Pt{CH<sub>2</sub>-C<sub>6</sub>H<sub>4</sub>-P(*o*-tolyl)<sub>2</sub>-κC,P}(S<sub>2</sub>CNMe<sub>2</sub>)HgI(μ-I)]<sub>2</sub> (**4**).** Important crystal data and data collection parameters for complex **4** are listed in Table 1. A parallelepiped-shaped crystal mounted on the tip of a glass fiber with epoxy cement was used for geometric and intensity data collection. All diffraction measurements were made at room temperature on a Siemens STOE/AED2 four circle diffractometer. Lattice dimensions and type were determined by routine procedures and verified by oscillation photography. Cell constants were refined from 2θ values of 70 reflections including Friedel pairs (17.3 < 2θ < 35.7°). Diffracted intensities were measured in a unique hemisphere of reciprocal space for 4.0 < 2θ < 50.0° by ω scans. During intensity data collection, three monitor reflections were measured at regular intervals, and they did not vary appreciably in intensity during the course of data collection. A measured absorption correction was applied on the basis of complete Ψ-scans of reflections with diffractometer angle χ near 90°. The structure of compound **4** was solved by direct methods and developed and refined in a series of alternating difference Fourier maps and least squares analyses using all data and the program package SHELXTL-PLUS.<sup>19a</sup> All non-hydrogen atoms were refined anisotropically. All hydrogen atoms were included in calculated positions and refined as riding atoms (C–H, 0.96 Å; *U* = 0.0917 Å<sup>2</sup>). Final difference electron density maps showed no peaks above 1 e Å<sup>-3</sup> (largest peak, 0.84 e Å<sup>-3</sup>; largest difference hole, -0.77 e Å<sup>-3</sup>). Residuals and other final refinement parameters are listed in Table 1. Residuals are defined in ref 19a.

**Crystal Structure Determination of [PdBr(S<sub>2</sub>COEt){μ-P(*o*-tolyl)<sub>2</sub>-C<sub>6</sub>H<sub>4</sub>-CH<sub>2</sub>-}HgBr]·0.5 HgBr<sub>2</sub>·C<sub>2</sub>H<sub>4</sub>Cl<sub>2</sub> (**11**·0.5 HgBr<sub>2</sub>·C<sub>2</sub>H<sub>4</sub>Cl<sub>2</sub>).** Crystal data and other details of the structure analysis are presented in Table 1. A crystal of compound **11**·0.5 HgBr<sub>2</sub>·C<sub>2</sub>H<sub>4</sub>Cl<sub>2</sub> was mounted at the end of a quartz fiber and glued in place with epoxy adhesive. All diffraction measurements were made at 143(1) K on an Enraf-Nonius

**Scheme 1**

CAD4 diffractometer, using graphite monochromated Mo K<sub>α</sub> X-radiation. Unit cell dimensions were determined from 25 centered reflections in the range 23.6 < 2θ < 31.8°. Diffracted intensities were measured in a unique quadrant of reciprocal space for 4.0 < 2θ < 50.0° by ω/θ scans. Three check reflections remeasured after every 3 h showed no decay of the crystal over the period of data collection. An absorption correction was applied on the basis of 555 azimuthal scan data (maximum and minimum transmission coefficients were 0.854 and 0.435). Lorentz and polarization corrections were applied. The structure was solved by Patterson and Fourier methods. All non-hydrogen atoms were assigned anisotropic displacement parameters and refined without positional constraints. The hydrogen atoms were constrained to idealized geometries and assigned isotropic displacement parameters 1.2 times the *U*<sub>iso</sub> value of their parent carbon atoms (1.5 times for the methyl hydrogen atoms). The carbon atoms of the 1,2-dichloroethane solvent molecule are disordered over two sets of positions (partial occupancy 0.5) and were refined with a common set of anisotropic thermal parameters. Full-matrix least squares refinement of this model against *F*<sup>2</sup> converged to final residuals given in Table 1. Final difference electron density maps showed three peaks above 1 e Å<sup>-3</sup> (1.24, 1.19, and 1.06 e Å<sup>-3</sup>; largest difference hole, -1.22) lying within 1.06 Å of the mercury atoms. Least squares calculations were carried out using the program SHELXL-93.<sup>19b</sup> Residuals are defined in ref 19b.

## Results and Discussion


**Reactivity of Complexes [Pt{CH<sub>2</sub>-C<sub>6</sub>H<sub>4</sub>-P(*o*-tolyl)<sub>2</sub>-κC,P}(S<sub>2</sub>CR)] [R = NMe<sub>2</sub> (**1**), OEt (**2**)] toward HgX<sub>2</sub> (X = Br, I).** The reactions of [Pt{CH<sub>2</sub>-C<sub>6</sub>H<sub>4</sub>-P(*o*-tolyl)<sub>2</sub>-κC,P}(S<sub>2</sub>CR)] (R = NMe<sub>2</sub> (**1**), OEt (**2**)) with HgX<sub>2</sub> (X = Br, I) (1:1 molar ratio) in dichloromethane afford the tetranuclear complexes [Pt{CH<sub>2</sub>-C<sub>6</sub>H<sub>4</sub>-P(*o*-tolyl)<sub>2</sub>-κC,P}(S<sub>2</sub>CR)HgX(μ-X)]<sub>2</sub> in high yields as air stable solids (Scheme 1a).

The IR spectra of complexes **3** and **4** show several absorptions corresponding to the Me<sub>2</sub>NCS<sub>2</sub><sup>-</sup> ligand in a bidentate coordination mode<sup>20</sup> (see Experimental Section). The values of the ν<sub>S-C</sub> (CN) [1554 cm<sup>-1</sup> (**3**), 1552 cm<sup>-1</sup> (**4**)] suggest a pronounced degree of C–N double bonding, even more so than in the starting complex (1543 cm<sup>-1</sup>). The IR spectra of complexes **5** and **6** also show the typical absorptions of the EtOCS<sub>2</sub><sup>-</sup> group.<sup>20</sup>

Figure 1 shows the molecular structure of compound **4** with the atom numbering scheme. General crystallographic information is collected in Table 1. Relevant bond distances and angles are listed in Table 2.

(19) (a) SHELXTL-PLUS Software Package for the Determination of Crystal Structures, Release 4.0; Siemens Analytical X-ray Instruments, Inc.: Madison, WI, 1990. (b) Sheldrick, G. M. SHELXL-93, a program for crystal structure determination; University of Göttingen: Göttingen, Germany, 1993.

(20) Coucouvanis, D. *Progress in Inorganic Chemistry*; John Wiley & Sons: New York, 1979; Vol. 26, p 424.



**Figure 1.** Drawing of the crystal structure of  $[\text{Pt}\{\text{CH}_2\text{-C}_6\text{H}_4\text{P}(\text{o-tolyl})_2\text{-}\kappa\text{C,P}\}(\text{S}_2\text{CNMe}_2)\text{HgI}(\mu\text{-I})]_2$  (**4**), showing the atom labeling scheme. Atoms are represented by their 50% probability ellipsoids.

**Table 2.** Selected Bond Lengths (Å) and Angles (deg) for  $[\text{Pt}\{\text{CH}_2\text{-C}_6\text{H}_4\text{P}(\text{o-tolyl})_2\text{-}\kappa\text{C,P}\}(\text{S}_2\text{CNMe}_2)\text{HgI}(\mu\text{-I})]_2$  (**4**)

|               |           |               |           |
|---------------|-----------|---------------|-----------|
| Pt—Hg         | 2.768(1)  | Pt—C(21)      | 2.081(14) |
| Pt—P          | 2.246(4)  | Pt—S(1)       | 2.413(4)  |
| Pt—S(2)       | 2.341(5)  | C(22)—N       | 1.315(21) |
| C(22)—S(1)    | 1.707(14) | C(22)—S(2)    | 1.733(14) |
| Hg—I(1)       | 2.661(1)  | Hg—I(2)       | 2.766(2)  |
| Hg—I(2')      | 3.041(1)  |               |           |
| S(1)—Pt—S(2)  | 73.9(1)   | C(22)—N—C(24) | 122.1(13) |
| S(1)—Pt—P     | 107.5(1)  | Pt—Hg—I(1)    | 113.8(1)  |
| P—Pt—C(21)    | 84.6(4)   | Pt—Hg—I(2)    | 112.4(1)  |
| S(2)—Pt—C(21) | 94.0(4)   | Pt—Hg—I(2')   | 109.5(1)  |
| S(1)—C(22)—N  | 125.5(1)  | I(1)—Hg—I(2)  | 122.7(1)  |
| S(2)—C(22)—N  | 122.2(10) | I(1)—Hg—I(2') | 105.0(1)  |
| C(22)—N—C(23) | 121.9(13) | I(2)—Hg—I(2') | 89.5(1)   |
| Hg—I(2)—Hg'   | 90.5(1)   |               |           |

The tetranuclear complex **4** can be regarded as being formed by two “ $\text{Pt}\{\text{CH}_2\text{-C}_6\text{H}_4\text{P}(\text{o-tolyl})_2\text{-}\kappa\text{C,P}\}(\text{S}_2\text{CNMe}_2)\text{HgI}_2$ ” units bridged by two I atoms and related to each other by a center of symmetry.

As can be seen, each unit contains one unsupported Pt—Hg bond. In each unit the five-coordinated Pt atom is located at the center of the base of a square pyramid with the Hg atom in the apical position. The angle between the Pt—Hg vector and the perpendicular to the Pt basal plane [Pt, C(21), P, S(1), S(2)] is  $9.19^\circ$ .<sup>21</sup>

Bond distances and angles corresponding to the “ $\text{Pt}\{\text{CH}_2\text{-C}_6\text{H}_4\text{P}(\text{o-tolyl})_2\text{-}\kappa\text{C,P}\}(\text{S}_2\text{CNMe}_2)$ ” fragment are similar to those observed in the Pt(II) complex which is used as starting material.<sup>16</sup>

The Pt—Hg distance [2.768(1) Å] and the geometry around the Pt center are similar to those observed in  $[\{2,6\text{-(Me}_2\text{NCH}_2\text{)}_2\text{-C}_6\text{H}_3\}\text{Pt}(\mu\text{-}\{p\text{-tol}\}\text{NC(H)N}(\text{Pr})\}\text{HgBrCl}]$  [2.8331(7) Å],<sup>22</sup>  $\text{trans-}[(\text{CH}_3\text{NH}_2)_2\text{Pt}(1,5\text{-diMeC}^-)_2\text{Hg}](\text{NO}_3)_2 \cdot 0.5 \text{H}_2\text{O}$  [2.765(1) Å],<sup>23</sup>  $\text{trans-}[(\text{CH}_3\text{NH}_2)_2\text{Pt}(1\text{-MeC}^-)_2\text{HgCl}(\text{NO}_3)]$  [2.835(1) Å],<sup>23</sup> and  $\text{trans-}[(\text{CH}_3\text{NH}_2)_2\text{Pt}(1\text{-MeC}^-)_2\text{Hg}](\text{NO}_3)_2$  [2.785(1) Å],<sup>23</sup> for which the Pt—Hg interaction has been described as a Pt→Hg donor bond with both metals in a formal oxidation state of 2. In contrast, the Pt—Hg distance is clearly different from the distances observed in mixed Pt—Hg compounds displaying a

covalent bond:  $[(2\text{-Me}_2\text{NCH}_2\text{-C}_6\text{H}_4)_2(\mu\text{-MeCO}_2)\text{PtHg}(\text{O}_2\text{CMe})]$ , 2.513(1) Å;<sup>24</sup>  $[(\text{PPh}_3)_2\text{R-Pt-Hg-R}]$ , 2.637(1) Å;<sup>25a</sup>  $[\text{PhCH}[\text{HgPt}(\text{Ph}_3\text{P})_2\text{Br}]\text{COOC}_{10}\text{H}_{19}]$ , 2.499 Å;<sup>25b</sup>  $[\text{N}(\text{CH}_2\text{CH}_2\text{PPh}_2)_3\text{Pt}(\text{HgMe})]\text{BPh}_4$ , 2.531(1) Å;<sup>25c</sup>  $[\text{PtCl}(\text{HgMe})(\text{dmphen})(\text{Z-MeO}_2\text{-CCH=CHCO}_2\text{Me})]$ , 2.558(1) Å.<sup>25d</sup>

It is noteworthy that, in complex **4**, the Pt→Hg bond is not supported by any bridging ligand, especially considering that the basic Pt complex  $[\text{Pt}\{\text{CH}_2\text{-C}_6\text{H}_4\text{P}(\text{o-tolyl})_2\text{-}\kappa\text{C,P}\}(\text{S}_2\text{CNMe}_2)]$  used as starting material contains two S atoms with pairs of electrons capable of forming a Hg—S bond. Mercury, furthermore, has a well-known affinity for sulfur.<sup>26</sup> The absence of bridging ligands between the Pt and Hg centers makes compound **4** different from the other complexes exhibiting Pt<sup>II</sup> to Hg<sup>II</sup> donor bonds.<sup>22,23</sup>

From the point of view of the Hg(II), the complex can be considered as a dimer of stoichiometry  $[\text{HgIL}(\mu\text{-I})]_2$ , L being the complex  $[\text{Pt}\{\text{CH}_2\text{-C}_6\text{H}_4\text{P}(\text{o-tolyl})_2\text{-}\kappa\text{C,P}\}(\text{S}_2\text{CNMe}_2)]$  with the Pt acting as the donor atom. Each Hg atom is located in a very distorted tetrahedral environment, the Hg—I terminal distance [2.661(1) Å] being shorter than the Hg—I bridging ones, which are in turn very different from each other [Hg—I(2), 2.766(2) Å, Hg—I(2'), 3.041(1) Å], in keeping with the structures of other  $[\text{LHgX}(\mu\text{-X})]_2$  complexes.<sup>27,28</sup> The Hg—I(2)—Hg—I(2') fragment is planar and the angles I(2)—Hg—I(2') and Hg—I(2)—Hg' are nearly  $90^\circ$ .

Although the literature contains numerous reports of 1:1 complexes of mercuric halides with neutral ligands such as  $[\text{HgX}(\text{Me}_3\text{SiCH}_2\text{SeMe})(\mu\text{-X})]_2$  (X = Cl, Br, I),<sup>27a</sup>  $[\text{HgI}(\text{Ph}_3\text{PC}_5\text{H}_4)(\mu\text{-I})]_2$ ,<sup>27b</sup>  $[\text{HgCl}(\text{PBu}_3)(\mu\text{-Cl})]_2$ ,<sup>28a</sup> or  $[\text{Ph}_3\text{PCHCOC}_6\text{H}_5\text{-HgX}_2]_2$  (X = Cl, I)<sup>28d</sup> showing discrete centrosymmetric halogen bridged dimeric structures, to our knowledge, there is none in which the neutral ligand is a metal complex with the metal center acting as donor atom, as in this case.

In agreement with the molecular symmetry observed in the crystal, the  $^3\text{P}\{^1\text{H}\}$  NMR spectra of complexes **3–6** (see Table 3) show only one singlet at about 25 ppm with the corresponding <sup>195</sup>Pt satellites. The P—Pt coupling constants in these complexes are smaller than those of the starting materials (see Table 3) in the range of about 200–400 Hz.

The <sup>1</sup>H NMR spectra of complexes **3–6** in CDCl $_3$  at room temperature (20 °C) also confirm the equivalence of the two halves of the molecule in each case. These spectra show a common pattern due to the cyclometalated group “ $\text{Pt}\{\text{CH}_2\text{-C}_6\text{H}_4\text{P}(\text{o-tolyl})_2\text{-}\kappa\text{C,P}\}$ ” (see Experimental Section): (a) The CH $_3$  groups of the o-tolylphosphine appear as two singlets indicating

(21) Nardelli, M. *Comput. Chem.* **1983**, 7, 95.

(22) van der Ploeg, A. F. M. J.; van Koten, G.; Vrieze, K.; Spek, A. L.; Duisenberg, A. J. M. *Organometallics* **1982**, 1, 1066.

(23) Krumm, M.; Zangrando, E.; Randaccio, L.; Menzer, S.; Danzmann, A.; Holtherrich, D.; Lippert, B. *Inorg. Chem.* **1993**, 32, 2183.

(24) (a) van der Ploeg, A. F. M. J.; van Koten, G.; Vrieze, K.; Spek, A. L.; Duisenberg, A. J. M. *J. Chem. Soc., Chem. Commun.* **1980**, 469. (b) van der Ploeg, A. F. M. J.; van Koten, G.; Vrieze, K.; Spek, A. L. *Inorg. Chem.* **1982**, 21, 2014.

(25) (a) Rossell, O.; Seco, M.; Torra, I.; Solans, X.; Font-Altaba, M. *J. Organomet. Chem.* **1984**, 270, C63. (b) Suleimanov, G. Z.; Bashilov, V. V.; Musaev, A. A.; Sokolov, V. I.; Reutov, O. A. *J. Organomet. Chem.* **1980**, 202, C61. (c) Ghilardi, C. A.; Midollini, S.; Moneti, S.; Orlandini, A.; Scapacci, G.; Daktariens, D. *J. Chem. Soc., Chem. Commun.* **1989**, 1686. (d) Cucciolito, E.; Giordano, F.; Panunzi, A.; Rufo, F.; de Felice, V. *J. Chem. Soc., Dalton Trans.* **1993**, 3421.

(26) (a) Christou, G.; Folting, K.; Huffman, J. C. *Polyhedron* **1984**, 3, 1247. (b) Bond, A. M.; Colton, R.; Hollenkamp, A. F.; Hoskins, B. F.; McGregor, K. *J. Am. Chem. Soc.* **1987**, 109, 1969.


(27) (a) Chadha, R. K.; Drake, J. E.; McManus, N. T.; Mislankar, A. *Can. J. Chem.* **1987**, 65, 2305. (b) Baenziger, N. C.; Flynn, R. M.; Swenson, D. C. *Acta Crystallogr.* **1978**, B34, 2300.

(28) (a) Bell, N. A.; Goldstein, M.; Jones, T.; Nowell, I. W. *Inorg. Chim. Acta* **1980**, 43, 87. (b) van Enckevort, W. J. P.; Beurskens, P. T.; Menger, E. M.; Bosman, W. P. *Cryst. Struct. Commun.* **1977**, 6, 417. (c) Castineiras, A.; Arquer, A.; Masaguer, J. R.; Martínez-Carrera, S.; García Blanco, S. *Anorg. Allg. Chem.* **1986**, 539, 219. (d) Kalyanasundari, M.; Panchanatheswaran, K.; Robinson, W. T.; Wen, H. *J. Organomet. Chem.* **1995**, 491, 103.

their inequivalence; (b) the signals corresponding to the CH<sub>2</sub>–Pt group are isochronous, showing however different values of  $^2J_{\text{Pt-H}}$ ; (c) the aromatic hydrogens give several multiplets between 6.5 and 7.8 ppm. The main difference observed with respect to the starting complex is the  $\delta$  value of the CH<sub>2</sub>–Pt signal together with the Pt–H coupling constant.

Square-pyramidal compounds containing Pt<sup>II</sup>→Hg<sup>II</sup> donor bonds (type II) have been proposed as intermediates in the formation of species with covalent Pt–Hg bonds (type I).<sup>24</sup> However, we have not observed the subsequent formation of type I complexes from compounds **3–6** (type II).



The reactions of compounds **3–6** with PPh<sub>3</sub> in 1:2 molar ratio afford the starting Pt(II) complexes [Pt{CH<sub>2</sub>–C<sub>6</sub>H<sub>4</sub>–P(*o*-tolyl)<sub>2</sub>– $\kappa$ C,P}(S<sub>2</sub>CR)] [R = NMe<sub>2</sub> (**1**), OEt (**2**)] as a result of the cleavage of the Pt<sup>II</sup>→Hg<sup>II</sup> bond rather than of the dihalogen bridging system (eq 1).



The reactivity of the platinum(II) complexes [Pt{CH<sub>2</sub>–C<sub>6</sub>H<sub>4</sub>–P(*o*-tolyl)<sub>2</sub>– $\kappa$ C,P}(S<sub>2</sub>CR)] [R = NMe<sub>2</sub> (**1**), OEt (**2**)] toward HgX<sub>2</sub> prompted us to study the reactions of the analogous palladium(II) complexes with the aim of preparing the corresponding derivatives with Pd→Hg bonds.

**Synthesis and Characterization of [Pd{CH<sub>2</sub>–C<sub>6</sub>H<sub>4</sub>–P(*o*-tolyl)<sub>2</sub>– $\kappa$ C,P}(S<sub>2</sub>CR)] (R = NMe<sub>2</sub> (**7**), OEt (**8**)).** These compounds have been prepared similarly to the previously reported platinum analogues.<sup>16</sup>

Solutions resulting from the reaction of [Pd{CH<sub>2</sub>–C<sub>6</sub>H<sub>4</sub>–P(*o*-tolyl)<sub>2</sub>– $\kappa$ C,P}(μ-Cl)]<sub>2</sub> with AgClO<sub>4</sub> (1:2 molar ratio) in THF at room temperature, presumably containing the [Pd{CH<sub>2</sub>–C<sub>6</sub>H<sub>4</sub>–P(*o*-tolyl)<sub>2</sub>– $\kappa$ C,P}(THF)<sub>2</sub>]ClO<sub>4</sub> species, react with equimolar amounts of sodium dimethyldithiocarbamate (NaS<sub>2</sub>CNMe<sub>2</sub>·2H<sub>2</sub>O) and potassium ethylxanthate (KS<sub>2</sub>COEt) to give the mononuclear complexes [Pd{CH<sub>2</sub>–C<sub>6</sub>H<sub>4</sub>–P(*o*-tolyl)<sub>2</sub>– $\kappa$ C,P}(S<sub>2</sub>CNMe<sub>2</sub>)] (**7**) and [Pd{CH<sub>2</sub>–C<sub>6</sub>H<sub>4</sub>–P(*o*-tolyl)<sub>2</sub>– $\kappa$ C,P}(S<sub>2</sub>COEt)] (**8**) as air stable solids (see Experimental Section; eqs 2 and 3).



The similarities of the IR and <sup>1</sup>H NMR spectra of compounds **7** and **8** to those of the analogous Pt complexes [Pt{CH<sub>2</sub>–C<sub>6</sub>H<sub>4</sub>–P(*o*-tolyl)<sub>2</sub>– $\kappa$ C,P}(S<sub>2</sub>CR)] [R = NMe<sub>2</sub> (**1**), OEt (**2**)] suggest a similar structure for all of them, consisting of square-planar mononuclear complexes with bidentate chelating ligands. As in the Pt derivatives **1** and **2**, the hindered rotation of the P–C<sub>ipso</sub> bonds at room temperature causes the inequivalence of the two CH<sub>3</sub> groups of the *o*-tolylphosphine. The  $\Delta G^\ddagger$  values at the coalescence temperature of the methyl resonances, for the P–C<sub>ipso</sub> bond rotation process in complexes **7** ( $\Delta G^\ddagger_{319\text{K}} = 64.2$  KJ mol<sup>–1</sup>) and **8** ( $\Delta G^\ddagger_{313\text{K}} = 63.2$  KJ mol<sup>–1</sup>) have been calculated using the approximation to Eyring's equation,<sup>29</sup> and they are very similar to the values found for the analogous platinum derivatives **1** and **2**.

**Table 3.** <sup>31</sup>P{<sup>1</sup>H} NMR<sup>a</sup> Data for Complexes **1–11**

| complex                                                                                   | δP (ppm)  | <sup>1</sup> J <sub>Pt–P</sub> (Hz) | Δ <sup>1</sup> J <sub>Pt–P</sub> (Hz) |
|-------------------------------------------------------------------------------------------|-----------|-------------------------------------|---------------------------------------|
| [Pt(C $\wedge$ P)(S <sub>2</sub> CNMe <sub>2</sub> )] ( <b>1</b> )                        | 24.70 (s) | 3969.4                              |                                       |
| [Pt(C $\wedge$ P)(S <sub>2</sub> COEt)] ( <b>2</b> )                                      | 23.73 (s) | 4083.8                              |                                       |
| [Pt(C $\wedge$ P)(S <sub>2</sub> CNMe <sub>2</sub> )HgBr(μ-Br)] <sub>2</sub> ( <b>3</b> ) | 26.06 (s) | 3594.93                             | 374.5 <sup>c</sup>                    |
| [Pt(C $\wedge$ P)(S <sub>2</sub> CNMe <sub>2</sub> )HgI(μ-I)] <sub>2</sub> ( <b>4</b> )   | 25.69 (s) | 3681.10                             | 288.3 <sup>c</sup>                    |
| [Pt(C $\wedge$ P)(S <sub>2</sub> COEt)HgBr(μ-Br)] <sub>2</sub> ( <b>5</b> )               | 25.16 (s) | 3745.80                             | 338.0 <sup>d</sup>                    |
| [Pt(C $\wedge$ P)(S <sub>2</sub> COEt)HgI(μ-I)] <sub>2</sub> ( <b>6</b> )                 | 24.63 (s) | 3862.75                             | 221.0 <sup>d</sup>                    |
| [Pd(C $\wedge$ P)(S <sub>2</sub> CNMe <sub>2</sub> )] ( <b>7</b> )                        | 35.58 (s) |                                     |                                       |
| [Pd(C $\wedge$ P)(S <sub>2</sub> COEt)] ( <b>8</b> )                                      | 35.93 (s) |                                     |                                       |
| [Pd(C $\wedge$ P)(S <sub>2</sub> CNMe <sub>2</sub> )HgI(μ-I)] <sub>2</sub> ( <b>9</b> )   | 36.76 (s) |                                     |                                       |
| [PdBr(S <sub>2</sub> CNMe <sub>2</sub> )(μ-P $\wedge$ C)HgBr] ( <b>10</b> ) <sup>b</sup>  | 20.24 (s) |                                     |                                       |
| [PdBr(S <sub>2</sub> COEt)(μ-P $\wedge$ C)HgBr] ( <b>11</b> ) <sup>b</sup>                | 20.71 (s) |                                     |                                       |

<sup>a</sup> C $\wedge$ P, CH<sub>2</sub>–C<sub>6</sub>H<sub>4</sub>–P(*o*-tolyl)<sub>2</sub>– $\kappa$ C,P; external reference H<sub>3</sub>PO<sub>4</sub>, 85%; CDCl<sub>3</sub>; room temperature. <sup>b</sup> 218 K. <sup>c</sup> Δ<sup>1</sup>J<sub>Pt–P</sub> = <sup>1</sup>J<sub>Pt–P</sub>(1) – <sup>1</sup>J<sub>Pt–P</sub>. <sup>d</sup> Δ<sup>1</sup>J<sub>Pt–P</sub> = <sup>1</sup>J<sub>Pt–P</sub>(2) – <sup>1</sup>J<sub>Pt–P</sub>.

**Reactivity of [Pd{CH<sub>2</sub>–C<sub>6</sub>H<sub>4</sub>–P(*o*-tolyl)<sub>2</sub>– $\kappa$ C,P}(S<sub>2</sub>CR)] [R = NMe<sub>2</sub> (**7**), OEt (**8**)] toward HgX<sub>2</sub> (X = Br, I).** The reactions of [Pd{CH<sub>2</sub>–C<sub>6</sub>H<sub>4</sub>–P(*o*-tolyl)<sub>2</sub>– $\kappa$ C,P}(S<sub>2</sub>CR)] (R = NMe<sub>2</sub> (**7**), OEt (**8**)) with HgX<sub>2</sub> (X = Br, I) proceed differently depending on the nature of X and the S<sub>2</sub>CR<sup>–</sup> ligands.

[Pd{CH<sub>2</sub>–C<sub>6</sub>H<sub>4</sub>–P(*o*-tolyl)<sub>2</sub>– $\kappa$ C,P}(S<sub>2</sub>CNMe<sub>2</sub>)] (**7**) reacts with HgI<sub>2</sub> at room temperature in a 1:1 molar ratio to give [Pd{CH<sub>2</sub>–C<sub>6</sub>H<sub>4</sub>–P(*o*-tolyl)<sub>2</sub>– $\kappa$ C,P}(S<sub>2</sub>CNMe<sub>2</sub>)HgI(μ-I)]<sub>2</sub> (**9**) in a very high yield (Scheme 1b), whereas the xanthate complex [Pd{CH<sub>2</sub>–C<sub>6</sub>H<sub>4</sub>–P(*o*-tolyl)<sub>2</sub>– $\kappa$ C,P}(S<sub>2</sub>COEt)] (**8**) does not react with HgI<sub>2</sub> under similar conditions or even when the reactants are refluxed in diethyl ether for several hours (Scheme 1c). The structure of **9** has been assigned on the bases of the similarities of its IR and <sup>31</sup>P{<sup>1</sup>H} and <sup>1</sup>H NMR spectra with its platinum-containing analogues, [Pt{CH<sub>2</sub>–C<sub>6</sub>H<sub>4</sub>–P(*o*-tolyl)<sub>2</sub>– $\kappa$ C,P}(S<sub>2</sub>CNMe<sub>2</sub>)HgBr(μ-Br)]<sub>2</sub> (**3**) and [Pt{CH<sub>2</sub>–C<sub>6</sub>H<sub>4</sub>–P(*o*-tolyl)<sub>2</sub>– $\kappa$ C,P}(S<sub>2</sub>CNMe<sub>2</sub>)HgI(μ-I)]<sub>2</sub> (**4**).


HgBr<sub>2</sub> reacts with both complexes **7** and **8**, giving similar complexes although different from the one obtained in the reaction with HgI<sub>2</sub>. Treatment of dichloromethane solutions of [Pd{CH<sub>2</sub>–C<sub>6</sub>H<sub>4</sub>–P(*o*-tolyl)<sub>2</sub>– $\kappa$ C,P}(S<sub>2</sub>CR)] [R = NMe<sub>2</sub> (**7**), OEt (**8**)] with HgBr<sub>2</sub> at low temperature (–25 °C) affords the air stable solids [PdBr(S<sub>2</sub>CNMe<sub>2</sub>){μ-P(*o*-tolyl)<sub>2</sub>–C<sub>6</sub>H<sub>4</sub>–CH<sub>2</sub>–}HgBr] (**10**) and [PdBr(S<sub>2</sub>COEt){μ-P(*o*-tolyl)<sub>2</sub>–C<sub>6</sub>H<sub>4</sub>–CH<sub>2</sub>–}HgBr] (**11**) in high yields (Scheme 1d). As can be seen in Figure 2, compounds **10** and **11** are the result of transmetalation processes in which Pd–Br and C–Hg bonds are formed. These reactions have to be performed at low temperature [–25 °C and –30 °C] in order to avoid the formation of an unidentified byproduct which appears as an impurity (δP = 42 ppm) in products **10** and **11** when the reactions are carried out at room temperature.

The geometry of the binuclear palladium–mercury complex [PdBr(S<sub>2</sub>COEt){μ-P(*o*-tolyl)<sub>2</sub>–C<sub>6</sub>H<sub>4</sub>–CH<sub>2</sub>–}HgBr] (**11**), as determined by X-ray diffraction studies, is illustrated in Figure 2 together with the atomic numbering scheme. General crystallographic information is collected in Table 1. Relevant bond distances and angles are listed in Table 4.

As can be seen, the molecule contains one palladium and one mercury atom bridged by the bidentate “P(*o*-tolyl)<sub>2</sub>–C<sub>6</sub>H<sub>4</sub>–CH<sub>2</sub>–” group, the Pd–Hg distance being 3.098(1) Å.

The palladium atom is located in a distorted square-planar environment formed by one bromine, two sulfur atoms of the ethylxanthate ligand, and the P of the bidentate bridging group; the S(1) atom is 0.159 Å away from the best plane<sup>21</sup> defined by Pd, Br(1), S(1), S(2), and P. The angles around the palladium center between *cis* ligand bonds range from 74.9(1) to 97.6(1)°, distortion which can be mainly due to the small bite angle

(29) Günther, H. *NMR Spectroscopy: An Introduction*; John Wiley & Sons: Chichester, U.K., 1980.



**Figure 2.** Drawing of the crystal structure of [PdBr(S<sub>2</sub>COEt){μ-P(o-tolyl)<sub>2</sub>-C<sub>6</sub>H<sub>4</sub>-CH<sub>2</sub>}HgBr] (**11**), showing the atom labeling scheme. Atoms are represented by their 50% probability ellipsoids.

**Table 4.** Selected Bond Lengths (Å) and Angles (deg) for [PdBr(S<sub>2</sub>COEt){μ-P(o-tolyl)<sub>2</sub>-C<sub>6</sub>H<sub>4</sub>-CH<sub>2</sub>}HgBr]·0.5 HgBr<sub>2</sub>·C<sub>2</sub>H<sub>4</sub>Cl<sub>2</sub> (**11**·0.5HgBr<sub>2</sub>·C<sub>2</sub>H<sub>4</sub>Cl<sub>2</sub>)


|                  |           |                   |           |
|------------------|-----------|-------------------|-----------|
| Pd—Br(1)         | 2.454(1)  | Pd—P              | 2.305(2)  |
| Pd—S(1)          | 2.288(3)  | Pd—S(2)           | 2.389(2)  |
| Pd—Hg            | 3.098(1)  | Hg—Br(2)          | 2.460(1)  |
| Hg—C(10)         | 2.106(10) | S(1)—C(1)         | 1.693(10) |
| S(2)—C(1)        | 1.706(10) | O—C(1)            | 1.299(11) |
| O—C(2)           | 1.489(14) | C(2)—C(3)         | 1.35(2)   |
| S(1)—Pd—P        | 94.93(9)  | S(1)—Pd—S(2)      | 74.90(9)  |
| Br(1)—Pd—P       | 92.58(7)  | S(2)—Pd—Br(1)     | 97.62(7)  |
| C(5)—C(10)—Hg(1) | 109.5(6)  | C(10)—Hg(1)—Br(2) | 171.1(3)  |
| S(1)—C(1)—S(2)   | 113.7(5)  | S(1)—C(1)—O       | 125.8(8)  |
| S(2)—C(1)—O      | 120.5(7)  | C(1)—O—C(2)       | 116.5(9)  |

of the chelate xanthate ligand (74.1°).<sup>30</sup> Pd—S distances are different, with Pd—S(2) [2.389(2) Å] longer than Pd—S(1) [2.288(3) Å], probably because of the higher *trans* influence of the P ligand compared to the bromide. The Pd—P distance [2.305(2) Å] is slightly longer than that observed in the complexes [Pd{CH<sub>2</sub>-C<sub>6</sub>H<sub>4</sub>-P(o-tolyl)<sub>2</sub>-κC,P}(μ-OOCCH<sub>3</sub>)<sub>2</sub>]<sup>18</sup> and [Pt{CH<sub>2</sub>-C<sub>6</sub>H<sub>4</sub>-P(o-tolyl)<sub>2</sub>-κC,P}(S<sub>2</sub>CNMe<sub>2</sub>)]<sup>16</sup> in which the CAP ligand acts as a chelate.

The xanthate ligand is essentially planar and almost coplanar with the Pd coordination plane (dihedral angle 6.3°). The C—S bond lengths [1.693(10), 1.706(10) Å] are very similar to those observed in other complexes with chelating xanthate ligands,<sup>31</sup> but the C—O distance [1.489(14) Å] is longer than those normally observed in this type of complexes and is as long as a typical C—O single bond.<sup>31</sup>

The Hg atom is located at 3.098(1) Å from the Pd atom and is close to the perpendicular to the Pd coordination plane; the angle between the perpendicular to the palladium coordination plane and the Pd—Hg vector is 17.5°. The Hg is coordinated to a Br and to the C donor atom of the bridging “P(o-tolyl)<sub>2</sub>-C<sub>6</sub>H<sub>4</sub>-CH<sub>2</sub>—” group in almost a linear mode [C(10)—Hg—Br(2) = 171.1(3)°] as is usual for RHgX compounds.<sup>32</sup> The Hg—Br<sup>5,27b</sup> and the Hg—C<sup>3,25</sup> distances are similar to those found in other organometallic Hg complexes.

The compound [PdBr(S<sub>2</sub>COEt){μ-P(o-tolyl)<sub>2</sub>-C<sub>6</sub>H<sub>4</sub>-CH<sub>2</sub>}-HgBr] (**11**) crystallizes with 0.5 molecules of HgBr<sub>2</sub>. The presence of HgBr<sub>2</sub> in the crystal can be explained in terms of the equilibrium represented in eq 4. The fact that complexes **10** and **11** dissociate in solution according to eq 4 is confirmed by the <sup>31</sup>P{<sup>1</sup>H} NMR data.




The <sup>31</sup>P{<sup>1</sup>H} NMR spectra of [PdBr(S<sub>2</sub>CR){μ-P(o-tolyl)<sub>2</sub>-C<sub>6</sub>H<sub>4</sub>-CH<sub>2</sub>}HgBr] [R = NMe<sub>2</sub> (**10**), OEt (**11**)] at room temperature show in both cases two signals, one due to complex **10** or **11**, respectively, and the other corresponding to the starting material **7** or **8**, respectively.

The spectrum of complex **10** at −55 °C does not show the signal corresponding to the starting material, while the spectrum of complex **11** at −55 °C shows the two signals observed at room temperature. The disappearance of the signal due to the starting material (at −55 °C) requires that HgBr<sub>2</sub> be added to the above mentioned solution of **11**. All these facts suggest that complexes **10** and **11** dissociate in solution at room temperature (**10**) or even low temperature (**11**) (eq 4).

As can be seen from the <sup>31</sup>P{<sup>1</sup>H} NMR data (Table 3), the new disposition of the CAP group, which now acts as a bridge between the Pd and Hg atoms, leads to an important change in the position of the <sup>31</sup>P signal (about 15 ppm) with respect to the cases in which this ligand acts as a chelate at the Pd center (complexes **7**–**9**). The new arrangement also causes important changes in the <sup>1</sup>H NMR spectrum, especially in the signals corresponding to the —CH<sub>2</sub>— group, which appears as an AX system with a <sup>2</sup>J<sub>H—H</sub> close to 10 Hz, and also for those of CH<sub>3</sub>—C<sub>6</sub>H<sub>4</sub>, which are shifted to lower δ by almost 1 ppm.

The formation of complexes **10** and **11** takes place as a result of a transmetalation process. Cleavage of platinum carbon σ bonds by electrophiles such as HgCl<sub>2</sub> have been observed previously.<sup>14b,c</sup> These reactions result in alkyl or aryl transfer from the platinum substrate to mercury and the formation of organomercurials (eq 5).



In our case, although the cleavage of the Pd—C σ bond takes place, the bidentate nature of the C,P-cyclometalated ligand leads to the formation of Pd—Hg complexes with the ligand acting as a bridge between the two metal centers.

Dirhodium(II) compounds with metalated arylphosphines acting as bridges between the two metal centers have been reported, with the phosphines forming three-atom bridges.<sup>33</sup> The formation of a four-atom bridge between two metal centers is very rare due to the stability of the five-membered metallocycles, and as far as we know only one such compound —[(C<sub>6</sub>F<sub>5</sub>)(8-mq)Pd(μ-8-mq)Pd(C<sub>6</sub>F<sub>5</sub>)(phen)] (8-mq = 8-quinolylmethyl)—containing a C,N metalated ligand (8-mq) with a four-atom bridge has previously been reported.<sup>34</sup>

The present results show the greater capacity of Pt(II), as compared to Pd(II), for the formation of donor–acceptor metal–metal bonds, as well as the predisposition of Hg to bond to carbon.

**Acknowledgment.** We thank the Dirección General de Enseñanza Superior (Spain) for financial support (Projects PB95-0003-CO2-01 and PB95-0792).

**Supporting Information Available:** For the crystal structure of **4**, full tables of crystallographic data, atomic coordinates, thermal parameters, and bond lengths and angles (5 pages). X-ray crystallographic files, in CIF format, for complex **11**·0.5HgBr<sub>2</sub>·C<sub>2</sub>H<sub>4</sub>Cl<sub>2</sub> are available on the Internet only. Ordering and access information is given on any current masthead page.

IC970584E

(30) Usón, R.; Forniés, J.; Falvello, L. R.; Tomás, M.; Ara, I.; Usón, I. *Inorg. Chim. Acta* **1995**, 232, 35.

(31) Orpen, G.; Brammer, L.; Allen, F. H.; Kennard, O.; Watson, D. G.; Taylor, R. *J. Chem. Soc., Dalton Trans.* **1989**, S1.

(32) Cotton, F. A.; Wilkinson, G. *Advanced Inorganic Chemistry*, 5th ed.; John Wiley & Sons: New York, 1988; p 618.

(33) Estevan, F.; González, G.; Lahuerta, P.; Martínez, M.; Peris, E.; van Eldik, R. *J. Chem. Soc., Dalton Trans.* **1996**, 1045.

(34) Forniés, J.; Navarro, R.; Sicilia, V.; Tomás, M. *Organometallics* **1990**, 9, 2422.