

Reactions of Carbohydrate α -Keto Toluene-*p*-sulphonates. Reaction of Methyl 4,6-*O*-Benzylidene-2-*O*-toluene-*p*-sulphonyl- α -D-*ribo*-hexopyranosid-3-ulose with Triethylamine-Methanol


By A. DMYTRACZENKO, W. A. SZAREK,* and J. K. N. JONES

(Department of Chemistry, Queen's University, Kingston, Ontario, Canada)

Summary. Reaction of methyl 4,6-*O*-benzylidene-2-*O*-toluene-*p*-sulphonyl- α -D-*ribo*-hexopyranosid-3-ulose (I) with triethylamine-methanol yields methyl 2,3-anhydro-4,6-*O*-benzylidene-3-methoxy- α -D-allopyranoside (II), methyl 4,6-*O*-benzylidene- α -D-*ribo*-hexopyranosid-3-ulose dimethyl acetal (III; R=H), and a derivative of 3-hydroxy-2-(hydroxymethyl)-4*H*-pyran-4-one (IV).

THE importance of sulphonic esters of carbohydrates as substrates for the preparation of a wide variety of sugars by nucleophilic displacement reactions has been firmly established.¹ The reactivity of a sulphonate group depends

critically upon the position of the group and upon the stereochemistry and conformation of the carbohydrate molecule.^{1,2} If a sulphonate is situated at C-2 of a pyranoside, then nucleophilic displacement with charged nucleophiles does not normally occur. Attempts have been made³ to achieve nucleophilic displacements of a 2-sulphonate by using a more activated sulphonate as a leaving group as in methyl 4,6-*O*-benzylidene-2-*O*-toluene-*p*-sulphonyl- α -*ribo*-hexopyranosid-3-ulose (I); it is known that a carbonyl group adjacent to the reaction centre assists bimolecular substitution. Here we report the unusual compounds obtained from the reaction of the α -keto toluene-*p*-sulphonate^{3,4} (I) with triethylamine-methanol.




SCHEME 1

A solution of compound (I) (2.5 g) in triethylamine (1.6 ml) and methanol (65 ml) was heated at reflux temperature; after 50 min t.l.c.† revealed that all of the starting material (R_F 0.65) had been consumed, and showed the presence of two new components, having R_F values of 0.61 and *ca.* 0.29 (elongated spot), in addition to material which did not migrate. The faster-moving component crystallized from chloroform-ethanol gave pure methyl 2,3-anhydro-4,6-*O*-benzylidene-3-methoxy- α -D-allopyranoside (II), m.p. 187–190°, $[\alpha]_D + 108^\circ$ (*c* 1.32); n.m.r. data: τ 2.4–2.7 (5H multiplet, Ph), 4.40 (1H singlet, benzylidene-methine H), 5.13 (1H doublet, $J_{1,2}$ 3 Hz, 1-H), 6.30 (1H doublet, 2-H), 6.55 and 6.60 (3H singlets, 1-OMe and 3-OMe); no OH or C=O absorptions in i.r. spectrum (Nujol mull). The *D-allo*-configuration was assigned to (II) on the basis of the magnitude of $J_{1,2}$; it has been shown⁵ in several 2,3-anhydroglycopyranosides that, when the 2,3-epoxy-group and the anomeric alkoxy-group were *trans*, $J_{1,2}$ was *ca.* 0, whereas when the groups were in the *cis*-arrangement, $J_{1,2}$ was 2.5–4.5 Hz. The formulation of (II) as an α -methoxyepoxide was substantiated by its conversion, on treatment with lithium aluminium hydride in tetrahydrofuran, into methyl 4,6-*O*-benzylidene-2-deoxy- α -D-*ribo*-hexopyranoside,⁶ m.p. 128–130°, $[\alpha]_D + 138^\circ$ (*c* 1.16). A possible mechanism for the formation of (II) and (I) involves initial epimerization to give the axial toluene-*p*-sulphonate, followed by equatorial attack by methanol on the carbonyl carbon, and finally internal displacement of the toluene-*p*-sulphonyloxy-group by the carbonyl oxygen.


The slower-moving component was isolated by evaporation of the filtrate, remaining after separation of com-

pound (II), to give a residue, which crystallized from chloroform-petroleum, yield 0.563 g (38%). Recrystallization afforded an analytically pure sample of the deriva-



SCHEME 2

tive of 3-hydroxy-2-(hydroxymethyl)-4*H*-pyran-4-one (IV), m.p. 115–117°; n.m.r. data $[(CD_3)_2SO]$: τ 1.90, 3.62 (1H doublets, $J_{5,6}$ 5.6 Hz, 5-H and 6-H), 2.60 (5H, Ph), 4.08 (1H singlet, acetal-methane H), 5.45 (2H singlet, CH_2), and 6.70 (3H singlet, OMe); ν_{max} (Nujol mull) 1658 and



SCHEME 3

† T.l.c. was performed with Silica Gel G as the adsorbent and 2:1 (v/v) ethyl acetate-petroleum (b.p. 60–80°) as the developing solvent. Optical rotations were measured at $23 \pm 2^\circ$ in chloroform. N.m.r. data, obtained at 60 MHz, refer to $CDCl_3$ solutions, unless otherwise stated. All new compounds gave satisfactory elemental analyses.

1640 cm^{-1} ($\cdot\text{C}=\text{C}\cdot\text{CO}\cdot\text{C}=\text{C}\cdot$); positive ferric chloride test. The structure assigned to (IV) was confirmed by the chemical transformations shown in Scheme 2. Thus, acid-catalysed hydrolysis of (IV) gave 3-hydroxy-2-(hydroxymethyl)-4H-pyran-4-one (V), which was converted into maltol (VII) essentially by the procedure described by Stodola,⁷ namely, by treatment of (V) with thionyl chloride to give the chloro-derivative (VI), followed by reduction of (VI) with zinc dust and acetic acid. A possible rationalization for the formation of the γ -pyrone (IV) from the α -keto toluene-*p*-sulphonate (I) on treatment with triethylamine-methanol is shown in Scheme 3.[†]

In a separate experiment, a solution of compound (I) in triethylamine-methanol was heated at reflux temperature for a period longer than 50 min; a third new component (R_F 0.54) was formed, which was shown to be methyl 4,6-*O*-benzylidene- α -D-ribo-hexopyranosid-3-ulose dimethyl acetal (III; R=H), a maximum yield of which was obtained in 48 h, at which time the presence of the α -methoxy-

epoxide (II) could no longer be detected by t.l.c. Compound (III; R=H) was isolated by column chromatography as a syrup, yield 21%, $[\alpha]_D + 112^\circ$ (*c* 1.24); n.m.r. data: τ 7.20 (1H doublet, J 12 Hz, disappeared on deuteration, OH), 5.35 (1H doublet, $J_{1,2}$ 4 Hz, 1-H), 6.38, 6.45, and 6.55 (3H singlets, 1-OMe and two 3-OMe's); ν_{max} (film) 3500 cm^{-1} (OH). The magnitude of the splitting observed for the 1-H signal in the n.m.r. spectrum is consistent with values reported⁸ for an equatorial-axial arrangement of 1-H and 2-H, respectively, in methyl 4,6-*O*-benzylidene- α -D-hexopyranosides. Compound (III; R=H) afforded a crystalline toluene-*p*-sulphonate (III; R=Ts), m.p. 136–138° (decomp.), $[\alpha]_D + 3.51^\circ$ (*c* 1.23).

Full details of these and other reactions of carbohydrate α -keto toluene-*p*-sulphonates will be published later.

We thank the National Research Council of Canada for financial support.

(Received, June 17th, 1971; Com. 993.)

[†] A comparable elimination reaction has recently been reported (W. Meyer zu Reckendorf, *Chem. Ber.*, 1970, **103**, 2418).

¹ R. S. Tipson, *Adv. Carbohydrate Chem.*, 1953, **8**, 107; D. H. Ball and F. W. Parrish, *ibid.*, 1968, **23**, 333; 1969, **24**, 139.

² L. Hough and A. C. Richardson in "Rodd's Chemistry of Carbon Compounds," Vol. 1F, ed. S. Coffey, Elsevier, Amsterdam, 1967, pp. 403–407; A. C. Richardson, *Carbohydrate Res.*, 1969, **10**, 395.

³ B. R. Baker and D. H. Buss, *J. Org. Chem.*, 1965, **30**, 2304; W. A. Szarek, B. T. Lawton, and J. K. N. Jones, *Tetrahedron Letters*, 1969, 4867.

⁴ K. Onodera, S. Hirano, and N. Kashimura, *Carbohydrate Res.*, 1968, **6**, 276; B. T. Lawton, W. A. Szarek, and J. K. N. Jones, *ibid.*, 1969, **10**, 456.

⁵ D. H. Buss, L. Hough, L. D. Hall, and J. F. Manville, *Tetrahedron*, 1965, **21**, 69.

⁶ D. A. Prins, *J. Amer. Chem. Soc.*, 1948, **70**, 3955.

⁷ F. H. Stodola, *J. Amer. Chem. Soc.*, 1951, **73**, 5912.

⁸ B. Coxon, *Tetrahedron*, 1965, **21**, 3481.