

FACILE PREPARATION OF UNSTABLE METABOLIC INTERMEDIATES; EPOXIDE(S) OF PYRAZOLO[1,5-a]PYRIDINE DERIVATIVES BY THE CYTOCHROME P-450 CHEMICAL MODEL

Yoshio NAGATSU, Tsunehiko HIGUCHI, and Masaaki HIROBE*

Faculty of Pharmaceutical Sciences, University of Tokyo, Hongo, Bunkyo-ku, Tokyo 113, Japan

Biomimetic oxidation of bioactive pyrazolo[1,5-a]pyridine derivatives carried out with the chemical model for cytochrome P-450, afforded 6,7-epoxides in relatively high yields, these appear to be chemically unstable precursors of the main metabolites, *i.e.*, 6,7-dihydro-6,7-diols.

KEYWORDS pyrazolo[1,5-a]pyridine derivative; epoxidation; metabolic intermediate; drug metabolism; P-450 chemical model; metalloporphyrin

Cytochrome P-450 plays a very important role in drug metabolism. A wide variety of chemical models for P-450 have been developed to elucidate the functions of the enzymes. However, there have been only a few reports on the application of these model systems to the study of the metabolism of practical drugs having many functional groups.¹⁾ From this point of view, we chose the 3-isobutryl-2-isopropylpyrazolo[1,5-a]pyridine (IBPP), an antiasthma²⁾ and cerebral vasodilator,³⁾ as the substrate to study the reaction by P-450 chemical models. We found that IBPP was oxidized by the TPPMnCl-NaOCl system⁴⁾ to preferentially afford 6,7-epoxide which appeared to be an intermediate of the main metabolite, 6,7-dihydro-6,7-diol, *in vivo*⁵⁾ and *in vitro*, which had not yet been isolated.

We report a novel epoxidation of the pyrazolo[1,5-a]pyridine ring by the chemical model system, metalloporphyrin-NaOCl. The derivatives of pyrazolo[1,5-a]pyridine were synthesized by the method of K. T. Potts *et al.*⁶⁾ Fig. 1 shows the time course of formation of the 6,7-epoxide and 4,5;6,7-diepoxyde of IBPP. Based on these results, the monoepoxide was selectively prepared using TPPMnCl⁷⁾ and the




Fig. 1. Time Course of Epoxidation of IBPP by the Catalyst-NaOCl Chemical Model

The yields of 6,7-epoxide and 4,5;6,7-diepoxyde were determined by HPLC.

diepoxide using mainly $T_{2,6F}PPFeCl^{8)}$ at 120 min. The rate of formation of 6,7-epoxide ($k_1 = 0.42$ mmol/min) was about three times higher than that of 4,5;6,7-diepoxide ($k_1 \times k_2 = 0.16$ mmol/min). In a typical experiment, substrate, 1 mmol; catalyst, 0.01 mmol; benzalkonium chloride (BKC), 0.1 mmol; 4'-imidazolyacetophenone (4'-ImAP), 0.12 mmol; NaOCl, 7 mmol and CH_2Cl_2 , 10 ml were stirred for 2 h at $0^\circ C$ under Ar atmosphere. When the catalyst was $T_{2,6F}PPFeCl$, 4'-ImAP was not used. The epoxidation of the pyrazolo [1,5-a]pyridine derivatives is summarized in Table I.⁹⁾

Table I. Epoxidation of Pyrazolo[1,5-a]pyridine Derivatives by Cytochrome P-450 Chemical Model

Compound I		II	III
R_1	R_2	TPPMnCl	$T_{2,6F}PPFeCl$
$CH(CH_3)_2$	$COCH(CH_3)_2$	16.2 % (16)	56.4 % (113)
$CH(CH_3)_2$	SCN	7.7 (8)	
C_6H_5	H	38.2 ^{a)} (19)	27.7 ^{a)} (28)
C_6H_5	COC_6H_5	9.5 (16)	
$p-SO_2CH_3C_6H_4$	I	12.6 (6)	27.9 (28)
$p-SO_2CH_3C_6H_4$	H	10.1 ^{b)}	
C_6H_5	NO	21.3 ^{c)}	

a) Position 3 of each epoxide was chlorinated; b) Epoxide was not isolated, and gave 3-Cl of compound I; c) Epoxide was not obtained, and gave 3- NO_2 of compound I.

Each value was an isolated yield based on the substrate.

Values in parentheses are turnover numbers based on catalyst.

The 2- C_6H_5 -3-H derivative was chlorinated at position 3 simultaneously, forming epoxides in high yields. The 2- $p-SO_2CH_3C_6H_4$ -3-H derivative was chlorinated at position 3 and the 2- C_6H_5 -3-NO derivative was oxidized to the 3- NO_2 form without epoxidation.

The ring protons of position 6 and 7 in the 6,7-epoxide shifted to 2~2.5 ppm higher magnetic field from IBPP.¹⁰⁾ In the mass spectrum, $M^+ = 246$ m/z of epoxide was 16 m/z higher than IBPP. This mass unit corresponded to one oxygen atom and $(M^+ - 43) = 203$ m/z was the base peak for which the fragment of 43 m/z agreed with $\cdot CH(CH_3)_2$. These results indicated the 6,7-epoxide structure of IBPP. The diepoxide was a single product which was of the syn or anti form from the 1H -NMR spectrum.¹¹⁾ The coupling constants were $J_{6,7} = J_{7,6} = 4.0$ Hz, $J_{4,5} = J_{5,4} = 5.0$ Hz, $J_{5,6} = J_{6,5} = 2.5$ Hz. E. Vogel et al.¹²⁾ showed that the coupling constants of position 2 and 3 of the syn and anti forms of the 1,2;3,4-diepoxide of naphthalene were 2.97 Hz and 1.75 Hz, respectively. From the coupling constant of position 5 and 6 in the diepoxide of pyrazolo[1,5-a]pyridine ring, diepoxide was indicated to be the syn form. This point must be studied further. $M^+ = 262$ m/z was the molecular peak in the mass spectrum of diepoxide and 32 m/z higher than IBPP. The base peak of the diepoxide was $(M^+ - 43) = 219$ m/z.¹³⁾ Other epoxides of pyrazolo[1,5-a]pyridines were identified by 1H -NMR and MS in a similar manner. The main

metabolite of IBPP in humans and various animals is the 6,7-diol form.⁵⁾ Its precursor was expected to be 6,7-epoxide, but has not been isolated and could not be synthesized from IBPP by common oxidizing reagents such as *m*-chloroperbenzoic acid, hydrogen peroxide or hypochlorite. We prepared for the first time, the 6,7-epoxide by the cytochrome P-450 chemical model in a one-step procedure. Because of instability of the 6,7-epoxide in buffer solution ($t_{1/2} = 1.8$ h), it was difficult to obtain appreciable amounts by dehydration with triphenylphosphine of 6,7-dihydro-6,7-diol, which had been isolated from rabbit urine by Miura *et al.*¹⁴⁾ In the present reaction system, the active species was considered to be a metalloxenoid.¹⁵⁾ Consequently, position 6 and 7 of the pyrazolo[1,5-*a*]pyridine ring was primarily epoxidized in a manner similar to those of olefins,⁴⁾ followed by epoxidation of position 4 and 5. Therefore, the epoxide did not seem to be formed *via* the 4,5-monoepoxide. In this reaction system, epoxidation did not occur without a catalyst. We succeeded in synthesizing the relatively unstable 6,7-epoxide efficiently by direct oxidation of IBPP with a chemical P-450 model system TPPMnCl-NaOCl. Chemical P-450 models could be used to obtain the unstable metabolites and to study the metabolism of drugs.

ACKNOWLEDGEMENTS

This work was supported in part by a Grant-in-Aid for Special Project Research from the Ministry of Education, Science and Culture, Japan. We thank the Central Research Laboratories of Kyorin Pharmaceutical Co., Ltd. for supplying the IBPP.

REFERENCES AND NOTES

- 1) a) H. Masumoto, K. Takeuchi, S. Ohta and M. Hirobe, *Chem. Pharm. Bull.*, submitted (1988); b) F. Pautet, R. Barret and M. Daudon, *Pharmazie*, **41**, 286 (1986).
- 2) M. Ohashi, R. Kanai, K. Nishino, T. Sato and I. Takayanagi, *Prostaglandins*, **32**, 875 (1986).
- 3) a) T. Irikura, Y. Kudo, M. Ohashi, R. Ishida, J. Kito, M. Kodaira, H. Uchida and K. Nishino, *Kiso to Rinsho*, **17**, 104 (1983); b) T. Irikura, Y. Kudo, H. Ohkubo, M. Ohashi, J. Kito and K. Nishino, *Oyo Yakuri*, **25**, 283 (1983).
- 4) J. P. Collman, J. I. Brauman, B. Meunier, T. Hayashi, T. Kodadek and S. A. Raybuck, *J. Am. Chem. Soc.*, **107**, 2000 (1985).
- 5) K. Takagi and K. Endo, *Oyo Yakuri*, **27**, 1167 (1985).
- 6) K. T. Potts, U. P. Singh and J. Bhattacharyya, *J. Org. Chem.*, **33**, 3766 (1968).
- 7) TPPMnCl: meso-Tetraphenylporphyrinatomanganese chloride was prepared according to the method in the following report; A. D. Adler, *J. Org. Chem.*, **32**, 467 (1967).
- 8) $T_{2,6F}PPFeCl$: meso-Tetrakis-2,6-difluorophenylporphyrinatoiron chloride was prepared according to the method in the following report; R. W. Wagner, D. S. Lawrence and J. S. Lindsey, *Tetrahedron Lett.*, **28**, 3069 (1987).
- 9) 6,7-Epoxide and 4,5;6,7-diepoxy were isolated by flash silica gel column chromatography. Some of the 6,7-epoxide was lost by decomposition on silica gel during column chromatography. The reaction condition in Table I was optimized while the condition in Fig. 1 was not. Therefore the yield of diepoxy in Table I was higher than that in Fig. 1.
- 10) 1H -NMR (400MHz, $CDCl_3$) δ ; 7.22 (1H, d, 4-H), 6.46 (1H, dd, 5-H), 5.18 (1H, d, 7-H), 4.11 (1H, t, 6-H), 3.46 (1H, m), 3.14 (1H, m), 1.34 (6H, d, 3- CH_3), 1.19 (6H, d, 2- CH_3).
- 11) 1H -NMR (400MHz, $CDCl_3$) δ ; 5.18 (1H, d, 7-H), 4.26 (1H, d, 4-H), 4.10 (1H, d, 5-H), 4.08 (1H, d, 6-H), 3.40 (1H, m), 3.15 (1H, m), 1.30 (6H, d, 3- CH_3), 1.20 (6H, d, 2- CH_3).
- 12) E. Vogel, H. H. Klug and M. S. Ridder, *Angew. Chem. Int. Ed. Engl.*, **15**, 229 (1976).
- 13) Elemental analysis of diepoxy was as follows: Calcd. for $C_{14}H_{18}N_2O_3$: C, 64.11; H, 6.92; N, 10.68. Found: C, 64.30; H, 6.89; N, 10.56.
- 14) Unpublished data.
- 15) R. J. M. Nolte, J. A. S. J. Razenberg and R. Schuurman, *J. Am. Chem. Soc.*, **108**, 2751 (1986).

(Received February 10, 1989)