

[Click for updates](#)

Organic Preparations and Procedures International: The New Journal for Organic Synthesis

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/uopp20>

Alternate and Step-Economic Synthesis of the β -Methylstyrene Chelating Pre-ligand of the Hoveyda-Grubbs' II Catalyst

Sebastian O. Simonetti^a, Enrique L. Larghi^a & Teodoro S. Kaufman^a

^a Instituto de Química Rosario (IQUIR, CONICET-UNR) and Facultad de Ciencias Bioquímicas y Farmacéuticas-Universidad Nacional de Rosario, Suipacha 531, S2002LRK Rosario, Argentina

Published online: 13 May 2015.

To cite this article: Sebastian O. Simonetti, Enrique L. Larghi & Teodoro S. Kaufman (2015) Alternate and Step-Economic Synthesis of the β -Methylstyrene Chelating Pre-ligand of the Hoveyda-Grubbs' II Catalyst, Organic Preparations and Procedures International: The New Journal for Organic Synthesis, 47:3, 227-231, DOI: [10.1080/00304948.2015.1025021](https://doi.org/10.1080/00304948.2015.1025021)

To link to this article: <http://dx.doi.org/10.1080/00304948.2015.1025021>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms &


OPPI BRIEF

Alternate and Step-Economic Synthesis of the β -Methylstyrene Chelating Pre-ligand of the Hoveyda-Grubbs' II Catalyst

Sebastian O. Simonetti, Enrique L. Larghi,
and Teodoro S. Kaufman

Instituto de Química Rosario (IQUIR, CONICET-UNR) and Facultad de Ciencias Bioquímicas y Farmacéuticas-Universidad Nacional de Rosario, Suipacha 531, S2002LRK Rosario, Argentina

The Hoveyda-Grubbs (HG)¹ series of olefin metathesis catalysts are more selective, recyclable, and heat, air and moisture stable than their predecessors.² β -Methylstyrenes, such as *E*-1-isopropoxy-2-propenylbenzene (**3**, *Scheme 1*), are currently preferred as pre-ligands for the synthesis of the second generation of the HG catalysts (HG-II),³ because they are less costly, easier to prepare, more stable and less prone to undergo spontaneous polymerization or homo-dimerization than the styrenes unsubstituted at the β -position, originally used for that purpose.^{4,5}




Previous syntheses of compound **3** (*Scheme 1*), the chelating pre-ligand of HG-II and related olefin metathesis catalysts,⁶ involved several steps, some of which were slow or tedious, requiring long reaction times,³ or proceeded only in moderate yields,³ employed high excess of alkylating agents, stoichiometric amounts of organometallic reagents,⁷ or afforded undesirable *E/Z* mixtures of olefins.⁸ In addition, an approach to related β -methylstyrenes via the Wittig reaction has been found to proceed in relatively low yields, and required the removal and disposal of large amounts of triphenylphosphine oxide; therefore, it was discarded as a viable route toward **3**.^{3,7}

We have recently reported an efficient one-pot sequential homo-bimetallic catalytic propenylation of haloarenes with allyltri(*n*-butyl)tin under palladium catalysis and optimized the reaction conditions.^{9,10} As a further application of this novel strategy, with impact in fine chemistry, herein we report an alternate and convenient, step-economic synthesis of **3**, from commercial 2-halophenols **8a,b** (*Scheme 2*).

Submitted by June 28, 2014

Address correspondence to Teodoro S. Kaufman, Instituto de Química Rosario (IQUIR, CONICET-UNR) and Facultad de Ciencias Bioquímicas y Farmacéuticas Universidad Nacional de Rosario, Suipacha 531, S2002LRK Rosario, Argentina. E-mail: kaufman@iquir-conicet.gov.ar



Scheme 1


a) K_2CO_3 , Cs_2CO_3 (cat.), *i*-PrI, DMF, 40°C, 24–48 h (93%); b) $\text{RhCl}_3 \cdot 3\text{H}_2\text{O}$ (cat.), *p*-TsOH·H₂O (cat.), 90% aq. EtOH, reflux, 5 h (71%, *E/Z* = 14.3:1) or $\text{CH}_2=\text{CHOTMS}$, Grubbs II (cat.), CH_2Cl_2 , 35°C, 24 h (66%); c) 1. $(\text{PPh}_3)_2\text{Cl}_2\text{Ru}$, 70°C, 17.5 h; 2. $\text{P}(\text{CH}_2\text{OH})_3$ [$> 99\%$ (GC), *E/Z* = 45:55]; d) K_2CO_3 (2 equiv.), *i*-PrBr (2 equiv.), DMF, 60°C, 48 h (93%); e) K_2CO_3 , *i*-PrBr, DMF, 60°C, 9 h; f) EtMgCl , THF, −15°C; g) TsOH, PhMe, 90°C, 1.5 h (58% overall, *E/Z* = 96:4).

The Williamson etherification of **8a** or **8b** was performed with 2-bromopropane and K_2CO_3 in EtOH¹ (Scheme 2), to afford **9a** and **9b** in 84% and 86% yield respectively, after 5 h of heating under reflux. This contrasted with previous preparations of **9a,b** involving comparatively poor eco-friendly solvents, strong bases,¹² or the relatively more expensive isopropyl iodide.

Next, the key homo-bimetallic catalytic reaction of allyltri(*n*-butyl)tin with the bromoethers **9a,b** was carried out in the presence of $\text{Pd}(\text{PPh}_3)_2\text{Cl}_2$ in diglyme at 130°C for 14 h. This process cleanly executed the Stille cross-coupling to the allyl intermediate³ which further underwent an *in situ* palladium-catalyzed conjugative double bond migration to give compound **3**.

Removal of triphenylphosphine by oxidation to triphenylphosphine oxide with H_2O_2 ,¹³ followed by treatment of the crude extract of the reaction with an activated fluorosilica gel phase (10% w/w),¹⁴ led to the complete elimination of all tin residues (GC-MS analysis), affording the expected product **3** in yields of 73% and 60% from **9a** and **9b**, respectively.

The present two-step alternative approach to **3** complements previously reported syntheses of this important and valuable β -methylstyrene derivative. It can be easily carried out in a short time transforming the sequence into a cost-competitive process, counterbalancing the expense of the chemicals involved. The wider scope of the present route and its adaptability to access similar compounds with various functional groups¹⁰ make it



Scheme 2

a) Me_2CHBr , K_2CO_3 , EtOH, reflux, 5 h (**9a**, 84%; **9b**, 86%); b) $n\text{-Bu}_3\text{SnCH}_2\text{CH}=\text{CH}_2$, Pd $(\text{PPh}_3)_2\text{Cl}_2$, LiCl, PPh_3 , Diglyme, 130°C, 6 h; c) 130°C, 8 h (73% from **9a**; 60% from **9b**).

useful for the preparation of related otherwise less accessible, pre-ligand members of this family.¹⁵ The availability of redundant pathways for the synthesis of small molecules with relevance in fine chemistry is always desirable, because of the potential economic value of such compounds. This is especially critical when intellectual property protection conditions restrict the use of the patent-protected processes.^{6,7}

Experimental Section

The reactions were carried out under dry argon atmospheres, employing oven-dried glassware. Ethanol was dried by treatment with Mg turnings and a catalytic amount of I₂, and distilled from the resulting magnesium ethoxide. Diglyme was distilled from sodium benzophenone ketyl.¹⁶ Anhydrous solvents were stored in dry Schlenk tubes. The remaining reagents were used as received. The IR spectra were acquired on a Shimadzu Prestige 21 spectrophotometer, as thin films. The ¹H and ¹³C NMR spectra were recorded at 300.13 and 75.48 MHz respectively, on a Bruker Avance 300 spectrometer, employing CDCl₃ [δ_{H} 7.26 (residual CHCl₃) and δ_{C} 76.9 ppm] as internal standard. Chemical shifts are reported in δ scale and the coupling constants (*J*) are given in Hertz. The GC-MS data were obtained on a Shimadzu QP2010 *plus* instrument (EI = 70 eV). The high resolution MS were obtained on a Bruker MicroTOF-Q II instrument. Detection of the ions was performed with electrospray ionization in positive ion mode. The compounds gave single spots when run on Kieselgel 60 GF₂₅₄ TLC plates, employing hexane-EtOAc as the solvent system. The chromatographic spots were detected by exposing the plates to UV light (254 nm), followed by spraying with *p*-anisaldehyde/sulfuric acid reagent in EtOH and careful heating up to 120°C.

General Procedure for the Synthesis of 1-halo-2-isopropoxybenzenes (9a and 9b)

A mixture of the 2-halophenol (**8a,b** 1.156 mmol), 192 mg (1.2 equiv. anhydrous K₂CO₃ (1.20 equiv.) and sodium iodide (4 mg) in absolute EtOH (3 mL) was magnetically stirred at ambient temperature under argon. The 2-bromopropane (1.20 equiv.), was added slowly and the mixture was then refluxed for 5 h, cooled to ambient temperature prior to the removal of the solvent under reduced pressure. Addition of EtOAc (50 mL) gave a solution which was washed successively with 1M NaOH and brine (15 mL each). The organic phase was dried over Na₂SO₄, and evaporated under reduced pressure to give a residue that was purified by flash column chromatography.

1-Bromo-2-isopropoxybenzene (9a)¹⁷

Oil; yield: 84%. IR (film): 602, 748, 953, 1030, 1107, 1275, 1385, 1472, 1587, 2978 cm⁻¹. ¹H NMR (300 MHz, CDCl₃): δ 1.38 (d, *J* = 5.9, 6H, CHMe₂), 4.55 (septet, *J* = 5.9, 1H, OCHMe₂), 6.81 (dt, *J* = 1.2 and 7.6, 1H, 3-H), 6.91 (dd, *J* = 0.6 and 7.9, 1H, 5-H), 7.22 (dt, *J* = 1.5 and 8.0, 1H, 4-H), 7.53 (dd, *J* = 1.6 and 7.9, 1H, 6-H). ¹³C NMR (75 MHz, CDCl₃): δ 22.1 (Me₂-CH), 72.1 (Me₂-CH), 113.8 (1-C), 115.9 (5-C), 121.9 (3-C), 128.2 (4-C), 133.5 (6-C), 154.6 (2-C). MS (EI, 70 eV): *m/z* (%) 172 (100), 174 (95), 214 [M]⁺ (25), 216 [M+2]⁺ (25).

1-Iodo-2-isopropoxybenzene (9b)¹⁸

Oil; yield: 86%. IR (film): 648, 748, 953, 1045, 1126, 1273, 1381, 1470, 1577, 2931, 2978 cm⁻¹. ¹H NMR (300 MHz, CDCl₃): δ 1.39 (d, *J* = 5.9, 6H, CHMe₂), 4.56 (septet, *J*

= 5.9, 1H, OCHMe₂), 6.69 (dt, $J = 1.6$ and 8.0 , 1H, 3-H), 6.85 (dd, $J = 1.6$ and 8.0 , 1H, 5-H), 7.25 (dt, $J = 1.6$ and 8.0 , 1H, 4-H), 7.78 (dd, $J = 1.6$ and 8.0 , 1H, 6-H). ¹³C NMR (75 MHz, CDCl₃): δ 22.1 (Me₂-CH), 72.1 (Me₂-CH), 88.6 (1-C), 114.4 (3-C), 122.5 (5-C), 129.3 (4-C), 139.6 (6-C), 156.8 (2-C). MS (EI, 70 eV): m/z (%) 65 (37), 93 (34), 220 [M-C₃H₆]⁺ (100), 262 [M]⁺ (25).

E-1-Isopropoxy-2-propenylbenzene (3)

A degassed solution containing the aryl bromide (**9a**, 183 mg, 0.852 mmol), anhydrous LiCl (286 mg, 6.82 mmol), Ph₃P (112 mg, 0.426 mmol) and Pd(Ph₃P)₂Cl₂ (71.7 mg, 0.102 mmol) in anhydrous diglyme (2 mL, final concentration of **9a** ca. 0.15 M) was treated dropwise with allyltri(*n*-butyl)tin (0.32 mL, 1.02 mmol). The reaction mixture was heated at 130°C under an argon atmosphere. A sample taken at 6 h and worked up allowed the isolation of *l*-allyl-2-isopropoxybenzene.³

IR (film): 605, 734, 864, 995, 1046, 1174, 1287, 1372, 1383, 1489, 1587, 1638, 2931, 2978, 3077 cm⁻¹. ¹H NMR (300 MHz, CDCl₃): δ 1.33 (d, $J = 6.0$ Hz, 6H), 3.37 (d, $J = 6.7$ Hz, 2H), 4.54 (septet, $J = 6.0$ Hz, 1H), 5.00–5.09 (m, 2H), 5.93–6.03 (m, 1H), 6.83–6.89 (m, 2H), 7.12–7.18 (m, 2H). ¹³C NMR (75 MHz, CDCl₃): δ 22.2, 34.6, 69.9, 113.0, 115.2, 120.2, 127.1, 129.7, 129.9, 137.2, 155.5. MS (EI, 70 eV): m/z (%) 91 (13), 115 (12), 119 (46), 133 (46), 134 (100), 135 (10), 176 [M]⁺ (66), 177 [M+1]⁺ (9). MS found: 176.1207; C₁₂H₁₆O requires m/z : 176.1201.

When the reaction was heated during 14 h, complete consumption of the starting material took place, as determined by GC analysis. Then, the reaction mixture was allowed to attain ambient temperature and was diluted with 25 mL of hexanes and filtered through a short pad of Florisil and Celite (1:1). The filtrate was washed successively with saturated NaF (4 × 10 mL), H₂O (3 × 10 mL), brine (15 mL) and H₂O₂ (10 mL of a 0.12% P/V solution), dried over Na₂SO₄ and evaporated *in vacuo*. The oily residue was re-dissolved in CH₂Cl₂ and stirred an additional period of 12 h with 200 mg activated fluorosilica. Filtration and evaporation of volatiles afforded 90 mg (73%) of the expected β -methylstyrene (**3**), as a colorless oil.

IR (film): 606, 782, 862, 959, 1049, 1176, 1289, 1383, 1484, 1596, 1655, 2876, 2933, 3070 cm⁻¹. ¹H NMR (300 MHz, CDCl₃): δ 1.35 (d, $J = 5.9$, 6H, Me₂-CH), 1.90 (dd, $J = 1.5$ and 6.6 , 3H, Me-CH=CH-Ar), 4.57 (septet, $J = 5.9$, 1H, Me₂-CH), 6.22 (dq, $J = 6.0$, 13.9 and 15.2, 1H, Me-CH=CH-Ar), 6.72 (dq, $J = 1.5$ and 15.9, 1H, CH₃-CH=CH-Ar), 6.85–6.90 (m, 2H, ArH), 7.14 (dt, $J = 1.5$ and 7.7, 1H, 4-H), 7.41 (dd, $J = 1.4$ and 7.5, 1H, 6-H). ¹³C NMR (75 MHz, CDCl₃): δ 18.9 (Me-CH=CH-Ar), 22.2 (Me₂-CH-O), 70.8 (Me₂-CH), 114.3 (6-C), 120.7 (5-C), 125.8 (2-C), 125.9 (CH₃-CH=CH-Ar), 126.4 (CH₃-CH=CH-Ar), 127.5 (3-C), 128.2 (5-C), 154.6 (1-C). MS (EI, 70 eV): m/z (%) 77 (10), 91 (24), 105 (13), 107 (9), 115 (17), 117 (12), 119 (69), 133 (54), 134 (100), 147 (12), 176 [M]⁺ (52), 177 [M+1]⁺ (7). MS found m/z = 176.1198; C₁₂H₁₆O (M⁺) requires m/z = 176.1201. The spectral data were in full agreement with those described.^{2,3,5,8}

Acknowledgements

The authors thank Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET) and Agencia Nacional de Promoción Científica y Tecnológica (ANPCyT) for financial support. S.O.S. thanks CONICET for his fellowship.

References

1. J. S. Kingsbury, J. P. A. Harrity, P. J. Bonitatebus Jr. and A. H. Hoveyda, *J. Am. Chem. Soc.*, **121**, 791 (1999).
2. R. H. Grubbs, *Handbook of Metathesis*, Vols. 1–3. Wiley-VCH Weinheim, 2003.
3. R. Bujok, M. Bieniek, M. Masnyk, A. Michrowska, A. Sarosiek, H. Steüpowaska, D. Arlt and K. Grela, *J. Org. Chem.*, **69**, 6894 (2004).
4. S. B. Garber, J. S. Kingsbury, B. L. Gray and A. H. Hoveyda, *J. Am. Chem. Soc.*, **122**, 8168 (2000).
5. S. Gessler, S. Randl and S. Blechert, *Tetrahedron Lett.*, **41**, 9973 (2000).
6. K. Skowerski, L. Gulajski and M. Bieniek, PCT Intl. Appl. WO 127880, 2013; *Chem. Abstr.*, **159**, 427938 (2013).
7. R. L. Pederson, J. K. Woertink, C. M. Haar, D. E. Gindelberger and Y. Schrodi, *US Patent* 6620955; *Chem. Abstr.*, **139**, 7375 (2003).
8. M. Bieniek, A. Michrowska, D. L. Usanov and K. Grela, *Chem. Eur. J.*, **14**, 806 (2008).
9. S. O. Simonetti, E. L. Larghi, A. B. J. Bracca and T. S. Kaufman, *Org. Biomol. Chem.*, **10**, 4124 (2012).
10. S. O. Simonetti, E. L. Larghi and T. S. Kaufman, *Org. Biomol. Chem.*, **12**, 3735 (2014).
11. A. B. J. Bracca and T. S. Kaufman, *Eur. J. Org. Chem.*, 5284 (2007).
12. G. Chiu, S. Li, P. J. Connolly, V. Pulito, J. Liu and S. A. Middleton, *Bioorg. Med. Chem. Lett.*, **17**, 3930 (2007).
13. C. R. Hilliard, N. Bhuvanesh, J. A. Gladysz and J. Blümel, *Dalton Trans.*, **41**, 1742 (2012).
14. D. C. Harrowven and I. L. Guy, *Chem. Commun.*, 1968 (2004).
15. A. Michrowska, R. Bujok, S. Harutyunyan, V. Sashuk, G. Dolgonos and K. Grela, *J. Am. Chem. Soc.*, **126**, 9318 (2004).
16. W. L. F. Armarego and C. L. L. Chai, *Purification of Laboratory Chemicals*, 6th Edition, Butterworth-Heinemann, Burlington, USA, 2009, p. 126 and 127.
17. A. Y. Kuposov, R. R. Karimov, I. M. Geraskin, V. N. Nemykin and V. V. Zhdankin, *J. Org. Chem.*, **71**, 8452 (2006).
18. T. Ishikawa, K. Shimooka, T. Narioka, S. Noguchi, T. Saito, A. Ishikawa, E. Yamazaki, T. Harayama, H. Seki and K. Yamaguchi, *J. Org. Chem.*, **65**, 9143 (2000).