



Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/gpss20>

Conversion of Oximes into Carbonyl Compounds with Thallium(III) Nitrate Supported onto HZSM-5 Zeolite Under Microwave Irradiation in Solventless System

Majid M. Heravi^a & Mitra Ghassemzadeh^b

^a Azzahra University, Vanak, Tehran, Iran

^b Chemistry and Chemical Engineering Research Center of Iran, Tehran, Iran

Published online: 27 Oct 2010.

To cite this article: Majid M. Heravi & Mitra Ghassemzadeh (2003) Conversion of Oximes into Carbonyl Compounds with Thallium(III) Nitrate Supported onto HZSM-5 Zeolite Under Microwave Irradiation in Solventless System, Phosphorus, Sulfur, and Silicon and the Related Elements, 178:1, 119-123, DOI: [10.1080/10426500307816](https://doi.org/10.1080/10426500307816)

To link to this article: <http://dx.doi.org/10.1080/10426500307816>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

CONVERSION OF OXIMES INTO CARBONYL COMPOUNDS WITH THALIUM(III) NITRATE SUPPORTED ONTO HZSM-5 ZEOLITE UNDER MICROWAVE IRRADIATION IN SOLVENTLESS SYSTEM

Majid M. Heravi^a and Mitra Ghassemzadeh^b
Azzahra University, Vanak, Tehran, Iran^a and Chemistry and
Chemical Engineering Research Center of Iran, Tehran, Iran^b

(Received June 20, 2002)

Oximes are oxidatively deprotected by thalium(III) nitrate supported onto HZSM-5 zeolite to the parent carbonyl compounds in high yields upon exposure to microwave irradiation.

Keywords: Carbonyl compounds; deoximation; microwave irradiation; thalium nitrate; zeolite

INTRODUCTION

Oximes frequently are used to protect carbonyl compounds and hence much attention has been paid to develop methods for their deprotection.¹ Since oximes can be prepared from noncarbonyl compounds, its conversion to carbonyl compounds can be considered a useful method for the synthesis of the latter.^{2,3}

The classical recovery of aldehydes or ketones from oximes consists of acid hydrolysis which removes the hydroxylamine from the equilibrium⁴ and often results in low yields. Although a large number of reagents are found in the literature for cleavage of oximes to carbonyl compounds,⁵ there still appears a need either to improve the existing methods or to introduce newer reagents to permit better selectivity with easy work up procedures.

M. M. Heravi is thankful for the partial financial support from Azzahra University Research Council.

Address correspondence to Majid M. Heravi, Department of Chemistry, School of Sciences, Azzahra University, Vanak, Tehran, Iran. E-mail: mmheravi@azzahra.ac.ir

Thalium(III) nitrate has been used as an oxidizing agent⁶ and Willgerodth-Kindler reaction.⁷ Recently, there has been growing interest in the application of microwave irradiation in chemical reaction enhancement,⁸ the salient features being improved reaction rates and formation of cleaner products.

When this technique is combined with solvent-free conditions, it would be especially appealing as it provides an opportunity to work with open vessels, thus avoiding the risk of high pressure development, and a possibility of carrying out the reaction on a preparative scale in addition to the associated selectivity and ease of manipulation.⁹ The microwave enhanced chemical reactions on mineral supports have gained much attention.¹⁰

In continuation of our ongoing program to develop environmentally benign reactions under solvent-free conditions using supported reagent combined with microwave,¹¹ in this article we report the deoxygenation reaction using thalium(III) nitrate supported onto HZSM-5 zeolite under microwave irradiation in solventless system.



















RESULTS AND DISCUSSION

Recently, we reported¹² on the use of HZSM-5 zeolite as a support for the oxidation of alcohols. Among various mineral supports examined such as alumina, clay, silica, and Na-zeolite, we found that HZSM-5 zeolite gives the best results. A weight equivalent of HZSM-5 zeolite and thalium(III) nitrate were crushed using pestle and mortar to give an intimate pair. Two equivalents of this mixture were crushed with the equivalent of neat benzaldoxime and exposed to microwave irradiation.

The progress of the reaction was monitored by TLC using petroleum ether-ethyl acetate, 8:2, as eluents. After 2 min deoxygenation of benzaldoxime was complete to afford benzaldehyde almost in quantitative yield.

To assess the generality of this deoxygenation method a variety of aldioximes and ketoximes were reacted under these conditions which yielded to excellent yields (Table I). The interesting feature of this method is the successful deoxygenation of ketones. As with some other reagents it results in the formation of Beckmann rearrangement¹³ products. However, this method is unsuccessful when applied to aryl aldioximes and ketoximes which contain strong electron donating groups such as $-\text{NH}_2$ or $-\text{OH}$ at *ortho* or *para* position most probably due


TABLE I Deoxygenation Reaction Using Thallium(III) Nitrate Under Microwave Irradiation in a Solventless System

| Entry | Substrate | Reaction time (sec) | Product ^a | Yield ^b (%) |
|-------|---|---------------------|---|------------------------|
| 1 |  | 120 |  | 98 |
| 2 |  | 120 |  | 83 |
| 3 |  | 80 |  | 85 |
| 4 |  | 180 |  | 82 |
| 5 |  | 240 |  | 80 |
| 6 |  | 240 |  | 85 |
| 7 |  | 180 |  | 92 |
| 8 |  | 240 |  | 83 |
| 9 |  | 240 |  | 90 |

^aAll products were known and identified by comparison of their physical data with those of authentic sample.

^bYields are based on isolated product.

to concomitant oxidation of the aromatic ring to quinone derivatives (Scheme 1).



SCHEME 1

This phenomenon has been noticed previously.¹⁴ This limitation, however, can be circumvented by acetylation of the OH and NH₂ groups before the formation of oximes.¹⁴

In conclusion we have developed a rapid and solventless method for cleavage of oxime compounds. This high yielding method is manipulative simple and a selective protocol when compared to the conventional solution phase or heterogeneous reactions.

EXPERIMENTAL SECTION

All substrates were synthesized by a known literature procedure. All products were characterized by comparison of their physical and spectroscopic data with those of authentic samples.

Deprotection of Oximes. Typical Procedure.

Thalium(III) nitrate (4 mmol) and benzylaldoxime (2 mmol) were crushed using a pestle and mortar. This mixture was transferred to a beaker and exposed to microwave irradiation. The progress of the reaction was monitored by TLC using petroleum ether: ethyl acetate 8:2. After completion of the reaction, the product was extracted with dichloromethane and filtered. The filtrate was evaporated to dryness to afford benzaldehyde. Yield 98%.

Caution: Although this reaction worked safely in our hand, using a microwave oven in an efficient hood is highly recommended.

REFERENCES

- [1] T. W. Greene and P. G. M. Wuts, *Protective Groups in Organic Synthesis* (John Wiley and Sons, New York, 1999), 3rd ed.
- [2] D. H. R. Barton, J. M. Beaton, L. E. Geller, and M. M. Pechel, *J. Am. Chem. Soc.*, **83**, 4076 (1961).
- [3] A. R. Katritzky, O. Meth-Cohn, and C. W. Rees, eds., *Comprehensive Organic Functional Groups Transformations* (Pergamon, New York, 1995), 1st ed., vol. 3.
- [4] E. B. Hersberg, *J. Org. Chem.*, **13**, 542 (1948).
- [5] a) N. B. Barhate, A. S. Gajare, R. D. Wakarkar, and A. Sudalai, *Tetrahedron Lett.*, **38**, 653 (1997); b) Y. S. Yadav, P. K. Sasmal, and P. K. Chand, *Synth. Commun.*, **29**, 3667 (1999); c) B. A. Nattier, K. J. Eash, and R. S. Mohan, *Synthesis*, 1010 (2001); d) A. Carsaro, A. Chiacchio, and V. Pistara, *Synthesis*, **13**, 1903 (2001).
- [6] A. McKillop, J. D. Hunt, R. D. Naylor, and F. C. Taylor, *J. Am. Chem. Soc.*, **93**, 4918 (1971).
- [7] A. McKillop, J. D. Hunt, R. D. Naylor, and F. C. Taylor, *J. Am. Chem. Soc.*, **93**, 4919 (1971).
- [8] R. A. Abramivoch, *Org. Prep. Proced. Int.*, **23**, 683 (1991).
- [9] R. S. Varma, R. Dahija, and R. K. Saini, *Tetrahedron Lett.*, **38**, 8819 (1997).
- [10] R. S. Varma and R. Dahija, *Tetrahedron Lett.*, **38**, 2043 (1997).
- [11] a) M. M. Heravi, D. Ajami, M. Ghassemzadeh, and M. M. Mojtahedi, *Tetrahedron Lett.*, **40**, 561 (1999); b) M. M. Heravi, D. Ajami, and M. Ghassemzadeh, *Synthesis*, **3**, 393 (1999); c) M. M. Heravi, D. Ajami, K. Aghapoor, and M. Ghassemzadeh, *J. Chem. Soc., Chem. Commun.*, 833 (1999); d) M. M. Heravi, D. Ajami, and M. M. Mojtahedi, *J. Chem. Res.*, 126 (2000).
- [12] M. M. Heravi, D. Ajami, K. Tabar-Hydar, and M. Ghassemzadeh, *J. Chem. Res.*, 334 (1999).
- [13] A. Bosch, P. Cruz, A. Diez-Barra, A. Loupy, and F. Langa, *Synlett.*, 1259 (1995).
- [14] A. McKillop, B. P. Swann, and E. C. Taylor, *Tetrahedron Lett.*, **26**, 4031 (1970).