Triphenylmethyldifluoramine: a stable reagent for the synthesis of *gem*-bis(difluoramines)[†]

G. K. Surya Prakash,*^a Markus Etzkorn,^a George A. Olah,*^a Karl O. Christe,^{ab} Stefan Schneider^a and Ashwani Vij^b

^a Loker Hydrocarbon Research Institute and Department of Chemistry, University of Southern California, University Park, Los Angeles, CA 90089-1661, USA. E-mail: gprakash@usc.edu; Fax: +1 213 740 6270; Tel: +1 213 740 5984

^b Air Force Research Laboratory, Edwards Air Force Base, CA, 93524, USA

Received (in Corvallis, OR, USA) 16th April 2002, Accepted 13th June 2002 First published as an Advance Article on the web 10th July 2002

The conversion of ketones into geminal bis(difluoramines) can be achieved under mild two-phase reaction conditions by employing triphenylmethyldifluoramine as an *in situ* source of difluoramine.

The introduction of a difluoramine functionality into organic moieties and the properties of the obtained target compounds have been investigated intensively.¹ More recently a new class of high-energy containing materials, *gem*-bis(difluoramine)-substituted heterocyclic nitramines has gained much attention as high-energy oxidizers: HNFX (1)² and TNFX (2)³ have been successfully synthesized from their corresponding ketone derivatives. Usually this transformation is achieved under strongly acidic conditions with an excess of condensed difluoramine (HNF₂),⁴ an unpredictably shock-sensitive and thermally unstable, gaseous compound.⁵ HNF₂ can be generated from different precursors, *e.g.* tetrafluorohydrazine,⁶ *N*,*N*-difluorourea⁷ or *N*,*N*-difluoroamine (**3**) mentions its hydrolysis to HNF₂.⁹ Except for **3**, a stable crystalline compound, all of the other HNF₂ precursors have only limited stability.



Compound 3 proved to be a convenient reagent for the conversion of ketones (A) into *gem*-bis(difluoramines) (B) thus circumventing the potential hazards associated with previous methodologies.



Triphenylmethyldifluoramine (**3**) is accessible from chlorotriphenylmethane and N₂F₄ in multi-gram quantities as stable, non-hygroscopic, colorless crystals.⁹† It can be stored without decomposition or any other hazards at ambient conditions. The complete characterization of **3** was carried out by IR, Raman, ¹H, ¹³C and ¹⁹F NMR spectroscopy. Additionally, we obtained single crystals suitable for a X-ray crystal structure determination.‡

[†] Electronic supplementary information (ESI) available: optimized preparation of **3**, analytical data for **3**, **18–21** and a plot of the unit cell of **3**. See http://www.rsc.org/suppdata/cc/b2/b203811k/ The structural features of **3** are similar to those reported for *p*bromophenyldiphenyl)methyldifluoramine.¹⁰ The C–N distance of 1.523(2) Å found in **3** is longer than the corresponding distance of 1.481 Å in triphenylmethylamine.¹¹ The presence of two highly electronegative fluorines on the nitrogen atom reduces back-donation from the nitrogen lone pair, thereby decreasing the C–N bond order. The torsion angles from the difluoramine group to the *ipso*-carbon atoms of the phenyl rings, namely F1–N1–C1–C8 and F2–N1–C1–C2 are 56.4(1)° and 39.1(1)°, respectively, and reflect the deviation of these atoms from perfect staggering.



The closest intramolecular H···F contacts found in the crystal structure of **3** result from H3···F1 and F2···F15 at 2.281(16) and 2.334(16) Å, respectively. The closest intermolecular H···N contacts result from H15···N1 and H9···N1 at 2.408(16) and 2.553(16) Å, respectively. All these contacts are shorter than the sum of the van der Waals radii of nitrogen and hydrogen (1.55 + 1.20 = 2.75 Å) or fluorine and hydrogen (1.47 + 1.20 = 2.67 Å). A plot of the unit cell is provided with the ESI.†‡

We have investigated two-phase organic solvent/oleum media bis(difluoraminations) with **3** under mild reaction conditions for some model ketones. Typically a solution of the ketone (**A**) and 2.2 equivalents of **3** in CH₂Cl₂, CHCl₃ or CFCl₃ were added dropwise to the same volume of oleum (30%) at 0 °C. Due to the potential hazards of the intermediate HNF₂ and any labile products, all reactions were carried out on a small scale (~1 mmol). The completion of the reactions was confirmed by NMR spectroscopy (¹⁹F, ¹H, ¹³C). Thus, they were usually worked up after 15–20 min by pouring into ice/ water, followed by extraction (CH₂Cl₂) and evaporation of the solvent from the dried organic phase.

The crude product contained only triphenylmethanol and the *gem*-bis(difluoramine) that (on a larger scale) can be purified by repeated crystallization, chromatography on neutral aluminum

Table 1 Conversion of ketones to gem-bis(difluoramines)

Entry	Ketone (A)	gem-Bis(difluoramine) (B)	Yield (%)	Ref.
1	Acetone (4)	2.2-Bis(difluoramino)propane (12)	90	4 <i>a</i>
2	Cyclopentanone (5)	1,1-Bis(difluoramino)cyclopentane (13)	80	4a
3	Cyclohexanone (6)	1,1-Bis(difluoramino)cyclohexane (14)	80	4a
4	Cyclohexa-1,4-dione (7)	1,1,4,4-Tetrakis(difluoramino)cyclohexane (15)	85	4a
5	N-Acetylpiperidinone (8)	4,4-Bis(difluoramino)- <i>N</i> -acetylpiperidine (16)	85	14
6	9a $(n = 1)$	17a	ab	_
7	9b $(n = 9)$	17b	80 ^b	_
8	9c $(n = 10)$	17c	85 ^b	_
9	10	18	90 ^b	_
10	11a (R = H)	20a	0^c	_
11	$11b(R = CH_3)$	20b	0^c	_

oxide or distillation, respectively. The given yields are based on ¹H NMR data of the crude product containing only the *gem*bis(difluoramine) (**B**) and triphenylmethanol. A variety of known (entries 1–5) as well as new (entries 6–11) difluoramine derivatives was prepared in good to excellent yields (Table 1).



Although **17a** could not yet be isolated on this small scale, it represents the smallest alicyclic geminal bis(difluoramine) so far characterized. The bisketone **11a** (**11b**) could only be converted to the bridged hemiaminal **19a** (**19b**); neither a larger excess of **3** nor extended reaction time furnished the tetrakis(difluoramine) **20a** (**20b**).

Mechanistically, the reactions involve cleavage of 3 to the triphenylmethyl cation and difluoramine. The latter or the in situ product, difluorsulfamic acid, reacts with the ketone A to provide gem-bis(difluoramine) B. It is noteworthy that our reaction did not proceed in conc. H₂SO₄, although acidolysis of 3 was observed spectroscopically: the ¹⁹F NMR signal of 3 disappeared and the trityl cation was observed in the ¹³C NMR spectrum. Also employing other strong acids such as FSO₃H, CF₃CO₂H, CF₃SO₃H or CF₃SO₃H/(CF₃SO₂)₂O resulted in similar behavior. In the case of FSO₃H/SbF₅ only decomposition of 3 was observed. We have never been able to observe free or protonated difluoramine.¹² It is also possible that under the reaction conditions SO₃ in oleum—besides being a water scavanger – might react with 3 to form F_2NSO_3H as the ultimate difluoraminating reagent (vide supra). The triphenylmethyl cation might also act like a Lewis acid catalyst, polarizing the carbonyl group (Mukayama type activation¹³).

In conclusion, the readily preparable, stable triphenylmethyldifluoramine (3) can be conveniently employed for the efficient conversion of ketones (A) under mild reaction conditions to the respective geminal bis(difluoramines) (B). This is the first difluoramination methodology that avoids the use of neat HNF₂ and therefore minimizes its potential hazards. Work is underway to further investigate the reaction mechanism, find alternative preparations of the reagent, and synthesize potential high-energy materials.

We gratefully acknowledge the financial support by the Office of Naval Research (Program Manager: Dr Judah Goldwasser), the Air Force Office of Scientific Research and the National Science Foundation.

CAUTION: *gem*-Bis(difluoramines) are potentially shocksensitive and thermolabile. They should be handled with care!

Notes and references

‡ *Crystal data*: the data were collected on a Bruker 3-circle platform diffractometer equipped with a SMART CCD detector with the χ-axis fixed at 54.74° from a fine-focus tube and an LT-3 apparatus for low temperature data collection: **3**, $C_{19}H_{15}F_2N$, $M_r = 295.32$, monoclinic, a = 10.3882(12), b = 9.8259(11), c = 14.6065(16) Å, $\beta = 92.402(2)^\circ$, V = 1489.6(3) Å³, T = 213(2) K, space group P_{21}/n , Z = 4, $D_c = 1.317$ Mg m⁻³, μ (Mo-Kα) = 0.094 mm⁻¹, 9649 reflections mesuared, 3035 unique ($R_{int} = 0.0321$). The final *wR2* was 0.1098 (all data). CCDC reference number 188416. See http://www.rsc.org/suppdata/cc/b2/b203811k/ for crystallographic data in CIF or other electronic format.

- W. Hanefeld, N,N-Dihalogen-amine, ed. D. Klaman in Methoden der Organischen Chemie (Houben-Weyl) Nitrogen Compounds, Vol. E16a/ Part 2, Georg-Thieme Verlag, Stuttgart, New York, 1990, pp. 893–925 and references therein.
- 2 (a) R. D. Chapman, M. F. Welker and C. B. Kreutzberger, J. Org. Chem., 1998, **63**, 1566; (b) R. D. Chapman, R. D. Gilardi, M. F. Welker and C. B. Kreutzberger, J. Org. Chem., 1999, **64**, 960.
- 3 T. Axenrod, X.-P. Guan, J. Sun, L. Qi, R. D. Chapman and R. D. Gilardi, *Tetrahedron Lett.*, 2001, **42**, 2621.
- 4 (a) K. Baum, J. Amer. Chem. Soc., 1968, 90, 7083; (b) W. H. Graham and J. P. Freeman, J. Org. Chem., 1969, 34, 2589; (c) A. V. Fokin, Y. M. Kosyrev, V. A. Makarov and N. P. Novoselov, Dokl. Akad. Nauk SSSR, 1969, 186, 112.
- 5 E. W. Lawless and I. C. Smith, *Inorganic High Energy Oxidizers* (Synthesis, Structure and Properties), Marcel Dekker Inc., New York, NY, USA, 1968, p. 69.
- 6 J. P. Freeman, A. Kennedy and C. B. Colburn, J. Am. Chem. Soc., 1960, 82, 5304.
- 7 (a) V. Grakauskas and K. Baum, J. Am. Chem. Soc., 1970, 92, 2096; (b)
 R. E. Banks, R. N. Hazeldine and J. P. Lalu, J. Chem. Soc. C, 1966, 1514.
- 8 (a) V. Grakauskas and K. Baum, J. Am. Chem. Soc., 1969, 91, 1679; (b)
 J. M. J. Lambert and G. M. Alain, (Societe Nationale des Poudres et Explosives), 1973, DE 2320119; (c) D. L. Klapotek and G. B. Hobrock, Inorg. Chem., 1967, 6, 1751.
- 9 (a) R. C. Petry and J. P. Freeman, J. Am. Chem. Soc., 1961, 83, 3912; (b)
 W. H. Graham and C. O. Parker, J. Org. Chem., 1963, 28, 850.
- 10 J. R. Surles, C. I. Bumgardner and J. Bordner, J. Fluorine Chem., 1975, 5, 467.
- 11 C. Gidwell and G. Furgeson, Acta Crystallogr., Sect. C, 1994, 50, 924.
- 12 K. O. Christe, Inorg. Chem., 1975, 14, 2821.
- 13 S. Kobayashi, M. Murakami and T. Mukaiyama, *Chem. Lett.*, 1985, 1535.
- 14 J. Zhang, J. Oxley, J. Smith, C. Bedford and R. Chapman, J. Mass Spectrom., 2000, 35, 841.