


Redox Chain Reaction—Indole and Pyrrole Alkylation with Unactivated Secondary Alcohols**

Xinping Han and Jimmy Wu*

The direct alkylation of indoles with alcohols would be of considerable synthetic utility because of the wide availability of both starting materials. While this is readily accomplished with activated substrates such as allylic,^[1] benzylic,^[2] and propargylic alcohols,^[3] the use of unactivated compounds such as secondary aliphatic alcohols is exceedingly rare and, to this day, represents a formidable challenge. Two examples were reported between 1940 and 1950.^[4,5] Despite their limited scope, the unique mechanism by which they operate (i.e., disproportionative condensation), piqued our curiosity.^[6] These transformations yielded C-alkylated products, but only under harsh reaction conditions (i.e., U.O.P. Ni catalyst, 30–600 mol % Na or K alkoxides, 174–220 °C, up to 64 h). Their synthetic utility, with respect to functional-group tolerance and product stability, would be considerably improved if milder reaction conditions could be identified.

We hypothesized that the fundamental mechanistic features of the base-promoted disproportionative condensation could be achieved under milder reaction conditions by employing an appropriate Brønsted acid catalyst (Scheme 1). As an illustrative example, indole **1** and alcohol **2** are treated with 5 mol % of the sacrificial ketone **3** and an appropriate acid catalyst. The reaction is initiated by condensation of **1** with **3** to furnish the indolyl cation **4** which is converted into the undesired by-product **5** and the new ketone **6** through a Meerwein–Ponndorf–Verley (MPV) reduction with **2** (Oppenauer oxidation of *i*PrOH).^[7] Then **6** condenses with **1** to furnish a different indolyl cation **7** which is similarly reduced to the desired product **8**.^[8] For every molecule of product that is formed, a molecule of **6** is regenerated, thereby propagating the reaction. We suggest *redox chain reaction* as an appropriate moniker for this process because of the similarities it shares with radical chain reactions,^[9] and also to distinguish it from base-promoted disproportionative condensation.

In line with our ongoing interest in the reactivity of indolyl cations,^[10] we report herein the successful application of the redox chain reaction to the alkylation of indole and pyrrole derivatives with unactivated secondary aliphatic alcohols. While catalytic variants are known,^[11,12] Meerwein–Pon-



Scheme 1. Acid-catalyzed redox chain reaction.

dorf–Verley–Oppenauer (MPVO) reactions typically require stoichiometric amounts of Lewis acids. To the best of our knowledge, this work is also likely to be the first instance of an MPVO process catalyzed by a Brønsted acid.^[13,14]



At the onset, we set out to identify the most efficient initiator and Brønsted acid catalyst, as well as their optimal loadings. After extensive experimentation (see the Supporting Information), we determined that 5 mol % of 2-methoxyacetophenone (**3a**) as an initiator and 10 mol % of TfOH in toluene at 100 °C provided the highest yield of **8ad** (88 %) for the reaction between *N*-benzylindole (**1a**) and *i*PrOH (**2d**; Scheme 2). Cyclohexanone and acetophenone (both 5 mol %) were also effective initiators, but because their by-products (corresponding to **5**) sometimes coeluted with the desired products on silica gel, we decided to adopt **3a** as the standard initiator. Importantly, control reactions in which either **3a** or TfOH was omitted resulted in no detectable product.

With the optimized reaction conditions in hand, we proceeded to explore the scope of the methodology. As is evident in Scheme 2, the reaction is tolerant to a variety of functional groups including esters, amides, nitro, nitriles, and ethers. The aminoalcohol **2g** can also be used, but in this case stoichiometric TfOH was required since up to 1 equivalent of acid could be consumed by the basic nitrogen center.

[*] X. Han, Prof. Dr. J. Wu
Department of Chemistry, Dartmouth College
6128 Burke Laboratories, Hanover, NH (USA)
E-mail: jimmy.wu@dartmouth.edu

[**] J.W. acknowledges financial support provided by Dartmouth College. We thank Dr. Richard J. Staples (Michigan State University, USA) for assistance with X-ray crystallography.

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/anie.201209810>.




Scheme 2. Scope of C3 alkylation of indoles. [a] 1.0 equiv alcohol, 1.1 equiv TFOH. [b] 2.0 equiv alcohol, 1.0 equiv TFOH, 10 mol% **3a**. [c] 2.0 equiv alcohol, 5 mol% TFOH, 10 mol% **3a**. Bn = benzyl, Tf = trifluoromethanesulfonyl, TBDPS = *tert*-butyldiphenylsilyl.

N-methylindole (**1e**) as well as 2-methyl-*N*-benzylindole (**1d**) are both compatible, thus producing **8dd** and **8eb** in good yields. The use of NH indole as the nucleophile resulted in no reaction.

As a testament to the improved efficiency of the redox chain reaction as compared to disproportionative condensation, we highlight the reaction between **1a** and cyclohexanol (**2b**) which occurs within only 1 hour at 100 °C to provide **8ab** in 93% yield upon isolation. The formation of three other products **8ac**, **8eb**, and **8an** (see Scheme 3) was also complete within 1 hour. In contrast, the alkylation of indole with cyclohexanol utilizing base-promoted disproportionative condensation requires 64 hours at 174–185 °C and furnishes the desired product in only 54% yield.^[4a] The use of primary

alcohols resulted in the formation of only **5** (< 5%). It is likely that the attenuated reactivity of primary alcohols as hydride donors, relative to secondary alcohols, precludes efficient propagation of the redox chain reaction.


Excellent levels of diastereoselectivity can be realized using alcohols with pre-existing stereocenters. As illustrated in Scheme 3, when we attempted the late-stage indoloylation of (+)-dihydrocholesterol (**2n**) with **1a**, the expected product **8an** was isolated in 81% yield as a single diastereomer



Scheme 3. Diastereoselective alkylation reactions. [a] Standard reaction conditions as described in Scheme 2. [b] Relative stereochemistry was confirmed by single-crystal X-ray analysis.^[17] [c] Confirmed by comparison of *J*_{HH} values to **8an**.


(stereochemical configuration was confirmed with single-crystal X-ray analysis). With *cis*-4-methylcyclohexanol (**2o**), high diastereoselectivity (**8ao**, 10:1 d.r.) was also achieved. Axial delivery^[15] of hydride to the respective indolyl carbocationic intermediates correctly predicts the observed selectivities for **8an** and **8ao**. Alkylation using **2p** (mixture *cis/trans*) furnished a 2:1 mixture of **8ap**. Curiously, alkylation with the 2-substituted cyclohexanol **2q** (mixture *cis/trans*) generated mainly the rearranged product **8aq**. By-products derived from 1,2-hydride shifts were not detected.

We wondered whether or not enantioenriched secondary alcohols would deliver hydride to the prochiral indolyl cation intermediates in a stereospecific fashion. Unfortunately, the reaction between **1a** and (*R*)-(-)-**2e** generated **8ae** as a racemate [Eq. (1)].^[16]



This methodology is also compatible with pyrrole nucleophiles (**9a-b**). Under the optimized reaction conditions, we


obtained good yields of the anticipated C2-substituted pyrroles **10a–f** (Scheme 4). Importantly, we did not detect any di- or polysubstituted products. We hypothesize that for steric reasons, after the first alkylation, the benzyl group is directed away from the newly installed C2 substituent and towards C5. Substantial nonbonding interactions experienced



Scheme 4. Scope of C2 alkylation of pyrroles. [a] Standard reaction conditions as described in Scheme 2. [b] 1.0 equiv alcohol, 1.1 equiv TfOH. [c] 2.0 equiv alcohol, 20 mol% TfOH, 10 mol% **3a**.

by electrophiles approaching C5 of **10** would disfavor a second alkylation event. For **9b**, we believe that alkylation takes place proximal to the C3 methyl group because C2 is more electron rich than C5. Furan and thiophene were not suitable nucleophiles as they resulted in either decomposition or no reaction, respectively.

Preliminary mechanistic studies are depicted in Scheme 5. The alkylation of **1a** with deuterated cyclohexanol **11** generated a 92:8 ratio of the isotopic isomers **14** and **15**. 1,4-Reduction would lead exclusively to **14**. Alternatively, 1,2-reduction and subsequent TfOH-promoted tautomerization/rearomatization would furnish **15**. Based on the 92:8 preference for the formation of **14**, we conclude that 1,4-reduction is



Scheme 5. Deuterium-labeling studies.

the predominant pathway. These results also provide evidence in support of the intermediacy of indolyl carbocation **12**.

In conclusion, we have developed the Brønsted acid catalyzed redox chain reaction as it pertains to the alkylation of indole and pyrrole derivatives with unactivated secondary alcohols. Broad functional-group tolerance has been demonstrated. Deuterium-labeling studies support the intermediacy of an indolyl carbocation which undergoes MPVO conjugate reduction as a likely reaction mechanism.

We anticipate that it might be possible to apply the concept of the redox chain reaction to the alkylation of other nucleophiles such as nonbasic nitrogen atoms, enamines, and electron-rich benzene derivatives. We are also exploring the possibility of using chiral Lewis and Brønsted acids to induce asymmetry.

Received: December 7, 2012

Revised: February 5, 2013

Published online: ■■■■■, ■■■■■

Keywords: alcohols · alkylation · heterocycles · redox chemistry · synthetic methods

- [1] a) B. M. Trost, J. Quancard, *J. Am. Chem. Soc.* **2006**, *128*, 6314; b) M. Kimura, M. Futamata, R. Mukai, Y. Tamaru, *J. Am. Chem. Soc.* **2005**, *127*, 4592; c) I. Usui, S. Schmidt, M. Keller, B. Breit, *Org. Lett.* **2008**, *10*, 1207; d) A. B. Zaitsev, S. Gruber, P. A. Plüss, P. S. Pregosin, L. F. Veiros, M. Wörle, *J. Am. Chem. Soc.* **2008**, *130*, 11604; e) M. Yasuda, T. Somyo, A. Baba, *Angew. Chem.* **2006**, *118*, 807; *Angew. Chem. Int. Ed.* **2006**, *45*, 793; f) K. J. Henry, Jr., P. A. Grieco, *J. Chem. Soc. Chem. Commun.* **1993**, 510; g) J. S. Yadav, B. V. Reddy, S. Aravind, G. G. K. S. N. Kumar, A. S. Reddy, *Tetrahedron Lett.* **2007**, *48*, 6117; h) J. Le Bras, J. Muzart, *Tetrahedron* **2007**, *63*, 7942; i) J. S. Yadav, B. V. S. Reddy, P. M. Reddy, C. Srinivas, *Tetrahedron Lett.* **2002**, *43*, 5185; j) D. Prajapati, M. Gohain, B. J. Gogoi, *Tetrahedron Lett.* **2006**, *47*, 3535.
- [2] a) M. Rueping, B. J. Nachtsheim, W. Ieawsuwan, *Adv. Synth. Catal.* **2006**, *348*, 1033; b) P. Vicennati, P. G. Cozzi, *Eur. J. Org. Chem.* **2007**, 2248; c) S. Shirakawa, S. Kobayashi, *Org. Lett.* **2007**, *9*, 311; d) J. Y. L. Chung, D. Mancheno, P. G. Dormer, N. Variankaval, R. G. Ball, N. N. Tsou, *Org. Lett.* **2008**, *10*, 3037.
- [3] a) J. S. Yadav, B. V. S. Reddy, K. V. R. Rao, G. G. K. S. Kumar, *Tetrahedron Lett.* **2007**, *48*, 5573; b) P. Srihari, D. C. Bhunia, P. Sreedhar, S. S. Mandal, J. S. S. R. Reddy, J. S. Yadav, *Tetrahedron Lett.* **2007**, *48*, 8120; c) R. Sanz, D. Miguel, J. M. Álvarez-Gutiérrez, F. Rodríguez, *Synlett* **2008**, 975.
- [4] a) E. F. Pratt, L. W. Botimer, *J. Am. Chem. Soc.* **1957**, *79*, 5248; b) R. H. Cornforth, R. Robinson, *J. Chem. Soc.* **1942**, 680.
- [5] Grigg and co-workers reported the use of iridium-catalyzed “hydrogen borrowing” to alkylate indole at C3 with primary benzylic and heterobenzylic alcohols. Two primary aliphatic alcohols were reported (three entries). In these instances, 0.2–2.0 equiv of KOH was required, the reaction time was 24–48 h at 110°C, and yields were 35, 53, and 74%. See: S. Whitney, R. Grigg, A. Derrick, A. Keep, *Org. Lett.* **2007**, *9*, 3299.
- [6] For other reactions which may proceed by base-promoted disproportionative condensation, see: a) E. F. Pratt, E. J. Frazza, *J. Am. Chem. Soc.* **1954**, *76*, 6174; b) K. L. Schoen, E. I. Becker, *J. Am. Chem. Soc.* **1955**, *77*, 6030; c) R. H. Cornforth, R. Robinson, *J. Chem. Soc.* **1942**, 682.

- [7] a) C. F. de Graauw, J. A. Peters, H. van Bekkum, J. Huskens, *Synthesis* **1994**, 1007; b) C. R. Graves, K. A. Scheidt, S. T. Nguyen, *Org. Lett.* **2006**, 8, 1229.
- [8] For the reductive alkylation between indole and ketones/aldehydes with Et₃SiH or H₂, see: a) J. R. Rizzo, C. A. Alt, T. Y. Zhang, *Tetrahedron Lett.* **2008**, 49, 6749; b) L.-L. Cao, D.-S. Wang, G.-F. Jiang, Y.-G. Zhou, *Tetrahedron Lett.* **2011**, 52, 2837.
- [9] F. A. Carey, R. J. Sundberg, *Advanced Organic Chemistry Part A: Structure and Mechanisms*, Springer Science + Business Media, New York, **2007**, pp. 965–1071.
- [10] X. Han, H. Li, R. P. Hughes, J. Wu, *Angew. Chem.* **2012**, 124, 10536; *Angew. Chem. Int. Ed.* **2012**, 51, 10390.
- [11] For lead references of MPVO reactions which are catalytic in Lewis acid, see: a) K.-J. Haack, S. Hashiguchi, A. Fujii, T. Ikariya, R. Noyori, *Angew. Chem.* **1997**, 109, 297; *Angew. Chem. Int. Ed. Engl.* **1997**, 36, 285; b) E. J. Campbell, H. Zhou, S. T. Nguyen, *Angew. Chem.* **2002**, 114, 1062; *Angew. Chem. Int. Ed.* **2002**, 41, 1020; c) D. A. Evans, S. G. Nelson, M. R. Gagné, A. R. Muci, *J. Am. Chem. Soc.* **1993**, 115, 9800.
- [12] For lead references on the intermolecular cross-Tishchenko reaction, see: a) L. Cronin, F. Manoni, C. J. O'Connor, S. J. Connon, *Angew. Chem.* **2010**, 122, 3109; *Angew. Chem. Int. Ed.* **2010**, 49, 3045; b) S. P. Curran, S. J. Connon, *Angew. Chem.* **2012**, 124, 11024; *Angew. Chem. Int. Ed.* **2012**, 51, 10866.
- [13] There is one example of an uncatalyzed MPVO reaction but requires 300 °C under supercritical conditions. See: L. Sominsky, E. Rozental, H. Gottlieb, A. Gedanken, S. Hoz, *J. Org. Chem.* **2004**, 69, 1492.
- [14] Brønsted acid additives to MPVO reactions promoted by aluminum-based reagents can enhance reaction rate. See: a) R. Kow, R. Nygren, M. W. Rathke, *J. Org. Chem.* **1977**, 42, 826; b) K. G. Akamanchi, N. R. Varalakshmy, *Tetrahedron Lett.* **1995**, 36, 3571; c) K. G. Akamanchi, N. R. Varalakshmy, B. A. Chaudhari, *Synlett* **1997**, 371.
- [15] a) B. W. Gung, *Tetrahedron* **1996**, 52, 5263; b) Y.-D. Wu, K. N. Houk, *J. Am. Chem. Soc.* **1987**, 109, 906; c) Y.-D. Wu, K. N. Houk, *J. Am. Chem. Soc.* **1987**, 109, 908.
- [16] The *ee* value determination performed by Lotus Separations, Princeton, NJ, USA.
- [17] CCDC 929479 (**8an**) contains the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif.


Communications



Synthetic Methods

X. Han, J. Wu* ————— ■■■■-■■■■

Redox Chain Reaction—Indole and Pyrrole Alkylation with Unactivated Secondary Alcohols



Secondary role: Indole and pyrrole derivatives are alkylated with unactivated secondary aliphatic alcohols by a Brønsted acid-catalyzed redox chain reaction mechanism. Broad functional-

group tolerance has been demonstrated and preliminary studies suggest that 1,4-reduction of a putative indolyl carbocation is the dominant mechanistic pathway.