

Synthesis and Antibacterial Activity of a Series of Heterocyclic Sulfonate Ligands and Their Metal Complexes

FENG XIAO-JUN¹, DAI BI-SHENG¹, SU CHENG-BAI¹, ZHENG-ZHENG¹, WANG LIN² and DONG HUA-ZE^{2,*}

¹Food and Drug Department, Qingyuan Polytechnic, Qingyuan 511510, P.R. China

²Department of Chemistry and Chemical Engineering, Hefei Normal University, Hefei 230061, P.R. China

*Corresponding author: E-mail: dapdong@163.com

(Received: 20 December 2011;

Accepted: 12 October 2012)

AJC-12273

A series of heterocyclic sulfonate ligands and their barium(II), silver(I), manganese(II), zinc(II) and cobalt(II) coordination compounds have been synthesized. Through antibacterial activities experiments of these complex compounds, metal salts as well as ligands, the rational conclusion is arrived that antibacterial activities of these compounds come from metal ions.

Key Words: Heterocyclic sulfonate ligand, Antibacterial activity, Coordination complex.

INTRODUCTION

Though organic sulfonate and their metal complexes have been studied by chemists¹⁻⁸, in pharmacy, the antibacterial activities of these compounds are explored scarcely. To perfect coordination and broad-spectrum antibacterial ability, sulfonic acid is modified with pyrimidine.


EXPERIMENTAL

All the solvents and chemicals were used without further purification. Infrared spectra (4000-400 cm⁻¹) were recorded with a Bruker Vector 22 FTIR spectrophotometer on KBr disks. Elemental analyses were done by using a Perkin-Elmer 1400C analyzer. The microbial strains were from Agricultural Culture Collection of China (ACCC).

Synthesis of L₁₋₃: The series of ligands{4-pyridin-2-yl-pyrimidine-2-sulfonate (**L₁**), 4-pyridin-3-yl-pyrimidine-2-sulfonate (**L₂**), 4-pyridin-4-yl-pyrimidine-2-sulfonate (**L₃**)} were synthesized according to the literature method⁹⁻¹¹ with synthetic route below (Scheme-I).

Synthesis of complexes: [BaL₁(NO₃)(H₂O)]_n (**1**), {[Ba(L₁)₂(H₂O)]·2H₂O}_n (**2**), [BaL₂(H₂O)(ClO₄)]_n (**3**), [Ag₃(L₁)₃(H₂O)₂]_n (**4**), [Ag₃(L₂)₃(H₂O)₂]_n (**5**), [Ag₃(L₁)₃(H₂O)₂]_n (**6**), [Mn(L₁)₂·H₂O] (**7**), [Mn(L₂)₂·H₂O] (**8**), [Mn(L₃)₂·4H₂O]·2H₂O (**9**), [Zn(L₁)₂·H₂O] (**10**), [Zn(L₂)₂·H₂O] (**11**), [Zn(L₃)₂·4H₂O]·2H₂O (**12**), [Co(L₁)₂·H₂O] (**13**), [Co(L₂)₂·H₂O] (**14**), [Co(L₃)₂·4H₂O]·2H₂O (**15**).

The single crystals of complexes **1-3** were used for detection of antibacterial activity directly (CCDC: **1**, 760305;



Scheme-I: Synthetic route of title ligands

2, 760306; **3**, 760307). Synthesis, characterization, crystal data and structure were reported in earlier thesis¹².

NaL₁, NaL₂, NaL₃ (26.0 mg, 0.1 mmol) were added, respectively in water (5 mL), then triplicate solutions of AgNO₃ (8.5 mg, 0.05 mmol) in acetonitrile (5 mL) were very slowly dropped in. Crystal products (**4**, **5**, **6**) formed after two-three weeks standing in darkness. The structures of **4**, **5** were confirmed by Liu and Hu¹³.

4: Yield 78 %. Anal. calcd. (%) for Ag₃(L₁)₃(H₂O)₂ {[Ag₃(C₉H₆N₃O₃S)₃(H₂O)₂]}: C, 30.35; H, 2.08; N, 11.80. Found. (%): C, 30.34; H, 2.10; N, 11.77. Main IR absorption (KBr, ν_{max}, cm⁻¹): 3431 (m), 1615 (m), 1578 (s), 1533 (m), 1496 (m), 1385 (m), 1225 (s), 1197 (s), 1043 (s).

TABLE-1
 DETECTION OF ANTIBACTERIAL ACTIVITY (“+” IS POSITIVE RESULT, “-” IS NEGATIVE RESULT)

Compound	Bacteria														
	<i>Escherichia coli</i> (ATCC01555)					<i>Bacillus subtilis</i> (ATCC01430)					<i>Staphylococcus aureus</i> (ATCC01331)				
L₁	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
L₂	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
L₃	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
BaCl ₂	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
3	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
AgNO ₃	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
4	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
5	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
6	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
MnCl ₂	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
7	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
8	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
9	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
ZnCl ₂	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
10	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
11	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
12	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
CoCl ₂	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
13	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
14	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
15	-	-	-	-	-	+	+	+	+	+	+	+	+	+	+

5: Yield 64 %. Anal. calcd. (%) for Ag₃(L₂)₃(H₂O)₂ {[Ag₃(C₉H₆N₃O₃S)₃(H₂O)₂]}: C, 30.35; H, 2.08; N, 11.80. Found (%): C, 30.33; H, 2.10; N, 11.83. Main IR absorption (KBr, $\nu_{\max}$, cm⁻¹): 3431 (m), 1617 (m), 1573 (s), 1530 (m), 1495 (m), 1386 (m), 1225 (s), 1197 (s), 1044 (s).

6: Yield 71 %. Anal. calcd. (%) for Ag₃(L₃)₃(H₂O)₂ {[Ag₃(C₉H₆N₃O₃S)₃(H₂O)₂]}: C, 30.35; H, 2.08; N, 11.80. Found. (%): C, 30.37; H, 2.07; N, 11.78. Main IR absorption (KBr, $\nu_{\max}$, cm⁻¹): 3432 (m), 1615 (m), 1574 (s), 1533 (m), 1495 (m), 1381 (m), 1222 (s), 1195 (s), 1041 (s).

The preparation of complexes (**7-15**) with similar way, the process as followed: the ligand (**L₁-L₃**, 0.0472 g, 0.2 mmol) in warm water (15 mL) and an aqueous solution (10 mL) of metal salt (0.1 mmol) [MnCl₂·4H₂O; Zn(ClO₄)₂·6H₂O; Co(ClO₄)₂·6H₂O] were mixed. The mixture was stirred for 1 h, the crystals were obtained in filtrate when the solution cooled to room temperature. CCDC: 9, 681493; 12, 681494; 15, 681495. The general formulas of these complexes are [M(L)₂]-H₂O (L = **L₁**, **L₂**)¹⁴ and [M(L)₂·4H₂O]·2H₂O (L = **L₃**)¹⁵. The structures of complexes **7**, **10**, **13** were reported by Dong *et al.*¹⁴.

8: Yield 86 %. Anal. calcd. (%) for C₁₈H₁₄MnN₆O₇S₂ {[Mn(L₂)₂]-H₂O}: C, 40.99; H, 2.29; N, 15.93. Found. (%): C, 40.97; H, 2.28; N, 15.89. Main IR absorption (KBr, $\nu_{\max}$, cm⁻¹): 3442 (vs), 1616 (w), 1577 (s), 1534 (m), 1430 (m), 1418 (m), 1370 (w), 1249 (vs), 1197 (s), 1034 (s).

11: Yield 63 %. Anal. calcd. (%) for C₁₈H₁₄ZnN₆O₇S₂ {[Zn(L₂)₂]-H₂O}: C, 38.89; H, 2.54; N, 15.12. Found. (%): C, 38.85; H, 2.53; N, 15.10. Main IR absorption (KBr, $\nu_{\max}$, cm⁻¹): 3445 (vs), 1614 (w), 1579(s), 1533(m), 1432(m), 1416(m), 1374(w), 1246(vs), 1199(s), 1034(s).

14: Yield 62 %. Anal. calcd. (%) for C₁₈H₁₄CoN₆O₇S₂ {[Co(L₂)₂]-H₂O}: C, 39.35; H, 2.57; N, 15.30. Found. (%): C, 39.33; H, 7.59; N, 15.32. Main IR absorption (KBr, $\nu_{\max}$, cm⁻¹):

3448 (vs), 1617 (w), 1574 (s), 1531 (m), 1434 (m), 1420 (m), 1372 (w), 1245 (vs), 1196 (s), 1032 (s).

Detection of antibacterial activity: After materials for detection of antibacterial activity autoclaved, each petri dish was infused proper nutrient agar *ca.* 40 °C. When the nutrient agar was cooled to room temperature, a solution of bacterial and a piece of circular paper (0.5 cm in diameter) carrying with sample were placed on it in order. Then petri dish were put into electro-thermal incubator at 36 °C for 24 h. The control group (n = 5) was established with every compound. At the same time, a blank control group (n = 5) without any sample was cultured.

As good water-solubility of **L₁₋₃** and inorganic compounds, their solutions (0.001 g/mL) were used in detecting antibacterial activity. The rest of samples (**1-15**) were performed in solid on watery papers for their poor solubility. The positive result is that the inhibition zone was present and *vice versa*. The blank control group assumed negative result.

RESULTS AND DISCUSSION

Three kinds of bacterias are not inhibited by ligands (**L₁₋₃**) and anions (Cl⁻, ClO₄⁻, NO₃⁻). Complexes and corresponding metal ions exhibits similar antibacterial activity. Enough metal ionic concentration is the cause of antimicrobial activity expect two kinds complexes that are Co(II)-**L₃** and Mn(II)-**L₁₋₃**. It is possible that dissociation constant of this two kinds metal complexes is too low to inhibit increase of microbe. These coordination compounds are potential sustained-release antibacterial drugs for their low dissociation constant.

ACKNOWLEDGEMENTS

This work was financially supported by Bureau of Science and Technology, Qingyuan (Project NO.2010A006, Project

No. 2009B057). It is also supported by the National Science Foundation of China (No. 21101053), Program for Excellent Young Talents in University of Anhui Province (2011SQRL128), Anhui Provincial Natural Science Research Project (kj20112311) and Key Disciplines Foundation of Hefei Normal University.

REFERENCES

1. B.J. Guanderman and P.J. Squattrito, *Inorg. Chem.*, **33**, 2924 (1994).
2. E.J. Kosnic, E.L. McClymont, R.A. Hodder and P.J. Squattrito, *Inorg. Chim. Acta*, **210**, 143 (1992).
3. A.J. Shubnell, E.J. Kosnic and P.J. Squattrito, *Inorg. Chim. Acta*, **216**, 101 (1994).
4. S. Marder, J. Peny and C. Yakymyshyn, *Chem. Mater.*, **6**, 1137 (1994).
5. V. Russell, M. Eter and M. Ward, *Chem. Mater.*, **6**, 1206 (1994).
6. J. Cai, C.H. Chen, C.Z. Liao, X.L. Feng and X.M. Chen, *Acta Crystallogr.*, **B57**, 520 (2001).
7. A.P. Côté and G.K.H. Shimizu, *Coord. Chem. Rev.*, **245**, 49 (2003).
8. J.W. Cai, *Coord. Chem. Rev.*, **248**, 1061 (2004).
9. M.M. Bio, F. Xu, M. Waters, J.M. Williams, K.A. Savary, C.J. Cowden, C. Yang, E. Buck, Z.J. Song, D.M. Tschaen, R.P. Volante, R.A. Reamer and E.J.J. Grabowski, *J. Org. Chem.*, **69**, 6257 (2004).
10. H.B. Zhu, H.Z. Dong, W. Huang and S.H. Gou, *J. Mol. Struct.*, **831**, 55 (2007).
11. X.B. Fang, H.Z. Dong and D.B. Tian, *Chin. J. Inorg. Chem.*, **25**, 47 (2009).
12. H.-Z. Dong, J.-H. Bi, H.-B. Zhu and S.-H. Gou, *J. Mol. Struct.*, **975**, 173 (2010).
13. X. Liu and D. Hu, *Acta Cryst.*, **E67**, m1067 (2011).
14. H.B. Zhu, H.Z. Dong, W. Huang and S.H. Gou, *J. Mol. Struct.*, **831**, 55 (2007).
15. X.-B. Fang, H.-Z. Dong and D.-B. Tian, *Chin. J. Inorg. Chem.*, **25**, 47 (2009).