

A Stereoselective Total Synthesis of (9*S*)-9-Dihydroerythronolide A *via* Coupling between the Right-Half (C1—C6) Aldehyde and the Left-Half (C7—C15) Sulfoxide^{1,2)}

Hitoshi TONE, Takao NISHI, Yuji OIKAWA, Masataka HIKOTA and Osamu YONEMITSU*

Faculty of Pharmaceutical Sciences, Hokkaido University, Kita-12, Nishi-6, Kitaku, Sapporo 060, Japan. Received August 26, 1988

As part of a study directed at the total synthesis of (9*S*)-9-dihydroerythronolide A, the C7—C15 sulfoxide, (2*S*,3*R*,4*S*,5*R*,6*R*,7*R*)-3,5-isopropylidenedioxy-7-(4-methoxybenzyloxy)-6-methoxymethoxy-1-phenylsulfinyl-2,4,6-trimethylnonane, was coupled with the C1—C6 aldehyde, (2*R*,3*S*,4*S*,5*R*)-6-*tert*-butyldiphenylsilyloxy-3,5-dimethyl-2,4-isopropylidenedioxyhexanal, to give the C1—C15 hydroxysulfoxide, which was converted to the seco-acid *via* a stereocontrolled methylation at the C6 position. Macrocyclization of the seco-acid by Yamaguchi's method gave the 14-membered lactone, which was converted to (9*S*)-9-dihydroerythronolide A.

Keywords macrolide antibiotic; erythromycin A; aglycone; erythronolide A; stereoselective synthesis; sulfone coupling; protecting group; seco-acid; macrolactonization; high dilution

As part of our continuing study directed toward the stereoselective synthesis of the well-known macrolide antibiotic erythromycin A (1),^{2,3)} in the preceding paper¹⁾ we reported the synthesis of the enone (4) having the whole carbon skeleton of (9*S*)-9-dihydroerythronolide A (3) by the Wittig-Horner coupling⁴⁾ between the aldehyde (5) and the β -ketophosphonate (6). However, the yield of 4 was too low to complete the synthesis of 3, although improvements are now in progress. In the present paper,²⁾ we report another approach *via* coupling between a C7—C15 sulfoxide (7) and a C1—C6 carbonyl compound (8),⁵⁾ leading to completion of the stereoselective total synthesis of 3. Actually, coupling of 9 and 10 followed by introduction of

the final methyl group at C6 and Yamaguchi's macrolactonization⁶⁾ gave 3.

Results and Discussion

Synthesis of the C7—C15 Sulfoxide (9), the C1—C6 Aldehyde (10), and the C1—C6 Methyl Ketone (11) Conversion of the C7-alcohol (12)^{1,7)} into the sulfide (14) was readily carried out *via* the tosylate (13).^{5,8)} Treatment of 12 with *p*-toluenesulfonyl chloride followed by sodium thiophenoxide gave 14 in high yield. Another method involving treatment of 12 with diphenyl disulfide and tributylphosphine⁹⁾ gave only poor results. Periodate oxidation of 14 gave the expected C7—C15 sulfoxide (9) as a






Chart 1



(i) TsCl, Et₃N, DMAP, CH₂Cl₂, reflux, 23 h; (ii) PhSNa, EtOH-DME (1:1), reflux, 2 h; (iii) NaIO₄, MeOH-H₂O, room temperature, 8.5 h.

Chart 2



(i) 1,3-Dithiane, *n*-BuLi, THF, -80°C, 40 min; (ii) 4N HCl, THF, room temperature, 11 h; (iii) TBDPSCl, imidazole, CH₂Cl₂, room temperature, 2 h; (iv) CH₂=CMe(OMe), PPTS, CH₂Cl₂, room temperature, 2.5 h; (v) MeI, NaHCO₃, 90% MeCN, 70°C, 1.5 h or NBS, 2,6-lutidine, 80% MeCN, room temperature, 1 h (71%); (vi) a) MeMgI, Et₂O, -15°C, 1 h (89%), b) (COCl)₂, DMSO, Et₃N (86%).

Chart 3

1:1 stereoisomeric mixture with regard to the sulfoxide position.

The diol (**15**), derived from D-glucose¹⁰ or methallyl alcohol,¹¹ was readily converted to the aldehyde (**16**),¹⁰ which was treated with the carbanion of 1,3-dithiane to give a 4.9:1 diastereoisomeric mixture of adducts mainly consisting of the non-chelation-controlled Cram adduct (**17**) in high yield. After purification of **17** by recrystallization, the isopropylidene protection of **17** was removed by treatment with 4N hydrochloric acid in tetrahydrofuran (THF) to give the triol (**18**), and selective protection of the primary alcohol of **18** with a *tert*-butyldiphenylsilyl (TBDPS) group gave **19**, the remaining diol of which was then protected as an acetonide to give **20**. Removal of the 1,3-dithiane protection of **20** proceeded rather smoothly by treatment with *N*-bromosuccinimide (NBS) in the presence of 2,6-lutidine,¹² but the yield of the expected C1—C6 aldehyde (**10**) was not so good. Treatment with methyl iodide in the presence of sodium hydrogencarbonate¹³ gave a better result.

The C1—C6 methyl ketone (**11**)⁵ was easily obtained from **10** by Grignard reaction followed by Swern oxidation.

Coupling between 9 and 10, and Total Synthesis of 3
Coupling between the sulfoxide (**9**) and the ketone (**11**) in the presence of lithium diisopropylamide (LDA)⁵ under several conditions was first examined, but the yield was always less than 20%. Therefore, the ketone (**11**) was replaced by the more reactive aldehyde (**10**), and the coupling product (**21**) was obtained in high yield as a mixture of four stereoisomers. Desulfurization with Raney nickel gave the alcohol (**22**) as a stereoisomeric mixture, which was subjected to Swern oxidation to give the ketone (**23**) as a single product.

Introduction of the final methyl group at the C6 position was achieved as follows. No reaction of **23** occurred with methylmagnesium iodide in ether, but when **23** was treated

with a large excess of methyllithium (MeLi) in ether at -85—-65°C, a chelation-controlled reaction proceeded quite smoothly to give a 4.9:1 stereoisomeric mixture in quantitative yield mainly consisting of the unexpected isomer (**25**; 83%), with the desired isomer (**24**) as the minor product (17%). However, when hexamethylphosphoramide (HMPA; 40 eq) was added to the above reaction system, the reaction changed clearly to a non-chelation-controlled reaction, and a 5.6:1 mixture of the desired Cram adduct (**24**; 73%) and its isomer (**25**; 13%) together with the recovered starting material (**23**; 9%) was isolated. The reaction of **23** with MeLi in THF gave a better result, that is, a 7.2:1 mixture mainly consisting of **24** (83%) was obtained. After purification by thin-layer chromatography (TLC), **24** was treated with methoxymethyl (MM) chloride to give quantitatively **26**, which was treated with fluoride anion and then oxidized at the resulting primary alcohol with Jones reagent to give the carboxylic acid (**27**). The *p*-methoxybenzyl (MPM) protective group of **27** was removed by catalytic hydrogenation over palladium charcoal or by oxidation with 2,3-dichloro-5,6-dicyanobenzoquinone (DDQ),¹⁴ and the seco-acid (**28**) required for macrolactonization was isolated in quantitative yield.

We were now ready to try macrolactonization. Two of Corey's methods using 4-*tert*-butyl-*N*-isopropyl-2-imidazolyl disulfide^{3a,15} and 2-pyridyl disulfide^{3b,16} were first applied, but unfortunately all attempts were unsuccessful. However, macrolactonization of **28** was achieved by Yamaguchi's method⁶ in the presence of a rather high concentration of 4-dimethylaminopyridine (DMAP). When 2 mM toluene solution of a mixed anhydride, prepared from **28** and 2,4,6-trichlorobenzoyl chloride in the presence of triethylamine in THF, was added very slowly to an equal volume of 50 mM toluene solution of 25 eq of DMAP under reflux over a period of 39 h, in spite of the fact that an axial methyl group of the 9,11-isopropylidene group hinders this




Chart 4

lactonization as pointed out by Stork and Rychnovsky,^{3e)} the macrolactonization proceeded gradually to give the expected lactone (**29**), though the yield was only 27%. In this high dilution macrolactonization, the yield of **29** was decisively dependent on the concentration of DMAP. When the final concentration of DMAP was 3 mM, no **29** was obtained. However, the yield increased with increasing concentration of DMAP, that is, yields of **29** at various final concentrations of DMAP were as follows: 0% at 3 mM, 13% at 6 mM, 14% at 9.5 mM, and 27% at 25 mM. All the protecting groups of **29** were removed with 50% acetic acid, and the title compound (**3**) was isolated in excellent yield. Compounds **3**, **28**, and **29** were identical with the respective authentic samples derived from natural erythromycin A in terms of infrared (IR), nuclear magnetic resonance (NMR) and mass spectra, and chromatographic mobilities.

Experimental

Physical data were measured as described in the previous paper.¹¹⁾

(2S,3R,4S,5R,6R,7R)-3,5-Isopropylidenedioxy-7-(4-methoxybenzyloxy)-6-methoxymethoxy-1-(4-toluenesulfonyloxy)-2,4,6-trimethylnonane (13) A solution of the alcohol (**12**) (327 mg, 0.721 mmol), Et₃N (0.90 ml, 6.48 mmol), DMAP (27 mg, 0.22 mmol), and *p*-toluenesulfonyl chloride (413 mg, 2.16 mmol) in CH₂Cl₂ (30 ml) was stirred under reflux for 23 h. It was then cooled, H₂O was added, and the mixture was stirred at room temperature for 3 h. The CH₂Cl₂ layer was washed with aqueous KHSO₄

and brine, dried (Na₂SO₄), and evaporated *in vacuo* to leave the tosylate (**13**) as a colorless solid (448 mg, 100%). Recrystallization from *n*-hexane gave pure **13** as colorless needles, mp 97–99°C. ¹H-NMR (CDCl₃) δ: 0.87 (3H, d, *J* = 7.0 Hz), 0.98 (3H, d, *J* = 7.0 Hz), 1.07 (3H, t, *J* = 7.5 Hz), 1.21 (6H, s), 1.29 (3H, s), 1.62 (1H, quintet, *J* = 7.5 Hz), 1.72–1.96 (3H, m), 2.45 (3H, s), 3.33 (1H, dd, *J* = 7.5, 2.0 Hz), 3.35 (3H, s), 3.45 (1H, dd, *J* = 8.0, 2.5 Hz), 3.80 (3H, s), 3.85 (1H, d, *J* = 4.5 Hz), 3.88 (1H, dd, *J* = 8.0, 2.5 Hz), 3.97 (1H, dd, *J* = 9.0, 8.0 Hz), 4.48, 4.61 (1H each, ABq, *J* = 11.0 Hz), 4.76, 4.98 (1H each, ABq, *J* = 7.0 Hz), 6.85 (2H, d, *J* = 9.0 Hz), 7.24 (2H, d, *J* = 9.0 Hz), 7.34 (2H, d, *J* = 8.0 Hz), 7.79 (2H, d, *J* = 8.0 Hz). MS *m/z* (relative intensity): 505 (M⁺ – 103, 0.1), 429 (0.7), 371 (7.9), 166 (13), 121 (100). [α]_D²⁵ –15.1° (*c* = 1.11, CHCl₃). Anal. Calcd for C₃₂H₄₈O₉S: C, 63.13; H, 7.95. Found: C, 62.94; H, 8.00.

(2S,3S,4S,5R,6R,7R)-3,5-Isopropylidenedioxy-7-(4-methoxybenzyloxy)-6-methoxymethoxy-1-phenylthio-2,4,6-trimethylnonane (14) NaH (60%, 80 mg, 2.0 mmol) was dissolved in EtOH (10 ml). After evolution of H₂ had ceased, thiophenol (206 μl, 2.0 mmol) was added, and the solution was stirred at room temperature for 10 min. A solution of **13** (304 mg, 0.5 mmol) in 1,2-dimethoxyethane (DME) (10 ml) was added, and the solution was refluxed under argon for 2 h, then cooled, Et₂O and H₂O were added. The Et₂O layer was washed with 1 N NaOH and brine, dried (Na₂SO₄), and evaporated *in vacuo*. The residue was chromatographed on a silica gel column with *n*-hexane–EtOAc (16:1) as the eluent to give **14** as a colorless oil (246 mg, 90%). ¹H-NMR (CDCl₃) δ: 0.98 (3H, d, *J* = 7.0 Hz), 1.01 (3H, d, *J* = 7.0 Hz), 1.07 (3H, t, *J* = 7.5 Hz), 1.28 (3H, s), 1.31 (3H, s), 1.34 (3H, s), 1.67 (1H, quintet, *J* = 7.5 Hz), 1.68–1.90 (3H, m), 2.85 (1H, dd, *J* = 13.0, 6.5 Hz), 2.99 (1H, dd, *J* = 13.0, 7.5 Hz), 3.46 (1H, dd, *J* = 8.5, 2.5 Hz), 3.79 (3H, s), 3.90 (1H, d, *J* = 4.0 Hz), 4.50, 4.61 (1H each, ABq, *J* = 11.0 Hz), 4.79, 5.00 (1H each, ABq, *J* = 7.0 Hz), 6.85 (2H, d, *J* = 9.0 Hz), 7.26 (2H, d, *J* = 9.0 Hz), 7.14–7.35 (5H, m). MS *m/z* (relative

intensity): 547 ($M^+ + 1$, 0.02), 546 (0.05), 457 (0.06), 443 (0.4), 368 (0.5), 348 (0.6), 309 (5.7), 221 (4.7), 121 (100). $[\alpha]_D^{25} = -24.6^\circ$ ($c = 2.95$, CHCl_3). Exact MS m/z Calcd for $\text{C}_{26}\text{H}_{35}\text{O}_4\text{S}$ ($M^+ - 103$): 443.2256. Found: 443.2263.

(2S,3R,4S,5R,6R,7R)-3,5-Isopropylidenedioxy-7-(4-methoxybenzyloxy)-6-methoxymethoxy-1-phenylsulfinyl-2,4,6-trimethylnonane (9) A solution of NaIO_4 (290 mg, 1.35 mmol) in H_2O (3 ml) was added to a stirred solution of **14** (246 mg, 0.451 mmol) in MeOH (20 ml) at room temperature. After 8.5 h, CH_2Cl_2 and H_2O were added, and the CH_2Cl_2 layer was dried (Na_2SO_4) and evaporated *in vacuo*. The residue was chromatographed on a silica gel column with CH_2Cl_2 as the eluent to give a 1:1 diastereoisomeric mixture of the sulfoxide (**9**) as a pale yellow oil (250 mg, 100%). $^1\text{H-NMR}$ (CDCl_3) δ : 1.01 (3H, d, $J = 7.0$ Hz), 1.08 (3H, t, $J = 6.5$ Hz), 1.08 (1.5H, d, $J = 6.5$ Hz), 1.17 (1.5H, d, $J = 6.5$ Hz), 1.24 (1.5H, s), 1.25 (1.5H, s), 1.29 (1.5H, s), 1.30 (1.5H, s), 1.32 (1.5H, s), 1.39 (1.5H, s), 1.50–1.75 (1H, m), 1.74–1.96 (2H, m), 2.10–2.34 (1H, m), 2.64–2.91 (2H, m), 3.19 (0.5H, dd, $J = 7.0$, 2.0 Hz), 3.35 (1.5H, s), 3.36 (1.5H, s), 3.46 (0.5H, t, $J = 7.5$ Hz), 3.47 (0.5H, t, $J = 7.5$ Hz), 3.60 (0.5H, dd, $J = 7.0$, 2.0 Hz), 3.79 (1.5H, s), 3.81 (1.5H, s), 3.89 (1H, dd, $J = 5.5$, 4.5 Hz), 4.76, 4.98 (0.5H each, ABq, $J = 6.5$ Hz), 4.79, 5.01 (0.5H each, ABq, $J = 6.5$ Hz), 6.86 (2H, d, $J = 9.0$ Hz), 7.25 (2H, d, $J = 9.0$ Hz), 7.47–7.70 (5H, m). MS m/z (relative intensity): 547 ($M^+ - 15$, 0.1), 545 (0.1), 487 (0.4), 477 (1.1), 383 (5.2), 325 (8.4), 121 (100). $[\alpha]_D^{25} = -12.2^\circ$ ($c = 3.53$, CHCl_3). Anal. Calcd for $\text{C}_{31}\text{H}_{46}\text{O}_7\text{S}$: C, 66.16; H, 8.23; S, 5.69. Found: C, 65.99; H, 8.18; S, 5.45. Exact MS m/z Calcd for $\text{C}_{30}\text{H}_{43}\text{O}_7\text{S}$ ($M^+ - 15$): 547.2729. Found: 547.2723.

(2R,3S,4S,5S)-3,5-Dimethyl-2-hydroxy-4,6-isopropylidenedioxyhexanal Trimethylene Dithioacetal (17) A 1.5 M solution of *n*-BuLi in hexane (2.21 ml, 3.32 mmol) was added dropwise to a stirred solution of 1,3-dithiane (399 mg, 3.32 mmol) in THF (5 ml) at -50°C under argon. The solution was stirred at -23 – -20°C for 2.5 h and then cooled to -80°C . A solution of the aldehyde (**16**) (308 mg, 1.66 mmol) in THF was added dropwise at -80 – -70°C . After 40 min, aqueous NH_4Cl was added, and the mixture was extracted with Et_2O . The extract was washed with brine, dried (Na_2SO_4), and evaporated. The residue was chromatographed on a silica gel column with CH_2Cl_2 as the eluent to give a 4.9:1 mixture of **17** and its C-2 isomer as a colorless oil (435 mg, 86%), which was solidified with MeOH. Recrystallization from MeOH gave pure **17** as colorless needles, mp 108–109°C. IR $\nu_{\text{max}}^{\text{KBr}}$ cm^{-1} : 3410. $^1\text{H-NMR}$ (CDCl_3) δ : 0.91 (3H, d, $J = 6.5$ Hz), 1.14 (3H, d, $J = 7.0$ Hz), 1.41 (3H, s), 1.44 (3H, s), 1.60–1.84 (1H, m), 1.90–2.20 (3H, m), 2.58 (1H, t, $J = 1.5$ Hz), 2.64–3.16 (4H, m), 3.61 (1H, dd, $J = 11.5$, 1.5 Hz), 3.84 (1H, d, $J = 11.0$ Hz), 3.91 (2H, dd, $J = 10.0$, 2.0 Hz), 4.14 (1H, dd, $J = 11.5$, 2.5 Hz). MS m/z (relative intensity): 307 ($M^+ + 1$, 0.2), 306 (1.1), 291 (9.5), 129 (94), 119 (100). $[\alpha]_D^{25} = +6.6^\circ$ ($c = 1.46$, CHCl_3). Exact MS m/z Calcd for $\text{C}_{14}\text{H}_{26}\text{O}_3\text{S}_2$ (M^+): 306.1323. Found: 306.1317.

(2R,3S,4R,5S)-3,5-Dimethyl-2,4,6-trihydroxyhexanal Trimethylene Dithioacetal (18) A 4N HCl solution (7.2 ml) was added to a stirred solution of the mixture of **17** and its C-2 isomer (435 mg, 1.43 mmol) in THF (15 ml) at room temperature. After 11 h, the solution was poured into aqueous NaHCO_3 , and extracted with CH_2Cl_2 . The extract was dried (Na_2SO_4) and evaporated *in vacuo*. The residue was chromatographed on a silica gel column with CH_2Cl_2 -MeOH (40:1) as the eluent to give a mixture of the triol (**18**) and its C-2 isomer as a colorless oil (350 mg, 92%), which was solidified with CH_2Cl_2 . Recrystallization from CH_2Cl_2 gave pure **18** as colorless needles, mp 140–141°C. $^1\text{H-NMR}$ (CDCl_3) δ : 0.98 (3H, d, $J = 7.0$ Hz), 1.05 (3H, d, $J = 7.0$ Hz), 1.90–2.20 (4H, m), 2.20–2.50 (1H, m), 2.50–3.20 (5H, m), 3.23 (1H, s), 3.67 (2H, d, $J = 5.0$ Hz), 3.80 (1H, d, $J = 10.0$ Hz), 3.89 (1H, dd, $J = 9.0$, 5.0 Hz), 4.02 (1H, dd, $J = 10.0$, 2.5 Hz). MS m/z (relative intensity): 266 ($M^+ + 0.2$), 250 (0.3), 248 (3.8), 207 (2.1), 160 (11), 129 (15), 119 (100). $[\alpha]_D^{25} = +2.2^\circ$ ($c = 1.06$, EtOH). Anal. Calcd for $\text{C}_{11}\text{H}_{22}\text{O}_3\text{S}_2$: C, 49.59; H, 8.32. Found: C, 49.05; H, 8.41. Exact MS m/z Calcd for $\text{C}_{11}\text{H}_{22}\text{O}_3\text{S}_2$ (M^+): 266.1010. Found: 266.1012.

(2R,3S,4S,5R)-6-tert-Butyldiphenylsilyloxy-2,4-dihydroxy-3,5-dimethylhexanal Trimethylene Dithioacetal (19) Imidazole (269 mg, 3.96 mmol) and *tert*-butyldiphenylsilyl chloride (688 μl , 2.64 mmol) were added to a stirred solution of the mixture of **18** and its C-2 isomer (350 mg, 1.32 mmol) in CH_2Cl_2 (30 ml) at room temperature. After 2 h, MeOH was added and the stirring was continued for 30 min. The reaction mixture was diluted with CH_2Cl_2 , washed with brine, dried (Na_2SO_4), and evaporated *in vacuo*. The residue was chromatographed on a silica gel column with CH_2Cl_2 as the eluent to give a mixture of the diol (**19**) and its C-2 isomer as a colorless oil (620 mg, 93%). IR $\nu_{\text{max}}^{\text{neat}}$ cm^{-1} : 3410. $^1\text{H-NMR}$ (CDCl_3) δ : 0.82 (3H, d, $J = 7.0$ Hz), 1.06 (9H, s), 1.09 (3H, d, $J = 7.0$ Hz), 1.64–2.20 (3H, m), 2.30–3.05 (5H, m), 3.14 (1H, s), 3.27 (1H, s), 3.64–4.00 (2H, m),

3.58 (1H, dd, $J = 10.5$, 4.0 Hz), 3.74 (1H, dd, $J = 10.5$, 4.0 Hz), 4.03 (1H, d, $J = 10.5$ Hz), 7.30–7.50 (6H, m), 7.54–7.76 (4H, m). $[\alpha]_D^{25} = +4.2^\circ$ ($c = 2.08$, CHCl_3).

(2R,3S,4S,5R)-6-tert-Butyldiphenylsilyloxy-3,5-dimethyl-2,4-isopropylidenedioxyhexanal Trimethylene Dithioacetal (20) 2-Methoxypropene (1.18 ml, 12.3 mmol) and pyridinium *p*-toluenesulfonate (PPTS) (31.2 mg, 0.123 mmol) were added to a stirred solution of the mixture of **19** and its C-2 isomer (620 mg, 1.23 mmol) in CH_2Cl_2 (60 ml). After 2.5 h, the solution was washed with aqueous NaHCO_3 , dried (Na_2SO_4), and evaporated *in vacuo*. The residue was chromatographed on a silica gel column with *n*-hexane-EtOAc (32:1) as the eluent to give **20** (528 mg, 79%) and its C-2 isomer (108 mg, 16%) as colorless oils.

Compound **20**: $^1\text{H-NMR}$ (CDCl_3) δ : 0.86 (3H, d, $J = 6.5$ Hz), 1.05 (3H, d, $J = 6.5$ Hz), 1.05 (9H, s), 1.44 (6H, s), 1.70–2.01 (3H, m), 2.05–2.20 (1H, m), 2.75–2.90 (4H, m), 3.51 (1H, dd, $J = 10.5$, 4.0 Hz), 3.57 (1H, dd, $J = 10.5$, 5.0 Hz), 3.76 (1H, dd, $J = 9.5$, 2.0 Hz), 3.96 (1H, dd, $J = 10.5$, 2.0 Hz), 4.15 (1H, d, $J = 10.5$ Hz), 7.34–7.47 (6H, m), 7.62–7.70 (4H, m). MS m/z (relative intensity): 545 ($M^+ + 1$, 1.5), 544 (3.8), 487 (0.8), 469 (3.1), 425 (26), 367 (69), 269 (100). $[\alpha]_D^{25} = +4.5^\circ$ ($c = 1.12$, CHCl_3). Exact MS m/z Calcd for $\text{C}_{30}\text{H}_{44}\text{O}_3\text{S}_2\text{Si}$ (M^+): 544.2501. Found: 544.2526.

(2R,3S,4S,5R)-6-tert-Butyldiphenylsilyloxy-3,5-dimethyl-2,4-isopropylidenedioxyhexanal (10) NaHCO_3 (1.29 g, 15.3 mmol) and MeI (477 μl , 7.62 mmol) were added to a solution of **20** (207 mg, 0.381 mmol) in 90% MeCN (10 ml), and the mixture was stirred at 70°C for 1.5 h, then cooled, CH_2Cl_2 and H_2O were added. The organic layer was dried (Na_2SO_4) and evaporated *in vacuo*. The residue was chromatographed on a silica gel column with *n*-hexane-EtOAc (8:1) as the eluent to give the aldehyde (**10**) as a colorless oil (153 mg, 88%). IR $\nu_{\text{max}}^{\text{neat}}$ cm^{-1} : 1735. $^1\text{H-NMR}$ (CDCl_3) δ : 0.76 (3H, d, $J = 6.5$ Hz), 1.04 (3H, d, $J = 5.0$ Hz), 1.06 (9H, s), 1.43 (3H, s), 1.50 (3H, s), 1.60–2.10 (2H, m), 3.55 (1H, dd, $J = 4.5$, 2.0 Hz), 3.74 (1H, dd, $J = 9.5$, 2.0 Hz), 4.22 (1H, d, $J = 2.5$ Hz), 7.30–7.50 (6H, m), 7.54–7.76 (4H, m), 9.48 (1H, s). MS m/z (relative intensity): 441 ($M^+ - 13$, 0.3), 440 (0.9), 439 (2.7), 397 (4.7), 367 (3.6), 339 (42), 309 (30), 269 (100). Exact MS m/z Calcd for $\text{C}_{26}\text{H}_{35}\text{O}_3\text{Si}$ ($M^+ - 15$): 439.2304. Found: 439.2295.

(2R,3S,4S,5R)-1-tert-Butyldiphenylsilyloxy-2,4-dimethyl-3,5-isopropylidenedioxyheptan-6-one (11) A 1.0 M solution of MeMgI in Et_2O (0.65 ml) was added dropwise to a stirred solution of **10** (58.5 mg, 0.129 mmol) in Et_2O (5 ml) at -15°C . After 1 h, the reaction mixture was poured into ice-cooled aqueous NH_4Cl and extracted with Et_2O . The extract was washed with brine, dried (Na_2SO_4), and evaporated *in vacuo*. The residue was chromatographed on a silica gel column with *n*-hexane-EtOAc (16:1) as the eluent to give **(2R,3S,4S,5R)-1-tert-butylidiphenylsilyloxy-2,4-dimethyl-6-hydroxy-3,5-isopropylidenedioxyheptane** as a colorless oil (54.1 mg, 89%). MS m/z (relative intensity): 456 ($M^+ - 14$, 0.1), 455 (0.4), 425 (0.2), 355 (19), 269 (16), 199 (58), 43 (100). Exact MS m/z Calcd for $\text{C}_{27}\text{H}_{39}\text{O}_4\text{Si}$ ($M^+ - 15$): 455.2618. Found: 455.2619.

Dimethyl sulfoxide (DMSO) (48 μl , 0.68 mmol) was added dropwise to a stirred solution of oxalyl chloride (30 μl , 0.34 mmol) in CH_2Cl_2 (3 ml) at -60°C , and the solution was stirred at -60 – -55°C for 30 min. A solution of the above alcohol (54.1 mg, 0.113 mmol) in CH_2Cl_2 (3 ml) was added dropwise at -70°C , and the mixture was stirred at -65 – -55°C for 1 h. Et_3N (141 μl , 1.02 mmol) was added dropwise at -70°C . The reaction mixture was allowed to warm to room temperature, washed with aqueous KHSO_4 and brine, dried (Na_2SO_4), and evaporated *in vacuo*. The residue was chromatographed on a silica gel column with *n*-hexane-EtOAc (16:1) as the eluent to give the recovered alcohol (4.7 mg, 8.7%) and the ketone (**11**) as a colorless oil (45.4 mg, 86%). IR $\nu_{\text{max}}^{\text{neat}}$ cm^{-1} : 1715. $^1\text{H-NMR}$ (CDCl_3) δ : 0.70 (3H, d, $J = 7.0$ Hz), 1.04 (3H, d, $J = 6.5$ Hz), 1.06 (9H, s), 1.41 (3H, s), 1.48 (3H, s), 1.60–2.08 (2H, m), 2.12 (3H, s), 3.53 (2H, dd, $J = 5.0$, 2.0 Hz), 3.74 (1H, dd, $J = 10.0$, 2.0 Hz), 4.23 (1H, d, $J = 2.5$ Hz), 7.30–7.50 (6H, m), 7.54–7.76 (4H, m). MS m/z (relative intensity): 455 ($M^+ - 13$, 0.3), 454 (0.7), 453 (2.0), 425 (1.4), 412 (2.0), 411 (6.0), 353 (34), 281 (34), 269 (100). $[\alpha]_D^{25} = +19.3^\circ$ ($c = 1.45$, CHCl_3). Exact MS m/z Calcd for $\text{C}_{27}\text{H}_{37}\text{O}_3\text{Si}$ ($M^+ - 15$): 453.2461. Found: 453.2474.

(2R,3S,4S,5R,6R,7R,9S,10S,11R,12R,13R)-3,5,9,11-Bis(isopropylidenedioxy)-1-tert-butylidiphenylsilyloxy-13-(4-methoxybenzyloxy)-12-methoxymethoxy-2,4,8,10,12-pentamethyl-7-phenylsulfinylpentadecan-6-ol (21) A 1.5 M solution of *n*-BuLi in hexane (0.34 ml, 0.52 mmol) was added dropwise to a stirred solution of diisopropylamine (80 μl , 0.57 mmol) in THF (2 ml) at 0°C under argon. After 20 min, a solution of **9** (282 mg, 0.502 mmol) in THF (2 ml) was added dropwise at -15°C . After 30 min, the solution was cooled at -80°C , and a solution of **10** (117 mg, 0.258 mmol) in THF (1.2 ml) was added dropwise at -81 – -77°C . The reaction mixture was stirred at -80°C for 1 h, then treated with aqueous NH_4Cl , and extracted with CH_2Cl_2 . The extract was washed with brine,

dried (Na_2SO_4), and evaporated *in vacuo*. The residue was chromatographed on a silica gel column with *n*-hexane-EtOAc (8:1–4:1) to give recovered **10** (11 mg, 10%), recovered **9** (138 mg, 49%), and a mixture of four diastereoisomers of **21** as a colorless oil (225 mg, 86%).

(2R,3S,4S,5R,6R,8R,9S,10S,11R,12R,13R)-3,5:9,11-Bis(isopropylidenedioxy)-1-tert-butylidiphenylsilyloxy-13-(4-methoxybenzyloxy)-12-methoxymethoxy-2,4,8,10,12-pentamethylpentadecan-6-ol (22) A solution of **21** (72.3 mg, 0.072 mmol) in EtOH (20 ml) was stirred in the presence of Raney Ni (W-2) (*ca.* 20 ml) at room temperature for 30 min. The catalyst was removed by filtration and washed with EtOH and CH_2Cl_2 . The filtrates were evaporated *in vacuo* to leave an oil, which was subjected to preparative TLC on silica gel. Development with *n*-hexane-EtOAc (4:1) gave the alcohol (**22**) as a colorless oil (45.6 mg, 71%). IR $\nu_{\text{max}}^{\text{neat}} \text{cm}^{-1}$: 3420. $^1\text{H-NMR}$ (CDCl_3) δ : 0.80 (3H, d, $J=6.5$ Hz), 0.96 (3H, d, $J=6.0$ Hz), 1.03 (3H, d, $J=7.0$ Hz), 1.04 (9H, s), 1.06 (3H, d, $J=8.0$ Hz), 1.07 (3H, t, $J=7.5$ Hz), 1.30 (3H, s), 1.34 (3H, s), 1.35 (3H, s), 1.37 (3H, s), 1.38 (3H, s), 1.63 (1H, quint, $J=7.5$ Hz), 1.72–2.00 (7H, m), 2.92 (1H, d, $J=3.0$ Hz), 3.35 (3H, s), 3.37–3.51 (3H, m), 3.54–3.68 (3H, m), 3.80 (3H, s), 3.88 (1H, d, $J=4.0$ Hz), 4.51, 4.60 (1H each, ABq, $J=10.5$ Hz), 4.79, 4.98 (1H each, ABq, $J=7.0$ Hz, 6.86 (2H, d, $J=9.0$ Hz), 7.26 (2H, d, $J=9.0$ Hz), 7.32–7.45 (6H, m), 7.60–7.70 (4H, m). MS m/z (relative intensity): 877 ($M^+ - 15$, 0.1), 835 (0.3), 623 (0.2), 597 (0.3), 579 (0.6), 565 (0.6), 535 (0.8), 367 (1.6), 121 (100). $[\alpha]_D^{25} - 3.4^\circ$ ($c=1.66$, CHCl_3). Exact MS m/z Calcd for $\text{C}_{31}\text{H}_{49}\text{O}_9$ ($M^+ - 327$): 565.3376. Found: 565.3365.

(2R,3S,4S,5R,6R,8R,9S,10S,11R,12R,13R)-3,5:9,11-Bis(isopropylidenedioxy)-1-tert-butylidiphenylsilyloxy-13-(4-methoxybenzyloxy)-12-methoxymethoxy-2,4,8,10,12-pentamethylpentadecan-6-one (23) The alcohol (**22**) (98.5 mg, 0.111 mmol) was oxidized with oxalyl chloride (40 μl , 0.45 mmol), DMSO (64 μl , 0.90 mmol), and Et_3N (188 μl , 0.135 mmol) as indicated for the preparation of **11**. The crude product was chromatographed on a silica gel column with *n*-hexane-EtOAc (16:1) as the eluent to give the ketone (**23**) as a colorless oil (90.2 mg, 92%). IR $\nu_{\text{max}}^{\text{neat}} \text{cm}^{-1}$: 1710. $^1\text{H-NMR}$ (CDCl_3) δ : 0.67 (3H, d, $J=7.0$ Hz), 0.91 (3H, d, $J=7.0$ Hz), 1.04 (3H, d, $J=4.5$ Hz), 1.05 (3H, d, $J=4.5$ Hz), 1.06 (9H, s), 1.07 (3H, t, $J=7.5$ Hz), 1.26 (3H, s), 1.27 (3H, s), 1.33 (3H, s), 1.40 (3H, s), 1.47 (3H, s), 1.55–1.90 (6H, m), 2.42 (1H, dd, $J=18.0$, 6.0 Hz), 2.56 (1H, dd, $J=18.0$, 7.5 Hz), 3.15 (1H, dd, $J=7.0$, 2.0 Hz), 3.36 (3H, s), 3.44 (1H, dd, $J=8.5$, 2.5 Hz), 3.48 (1H, dd, $J=10.5$, 5.5 Hz), 3.56 (1H, dd, $J=10.5$, 4.0 Hz), 3.73 (1H, dd, $J=9.5$, 2.0 Hz), 3.80 (3H, s), 3.87 (1H, d, $J=4.0$ Hz), 4.21 (1H, d, $J=2.0$ Hz), 4.52, 4.61 (1H each, ABq, $J=10.5$ Hz), 4.78, 4.99 (1H each, ABq, $J=7.0$ Hz), 6.86 (2H, d, $J=9.0$ Hz), 7.26 (2H, d, $J=9.0$ Hz), 7.34–7.48 (6H, m), 7.60–7.80 (4H, m). MS m/z (relative intensity): 833 ($M^+ - 57$, 0.3), 800 (0.2), 782 (0.3), 742 (0.4), 653 (0.5), 621 (0.5), 595 (0.7), 563 (0.9), 269 (10), 121 (100). $[\alpha]_D^{25} + 3.4^\circ$ ($c=3.37$, CHCl_3). Exact MS m/z Calcd for $\text{C}_{35}\text{H}_{57}\text{O}_9$ ($M^+ - 269$): 621.4002. Found: 621.3954.

(2R,3S,4S,5R,6R,8R,9S,10S,11R,12R,13R)-3,5:9,11-Bis(isopropylidenedioxy)-1-tert-butylidiphenylsilyloxy-2,4,6,8,10,12-hexamethyl-13-(4-methoxybenzyloxy)-12-methoxymethoxypentadecan-6-ol (24) A 1.40 M solution of MeLi in Et_2O (0.36 ml, 0.51 mmol) was added dropwise to a stirred solution of **23** (45.0 mg, 0.056 mmol) in THF (8 ml) at -85°C under argon. After 1 h, aqueous NH_4Cl was added, and the mixture was extracted with Et_2O . The extract was washed with brine, dried (Na_2SO_4), and evaporated *in vacuo* to leave an oil, which was subjected to preparative TLC on silica gel. Development with CH_2Cl_2 -MeOH (80:1) gave **24** (38.0 mg, 82.9%) and its C-6 isomer (**25**) (5.3 mg, 11.6%) as colorless oils.

Compound 24: IR $\nu_{\text{max}}^{\text{neat}} \text{cm}^{-1}$: 3550. $^1\text{H-NMR}$ (CDCl_3) δ : 0.91 (3H, d, $J=6.5$ Hz), 0.95 (3H, d, $J=7.0$ Hz), 1.03 (3H, d, $J=7.5$ Hz), 1.05 (9H, s), 1.06 (3H, d, $J=6.5$ Hz), 1.07 (3H, t, $J=6.0$ Hz), 1.25 (3H, s), 1.28 (3H, s), 1.33 (3H, s), 1.34 (3H, s), 1.40 (3H, s), 1.41 (3H, s), 1.50–2.00 (8H, m), 2.40 (1H, s), 3.28 (1H, dd, $J=6.5$, 2.5 Hz), 3.35 (3H, s), 3.41 (1H, dd, $J=6.0$, 2.5 Hz), 3.43 (1H, s), 3.54 (2H, d, $J=5.0$ Hz), 3.67 (1H, d, $J=9.0$ Hz), 3.80 (3H, s), 3.87 (1H, d, $J=4.0$ Hz), 4.52, 4.60 (1H each, ABq, $J=11.0$ Hz), 4.80, 4.99 (1H each, ABq, $J=7.0$ Hz), 6.86 (2H, d, $J=8.5$ Hz), 7.27 (2H, d, $J=8.5$ Hz), 7.32–7.50 (6H, m), 7.58–7.70 (4H, m). MS m/z (relative intensity): 849 ($M^+ - 57$, 0.1), 621 (0.2), 549 (1.2), 425 (0.5), 411 (1.3), 367 (2.5), 269 (5.8), 121 (100). $[\alpha]_D^{25} + 14.1^\circ$ ($c=1.24$, CHCl_3). Exact MS m/z Calcd for $\text{C}_{31}\text{H}_{49}\text{O}_8$ ($M^+ - 357$): 549.3426. Found: 549.3414.

(2R,3S,4S,5R,6R,8R,9S,10S,11R,12R,13R)-3,5:9,11-Bis(isopropylidenedioxy)-6,12-bis(methoxymethoxy)-1-tert-butylidiphenylsilyloxy-2,4,6,8,10,12-hexamethyl-13-(4-methoxybenzyloxy)pentadecane (26) Diisopropylethylamine (0.60 ml, 3.4 mmol) and then chloromethyl methyl ether (0.13 ml, 1.7 mmol) were added dropwise to a stirred solution of **24** (61.8 mg, 0.0682 mmol) in CH_2Cl_2 (2 ml) at room temperature. The solution was stirred at 50 – 55°C for 11 h. MeOH was added, and the

stirring was continued at room temperature for 30 min. The reaction mixture was diluted with Et_2O , washed with aqueous KHSO_4 and brine, dried (Na_2SO_4), and evaporated *in vacuo* to leave **26** as a pale yellow oil (66.0 mg, 100%). $^1\text{H-NMR}$ (CDCl_3) δ : 0.83 (3H, d, $J=6.5$ Hz), 1.01 (3H, d, $J=7.0$ Hz), 1.03 (3H, d, $J=6.0$ Hz), 1.04 (9H, s), 1.05 (3H, d, $J=7.0$ Hz), 1.08 (3H, d, $J=7.5$ Hz), 1.25 (6H, s), 1.28 (3H, s), 1.32 (3H, s), 1.40 (6H, s), 1.54 (1H, s), 1.60–1.92 (7H, m), 3.27 (1H, dd, $J=6.5$, 2.0 Hz), 3.34 (3H, s), 3.36 (3H, s), 3.45 (1H, dd, $J=8.5$, 2.0 Hz), 3.50 (2H, d, $J=3.5$ Hz), 3.68 (1H, dd, $J=9.5$, 1.0 Hz), 3.74 (1H, d, $J=1.0$ Hz), 3.80 (3H, s), 3.89 (1H, d, $J=4.0$ Hz), 4.52, 4.61 (1H each, ABq, $J=10.5$ Hz), 4.75, 4.84 (1H each, ABq, $J=6.5$ Hz), 4.78, 5.00 (1H each, ABq, $J=7.0$ Hz), 6.87 (2H, d, $J=8.5$ Hz), 7.27 (2H, d, $J=8.5$ Hz), 7.30–7.50 (6H, m), 7.58–7.70 (4H, m). MS m/z (relative intensity): 637 ($M^+ - 313$, 0.2), 607 (0.5), 593 (0.6), 579 (0.7), 549 (1.5), 425 (1.1), 367 (5.1), 269 (7.5), 121 (100). FD MS m/z : 951 ($M^+ + 1$). $[\alpha]_D^{25} + 11.4^\circ$ ($c=1.18$, CHCl_3).

(2R,3S,4S,5R,6R,8R,9S,10S,11R,12R,13R)-3,5:9,11-Bis(isopropylidenedioxy)-6,12-bis(methoxymethoxy)-2,4,6,8,10,12-hexamethyl-13-(4-methoxybenzyloxy)pentadecanoic Acid (27) A 1.0 M solution of *n*- Bu_4NF in THF (0.40 ml, 0.40 mmol) was added to a stirred solution of **26** (63.1 mg, 0.0664 mmol) in THF (4 ml) at room temperature. The solution was stirred at 55 – 60°C for 2 h, then diluted with Et_2O , washed with brine, dried (Na_2SO_4), and evaporated *in vacuo*. The residue was chromatographed on a silica gel column with CH_2Cl_2 -MeOH (80:1) as the eluent to give **(2R,3S,4S,5R,6R,8R,9S,10S,11R,12R,13R)-3,5:9,11-bis(isopropylidenedioxy)-6,12-bis(methoxymethoxy)-2,4,6,8,10,12-hexamethyl-13-(4-methoxybenzyloxy)pentadecan-1-ol** as a colorless oil (44.4 mg, 94%). IR $\nu_{\text{max}}^{\text{neat}} \text{cm}^{-1}$: 3480. $^1\text{H-NMR}$ (CDCl_3) δ : 0.94 (3H, d, $J=6.5$ Hz), 1.00 (3H, d, $J=6.5$ Hz), 1.03 (3H, d, $J=6.5$ Hz), 1.05 (3H, d, $J=5.5$ Hz), 1.07 (3H, t, $J=7.5$ Hz), 1.27 (3H, s), 1.30 (3H, s), 1.32 (3H, s), 1.35 (3H, s), 1.39 (3H, s), 1.40 (3H, s), 1.52 (1H, s), 1.60–2.00 (7H, m), 3.27 (1H, dd, $J=6.5$, 2.0 Hz), 3.33 (3H, s), 3.36 (3H, s), 3.47 (1H, dd, $J=8.0$, 2.0 Hz), 3.52 (1H, d, $J=5.0$ Hz), 3.58 (2H, s), 3.62 (1H, d, $J=1.5$ Hz), 3.80 (3H, s), 3.89 (1H, d, $J=4.0$ Hz), 4.52, 4.62 (1H each, ABq, $J=10.5$ Hz), 4.76, 4.82 (1H each, ABq, $J=6.5$ Hz), 4.80, 5.00 (1H each, ABq, $J=6.5$ Hz), 6.86 (2H, d, $J=8.5$ Hz), 7.26 (2H, d, $J=8.5$ Hz). MS m/z (relative intensity): 697 ($M^+ - 15$, 0.1), 457 (0.2), 311 (3.1), 269 (1.5), 121 (100). $[\alpha]_D^{25} + 0.6^\circ$ ($c=0.80$, CHCl_3). Exact MS m/z Calcd for $\text{C}_{25}\text{H}_{45}\text{O}_7$ ($M^+ - 255$): 457.3165. Found: 457.3168.

Jones reagent (2.67 M, 37 μl , 0.099 mmol) was added to a stirred solution of the above alcohol (17.6 mg, 0.0247 mmol) in acetone (2 ml) at -17°C . After 2 h, isopropanol was added. The reaction mixture was diluted with CH_2Cl_2 , washed with aqueous KHSO_4 and brine, dried (Na_2SO_4), and evaporated *in vacuo*. The residue was chromatographed on a silica gel column with CH_2Cl_2 -MeOH (40:1) as the eluent to give the carboxylic acid (**27**) as a colorless oil (14.0 mg, 78%). IR $\nu_{\text{max}}^{\text{neat}} \text{cm}^{-1}$: 1735, 1710. $^1\text{H-NMR}$ (CDCl_3) δ : 0.97 (3H, d, $J=7.0$ Hz), 0.99 (3H, d, $J=7.5$ Hz), 1.04 (3H, d, $J=6.5$ Hz), 1.08 (3H, t, $J=7.5$ Hz), 1.24 (3H, d, $J=7.0$ Hz), 1.27 (3H, s), 1.29 (3H, s), 1.31 (3H, s), 1.34 (3H, s), 1.43 (6H, s), 1.55–1.95 (7H, m), 2.67 (1H, dt, $J=16.5$, 7.0 Hz), 3.24 (1H, dd, $J=7.0$, 2.0 Hz), 3.34 (3H, s), 3.37 (3H, s), 3.43 (1H, dd, $J=8.0$, 2.5 Hz), 3.80 (3H, s), 3.84 (1H, d, $J=2.0$ Hz), 3.86 (1H, d, $J=3.0$ Hz), 3.89 (1H, d, $J=3.5$ Hz), 4.75, 4.80 (1H each, ABq, $J=10.5$ Hz), 4.85, 4.96 (1H each, ABq, $J=6.5$ Hz), 6.86 (2H, d, $J=8.5$ Hz), 7.26 (2H, d, $J=8.5$ Hz). MS m/z (relative intensity): 471 ($M^+ - 255$, 0.2), 457 (0.2), 413 (7.9), 121 (100). FD MS m/z : 727 ($M^+ + 1$). $[\alpha]_D^{25} - 1.7^\circ$ ($c=0.58$, CHCl_3). Exact MS m/z Calcd for $\text{C}_{24}\text{H}_{39}\text{O}_9$ ($M^+ - 255$): 471.2594. Found: 471.2577.

3,5:9,11-Di-O-isopropylidene-6,12-di-O-methoxymethyl-(9S)-9-dihydroerythronolide A Seco-acid (28) A solution of **27** (5.8 mg, 0.0080 mmol) in EtOH (2 ml) was hydrogenated in the presence of 10% Pd-C (10 mg) at 55°C for 13 h. After removal of the catalyst by filtration, the filtrate was evaporated *in vacuo* to leave the seco-acid (**28**) as a colorless oil (4.8 mg, 99%). IR $\nu_{\text{max}}^{\text{neat}} \text{cm}^{-1}$: 1730, 1715. $^1\text{H-NMR}$ (CDCl_3) δ : 0.97 (3H, d, $J=7.0$ Hz), 1.00 (3H, d, $J=6.5$ Hz), 1.04 (3H, t, $J=7.0$ Hz), 1.09 (3H, d, $J=6.5$ Hz), 1.24 (3H, d, $J=7.0$ Hz), 1.27 (3H, s), 1.28 (3H, s), 1.34 (3H, s), 1.36 (3H, s), 1.43 (6H, s), 1.50–1.80 (4H, m), 1.80–2.10 (3H, m), 2.66 (1H, dt, $J=17.0$, 7.0 Hz), 3.29 (1H, dd, $J=7.5$, 2.0 Hz), 3.34 (3H, s), 3.41 (3H, s), 3.69 (1H, dd, $J=10.5$, 1.5 Hz), 3.86 (1H, d, $J=2.0$ Hz), 3.87 (1H, dd, $J=9.0$, 2.0 Hz), 3.93 (1H, d, $J=4.5$ Hz), 4.74, 4.82 (1H each, ABq, $J=7.0$ Hz), 4.78, 4.85 (1H each, ABq, $J=7.5$ Hz). MS m/z (relative intensity): 591 ($M^+ - 15$, 0.1), 485 (0.1), 471 (0.3), 413 (1.2), 343 (2.8), 325 (2.7), 285 (7.1), 223 (14), 45 (100). $[\alpha]_D^{25} + 18.2^\circ$ ($c=0.44$, CHCl_3). Exact MS m/z Calcd for $\text{C}_{30}\text{H}_{55}\text{O}_{11}$ ($M^+ - 15$): 591.3744. Found: 591.3770.

3,5:9,11-Di-O-isopropylidene-6,12-di-O-methoxymethyl-(9S)-9-dihydroerythronolide A (29) Et_3N (7.7 μl , 0.055 mmol) was added to a stirred solution of **28** (30.3 mg, 0.050 mmol) in THF (1 ml) at room temperature

under argon. After 10 min, 2,4,6-trichlorobenzoyl chloride (8.0 μ l, 0.050 mmol) was added, the stirring was continued for 2 h at room temperature, and then the solution was made up to 25 ml with toluene. The solution was added very slowly to a stirred solution of DMAP (151 mg, 1.25 mmol) in refluxing toluene (25 ml) over a period of 39 h under argon. The stirring was continued for 11 h. After being cooled, the reaction mixture was washed with aqueous KHSO_4 and brine, dried (Na_2SO_4), and evaporated *in vacuo*. The residue was chromatographed on a silica gel column with *n*-hexane–EtOAc (4:1) as the eluent to give crude **29**, which was subjected to preparative TLC on silica gel. Development with *n*-hexane–EtOAc (4:1) gave pure **29** as a colorless solid (7.9 mg, 27%), mp 128–129 °C (from *n*-hexane). IR $\nu_{\text{max}}^{\text{KBr}}$ cm^{-1} : 1725. $^1\text{H-NMR}$ (CDCl_3) δ : 0.85 (3H, t, $J=7.5$ Hz), 1.02 (3H, d, $J=6.5$ Hz), 1.16 (3H, s), 1.18 (3H, d, $J=6.0$ Hz), 1.26 (3H, d, $J=6.0$ Hz), 1.29 (3H, s), 1.30 (3H, d, $J=7.0$ Hz), 1.05–1.40 (2H, m), 1.42 (3H, s), 1.43 (3H, s), 1.46 (6H, s), 1.45–1.68 (2H, m), 1.74–2.02 (2H, m), 2.03–2.29 (1H, m), 2.74 (1H, dq, $J=10.5, 6.5$ Hz), 3.10 (1H, d, $J=11.0$ Hz), 3.33 (3H, s), 3.39 (3H, s), 3.62 (1H, s), 3.78 (1H, d, $J=10.5$ Hz), 4.02 (1H, s), 4.69, 4.80 (1H each, ABq, $J=7.0$ Hz), 4.79, 4.87 (1H each, ABq, $J=6.0$ Hz), 5.47 (1H, dd, $J=11.5, 2.0$). MS m/z (relative intensity): 573 ($M^+ - 15$, 0.7), 481 (0.3), 413 (0.6), 253 (1.1), 45 (100). $[\alpha]_D^{25} - 15.0^\circ$ ($c=0.18$, CHCl_3). Anal. Calcd for $\text{C}_{31}\text{H}_{56}\text{O}_{10}$: C, 63.23; H, 9.58. Found: C, 63.45; H, 9.56. Exact MS m/z Calcd for $\text{C}_{30}\text{H}_{53}\text{F}_{10}\text{O}$ ($M^+ - 15$): 573.3638. Found: 573.3633.

(9S)-9-Dihydroerythronolide A (3) A solution of **29** (7.9 mg, 0.013 mmol) in AcOH (1 ml) and H_2O (0.5 ml) was stirred at 50–55 °C for 2 h. The solvent was evaporated off *in vacuo*. Benzene was added to the residue, and the solution was evaporated again *in vacuo* to leave **3** as a colorless solid (6.1 mg, 100%), which was recrystallized from CHCl_3 –petroleum ether to give colorless needles, mp 201–203 °C. IR $\nu_{\text{max}}^{\text{KBr}}$ cm^{-1} : 3400, 2975, 2940, 1715, 1185, 1080, 975. $^1\text{H-NMR}$ (CDCl_3) δ : 0.91 (3H, t, $J=7.5$ Hz), 1.04 (3H, d, $J=7.0$ Hz), 1.05 (3H, s), 1.24 (3H, d, $J=7.0$ Hz), 1.24 (3H, s), 1.30 (6H, d, $J=6.5$ Hz), 1.12–1.35 (1H, m), 1.38–1.65 (4H, m), 1.76 (1H, s, OH), 1.94 (1H, dq, $J=7.5, 1.5$ Hz), 1.98–2.03 (1H, m), 2.52 (1H, s, OH), 2.79 (1H, dq, $J=10.5, 6.5$ Hz), 2.90 (1H, s, OH), 2.96 (1H, dt, $J=9.5, 2.5$ Hz), 3.39 (1H, s, OH), 3.41 (1H, d, $J=7.5$ Hz, OH), 3.49 (1H, dd, $J=4.5, 1.5$ Hz), 3.86 (1H, d, $J=10.5$ Hz), 3.96 (1H, s), 4.25 (1H, d, $J=4.5$ Hz, OH), 4.61 (1H, dd, $J=11.0, 1.5$ Hz). MS m/z (relative intensity): 402 ($M^+ - 18$, 0.3), 384 (1.6), 366 (1.9), 327 (5.2), 43 (100). $[\alpha]_D^{25} + 10.8^\circ$ ($c=0.11$, CHCl_3) (lit.¹⁷) mp 199–200 °C; $[\alpha]_D^{25} + 9.5^\circ$ (MeOH).

References and Notes

- 1) Chiral synthesis of polyketide-derived natural products. XXII. For part XXI, see: H. Tone, T. Nishi, Y. Oikawa, M. Hikota and O. Yonemitsu, *Chem. Pharm. Bull.*, **37**, 1160 (1989).
- 2) A preliminary report of this work: H. Tone, T. Nishi, Y. Oikawa, M. Hikota and O. Yonemitsu, *Tetrahedron Lett.*, **28**, 4569 (1987).
- 3) Successful total synthetic studies directed toward **1**, **2**, and **3**: a) E. J. Corey, P. B. Hopkins, S. Kim, S. Yoo, K. P. Nambiar and J. R. Falk, *J. Am. Chem. Soc.*, **101**, 7131 (1979); b) R. B. Woodward, *et al.*, *ibid.*, **103**, 3210, 3213, 3215 (1981); c) B. Bernet, P. M. Bishop, M. Caron, T. Kawamata, B. L. Roy, L. Ruest, G. Sauve, P. Soucy and P. Deslongchamps, *Can. J. Chem.*, **63**, 2810, 2814, 2818 (1985); d) M. Kinoshita, M. Arai, K. Tomioka and M. Nakata, *Tetrahedron Lett.*, **27**, 1811, 1815 (1986); e) G. Stork and S. D. Rychnovsky, *J. Am. Chem. Soc.*, **109**, 1565 (1987); f) T. Nakata, M. Fukui and T. Oishi, *Tetrahedron Lett.*, **29**, 2219, 2223 (1988).
- 4) J. Boutagy and R. Thomas, *Chem. Rev.*, **74**, 87 (1974); W. S. Wadsworth, Jr., *Org. React.*, **25**, 73 (1977).
- 5) G. Stork, I. Paterson and F. K. C. Lee, *J. Am. Chem. Soc.*, **104**, 4686 (1982).
- 6) J. Inanaga, K. Hirata, H. Saeki, T. Katsuki and M. Yamaguchi, *Bull. Chem. Soc. Jpn.*, **52**, 1989 (1979).
- 7) Unless otherwise noted, the numberings are based on that of **3**.
- 8) Cf. I. Paterson, *Tetrahedron Lett.*, **24**, 1311 (1983).
- 9) I. Nakagawa, K. Aki and T. Hata, *J. Chem. Soc., Perkin Trans. 1*, **1983**, 1315.
- 10) Y. Oikawa, T. Nishi and O. Yonemitsu, *J. Chem. Soc., Perkin Trans. 1*, **1985**, 19.
- 11) H. Tone, M. Hikota, T. Hamada and O. Yonemitsu, *Chem. Pharm. Bull.*, **37**, 1155 (1989).
- 12) E. J. Corey and B. W. Erickson, *J. Org. Chem.*, **36**, 3553 (1971).
- 13) M. Fetizon and M. Jurion, *J. Chem. Soc., Chem. Commun.*, **1972**, 382.
- 14) Y. Oikawa, T. Yoshioka and O. Yonemitsu, *Tetrahedron Lett.*, **23**, 885 (1982); K. Horita, T. Yoshioka, T. Tanaka, Y. Oikawa and O. Yonemitsu, *Tetrahedron*, **42**, 3021 (1986).
- 15) E. J. Corey and D. J. Brunelle, *Tetrahedron Lett.*, **1976**, 3409.
- 16) E. J. Corey and K. C. Nicolaou, *J. Am. Chem. Soc.*, **96**, 5614 (1974).
- 17) P. H. Jones and E. K. Rowley, *J. Org. Chem.*, **33**, 665 (1968).