


A Convenient Method for Synthesis of Optically Active β -Hydroxyamines from Primary Amines through Enecarbamates as Key Intermediates[†]

Yoshihiro Matsumura,^{1*} Takahiro Ohishi,¹ Chihiro Sonoda,¹ Toshihide Maki,¹ and Mitsuaki Watanabe²

*Faculty of Pharmaceutical Sciences¹ and Center for Instrumental Analysis,²
Nagasaki University, 1-14 Bunkyo-machi, Nagasaki 852, Japan*

Abstract: This report describes a new method to prepare optically active β -hydroxyamines starting from primary amines. The method consists of a transformation of *N*-methoxycarbonylated primary amines to the corresponding enecarbamates utilizing electrochemical oxidation and an asymmetric hydroboration of the enecarbamates to produce optically active β -hydroxyamines. © 1997 Elsevier Science Ltd.


Optically active β -hydroxyamines **2** occur widely in nature¹ and are often used in organic synthesis as exemplified by the use as chiral auxiliaries.² Thus, exploiting convenient methods for the synthesis of optically active **2** is worthwhile. Although a variety of synthetic methods for racemic **2** have been reported so far,³ there have been only a few methods for the synthesis of optically active ones.^{4,5} We report a new route to prepare optically active **2** from easily available achiral primary amines **1** (Scheme 1). There have not been so far any methods for the conversion of **1** to **2**.



Scheme 1

Our strategy is the transformation of *N*-protected primary amines **3** to the corresponding enecarbamates **5**⁶ followed by the asymmetric introduction of a hydroxy group to the β position to produce *N*-protected optically active β -hydroxyamines **6** (eq 1). This strategy is based on our previous finding that electrochemical α -methoxylation of carbamates **3** followed by acid-catalyzed elimination of methanol from the α -methoxylated carbamates **4** afford **5**⁷ as well as on the well-known asymmetric hydroboration of alkenes.⁸ However, the yields of the acid-catalyzed conversion of **4** to **5** were not always satisfactory and there have not been any

precedents on hydroboration of the encarbamates **5**.⁵ This report describes both an improved procedure for the methanol elimination (**4** to **5**) and an asymmetric hydroboration of **5** to give **6**.



Results and Discussion

The first step in our method is electrochemical oxidation of **3a-e** in methanol, which was achieved by a reported procedure⁷ to give α -methoxycarbamates **4a-e** in the yields shown in Table 1. Although **4f** was hardly obtainable by the electrochemical α -methoxylation method,⁹ it could be prepared by electrochemical decarboxylation of *N*-methoxycarbonylphenylalanine.¹⁰

Table 1. Electrochemical Oxidation of Carbamates **3a-e** in MeOH

Entry	Carbamate 3a-e	R	R'	Supporting Electrolyte	F/mol	Product 4a-e	Yield (%)
1	3a	Me	H	Et ₄ NOTs	20	4a	76
2	3b	Me	Me	Et ₄ NOTs	15	4b	63
3	3c	<i>i</i> -Pr	H	Et ₄ NOTs	23	4c	58
4	3d	Me(CH ₂) ₅	H	Et ₄ NBF ₄	8	4d	86
5	3e	CO ₂ Me	H	Et ₄ NBF ₄	6	4e	81
6	- ^a	Ph	H	AcONa	6	4f	81

^a The starting compound was *N*-methoxycarbonylphenylalanine.

The second step is the conversion of **4** to **5**. We had already found that heating cyclic α -methoxycarbamates in the presence of a small amount of NH₄Cl gave the corresponding encarbamates in good yields^{7a} but the reaction conditions were not always satisfactory for acyclic α -methoxycarbamates **4a-f**, affording **5a-f** in low yields (Table 2). On the other hand, we found herein that the treatment of **4a-f** with NaH in THF gave **5a-f** in good yields (Entries 1~4), while the yields of **5e, f** were comparable with those obtained by acid-catalyzed procedures (Entries 5 and 6).


Table 2. Elimination of Methanol from α -Methoxycarbamates **4a-f**

Entry	α -Methoxycarbamate			Enecarbamate 5a-f	Yield (%) (<i>cis/trans</i>)	
	4a-f	R	R'		$\Delta/\text{NH}_4\text{Cl}^a$	NaH ^b
1	4a	Me	H	5a	33 (13/87)	56 (48/52)
2	4b	Me	Me	5b	29 (-)	84 (-)
3	4c	<i>i</i> -Pr	H	5c	52 (0/100)	73 (0/100)
4	4d	Me(CH ₂) ₅	H	5d	57 (40/60)	97 (45/55)
5	4e	CO ₂ Me	H	5e	80 (79/21)	72 (77/23)
6	4f	Ph	H	5f	60 (12/88)	60 (17/83)

^a 5h reflux in toluene with a cat. amount of NH₄Cl.


^b 1h reflux in THF with 3.0 equiv. of NaH.

The third step is the asymmetric introduction of a hydroxy group into the β position to the nitrogen atom of **5a-f**. Before the asymmetric hydroxylation, we examined the possibility of hydroboration of **5a-f** using borane in THF (eq 2), and found that these enecarbamates **5** except **5e** could be converted to **6a-f** (Table 3).

Table 3. Hydroboration of Enecarbamates **5a-f**

Entry	Enecarbamate 5a-f	R	R'	β -Hydroxyamine 6a-f	Yield (%)
1	5a	Me	H	6a	50
2	5b	Me	Me	6b	30
3	5c	<i>i</i> -Pr	H	6c	55
4	5d	Me(CH ₂) ₅	H	6d	69
5	5e	CO ₂ Me	H	6e	0
6	5f	Ph	H	6f	76

The reason why **5e** did not give **6e** by hydroboration is explainable in terms of the selective formation of a borane enolate **7**. In fact, the reaction of **5e** with borane gave a hydrogenation product, methyl 3-(*N*-methoxycarbonylamino)propionate **3e**, in a quantitative yield (eq 3).



Optically active β -hydroxyamines are our final targets. Since it has been known that asymmetric hydroboration of alkenes largely depends on the stereochemistry,^{8, 11} *cis* and *trans* stereoisomers of enecarbamates **5a, c, d, f** were isolated by column chromatography. At first, the *trans* stereoisomers *trans*-**5** were subjected to the hydroboration using (+)-Ipc-BH₂ (eq 4). The results are shown in Table 4 in which optically active **6c, d, f** were obtained with moderate enantiomeric excesses (ee's) (entries 2-4 in Table 4), while the ee of **6a** was low (entry 1 in Table 4).





Table 4. Asymmetric Hydroboration of *trans*-**5a, c, d, f**


Entry	Enecarbamate <i>trans</i> - 5	R	β -Hydroxyamine 6	Yield (%)	$[\alpha]_D^{23}$ (c in MeOH)	%ee
1	5a	Me	(<i>S</i>)-(+)- 6a	56	+1.88° (1.33)	7
2	5c	<i>i</i> -Pr	(<i>S</i>)-(+)- 6c	56	+30.90° (1.10)	66
3	5d	Me(CH ₂) ₅	(<i>S</i>)-(+)- 6d	92	+3.35° (1.40)	60
4	5f	Ph	(<i>S</i>)-(+)- 6f	94	+2.80° (0.75)	70

Those results are explainable by assuming intermediates **8** and **9** similar to those proposed in the hydroboration of simple *trans*-alkenes using (+)-Ipc-BH₂ (Scheme 2).⁸ Namely, plausible transition states leading to (*S*)- and (*R*)-**6** may be **8** and **9**, respectively, in which R group is disposed *anti* to the pinanyl group, and the difference of steric repulsion between R and the pinanyl group in **8** and **9** may reflect the ee's. In a case where R is a small group such as a methyl group, the ee was very low (entry 1 in Table 4), and relatively high ee's were observed in cases of larger R groups such as *i*-propyl, *n*-hexyl and phenyl groups (entries 2-4 in Table 4).





Scheme 2


In contrast, the hydroboration of *cis*-5f with (+)-Ipc·BH₂ resulted in a low ee even though R was a phenyl group (eq 5). Many possible transition states could be conceivable in the hydroboration of *cis*-5 with (+)-Ipc·BH₂⁸ to explain the reason why the low ee was observed in the hydroboration of *cis*-5f with (+)-Ipc·BH₂. The difference in relative stabilities between such many transition states would be less than that between 8 and 9.




Also, the reaction of *cis*- and *trans*-5f with (-)-Ipc₂BH gave low ee's (eqs 6 and 7). Since it has been known that (-)-Ipc₂BH gives higher stereoselectivities in hydroboration of simple *cis*-alkenes than simple *trans*-alkenes,¹¹ the methoxycarbonylamino group of *cis*- and *trans*-5f may be responsible for the low ee's, though it is not clear yet what effects the methoxycarbonylamino group brought about on the stabilities of the transition states.



Both (*S*)-2-hydroxypropylamine and (*S*)-2-hydroxy-2-phenylethylamine have been known to possess the positive optical rotations.^{12, 13} Since (*S*)-(+)-2-hydroxypropylamine **10** was easily available, the absolute configuration and ee of the obtained **6a** (entry 1 in Table 4) were determined by comparison with authentic (*S*)-(+)-**6a** prepared from **10** (eq 8), and the absolute configuration of the obtained **6f** (entry 4 in Table 4) to be *S* was determined by deprotecting **6f** to 2-hydroxy-2-phenylethylamine followed by measuring the optical rotation.¹⁴ The fact that the absolute configuration of **6a, f** was *S* supports our working hypothesis (Scheme 2) in which the main reaction proceeded *via* an intermediate **8**.



The ee's of **6c, d, f** were determined on the bases of ¹H NMR of Mosher's esters **11c, d, f** which were prepared by condensation of **6c, d, f** with Mosher's acid in the presence of DCC and DMAP (eq 9).




	R	Yield (%)	%de
11c	<i>i</i> -Pr	87	66
11d	Me(CH ₂) ₅	86	60
11f	Ph	85	70

Although the absolute configurations of **6c, d** could not be determined directly, they were estimated to be *S* by assuming that the hydroboration of **5c, d** might proceed through intermediates **8** in a similar way to that of

5a, f giving (*S*)-(+)-**6a, f**.

As shown in eqs 4-7, optically active β -hydroxyamines could be obtained with moderate ee's by the reactions of *trans*-**5** with (+)-Ipc·BH₂ as a chiral hydroboration reagent. Usefulness of our method was demonstrated by the transformation of (*S*)-(+)-**6f** to resedine **12**, an alkaloid isolated from *Redeeda Lutola* (eq 10).¹⁵



Further studies on the application of our method to more complexed primary amines and on the improvement of ee are under investigation.

Experimental

Melting points were determined on a Yanagimoto micro-melting point apparatus and are uncorrected. IR spectra were measured with JASCO 810 and Shimadzu FTIR-8100A spectrometers. ¹H NMR spectra were recorded on Varian Gemini-200 (200MHz), 300 (300MHz) and UNITY plus 500 (500MHz) spectrometers with tetramethylsilane as an internal reference and CDCl₃ as a solvent. Optical rotations were measured by JASCO DIP-370. Elemental analyses were performed at the Microanalytical Laboratory of the Center for Instrumental Analysis in Nagasaki University.

Materials. Carbamates **3a-e**^{7b, 16} and *N*-methoxycarbonylphenylalanine¹⁷ were known compounds. These compounds were prepared by conventional method; Methyl chlorocarbonate was added to the corresponding primary amines in aqueous NaOH. After usual workup, the carbamates were isolated by column chromatography. DCC(dicyclohexylcarbodiimide), DMAP[4-(dimethylamino)pyridine] and other chemical reagents were commercially available and used without further purification.

Electrochemical Oxidation of Carbamates 3a-e and *N*-Methoxycarbonylphenylalanine. The oxidation was carried out according to the reported procedure.^{7,10} A typical procedure for the electrochemical oxidation of **3a-e** was as follows. In an undivided cell equipped with platinum electrodes (2cm x 2cm), a solution of **3d** (6.0g, 32mmol) in methanol (80 mL) containing tetraethylammonium tetrafluoroborate (1.0g, 4.6mmol) was electrolyzed at 300mA for 23hrs (8.0 F/mol). During the electrolysis, the solution was kept at -10°C by standing the cell in a cooling bath. After the electrolysis, the solution was added to water and the

organic portion was extracted with ether. The ethereal solution was dried over MgSO_4 , and then the solvent was evaporated *in vacuo* to give a residue, which was subjected on a column chromatography to isolate **4d**.

***N*-Methoxycarbonyl-1-methoxypropylamine (4a).**

76% Yield; oil; IR (neat) 3325, 2950, 2850, 1710, 1530, 1460, 1360, 1280, 1240, 1200, 1150, 1090, 1060, 1015, 960, 890 cm^{-1} ; ^1H NMR (CDCl_3) δ 0.94 (t, $J=7.5$ Hz, 3H), 1.49-1.76 (m, 2H), 3.36 (s, 3H), 3.70 (s, 3H), 4.70-5.05 (m, 2H).; Anal. Calcd for $\text{C}_6\text{H}_{13}\text{NO}_3$: C, 48.97; H, 8.90; N, 9.52. Found: C, 49.38; H, 8.52; N, 9.54.

***N*-Methoxycarbonyl-1-methoxy-2-methylpropylamine (4b).^{7b}**

63% Yield; mp 154-157°C (ethyl acetate/hexane); IR (KBr) 2960, 1735, 1530, 1460, 1300, 1240, 1200, 1090, 1035 cm^{-1} ; ^1H NMR (CDCl_3) δ 0.92 (d, $J=6.6$ Hz, 3H), 0.95 (d, $J=6.6$ Hz, 3H), 1.76-1.90 (m, 1H), 3.34 (s, 3H), 3.70 (s, 3H), 4.54-4.66 (m, 1H), 4.92-5.00 (bs, 1H); Anal. Calcd for $\text{C}_7\text{H}_{15}\text{NO}_3$: C, 52.14; H, 9.38; N, 8.69. Found: C, 52.04; H, 8.92; N, 8.91.

***N*-Methoxycarbonyl-1-methoxy-3-methylbutylamine (4c).^{7b}**

58% Yield; oil; IR (neat) 3323, 2957, 1709, 1529, 1450, 1369, 1259, 1194, 1091, 1055, 970 cm^{-1} ; ^1H NMR (CDCl_3) δ 0.92 (d, $J=6.2$ Hz, 6H), 1.33-1.80 (m, 3H), 3.35 (s, 3H), 3.70 (s, 3H), 4.85-4.95 (bs, 2H); Anal. Calcd for $\text{C}_8\text{H}_{17}\text{NO}_3$: C, 54.84; H, 9.78; N, 7.99. Found: C, 54.58; H, 9.44; N, 7.89.

***N*-Methoxycarbonyl-1-methoxyoctylamine (4d).^{16c}**

86% Yield; oil; IR 3319, 2928, 2856, 1707, 1529, 1360, 1236, 1194, 1093 cm^{-1} ; ^1H NMR (CDCl_3) δ 0.88 (t, $J=6.9$ Hz, 3H), 1.20-1.77 (m, 12H), 3.35 (s, 3H), 3.70 (s, 3H), 4.62-5.08 (bs, 2H); Anal. Calcd for $\text{C}_{11}\text{H}_{23}\text{NO}_3$: C, 60.80; H, 10.67; N, 6.45. Found: C, 60.55; H, 10.45; N, 6.29.

Methyl 3-methoxy-3-[(methoxycarbonyl)amino]propionate (4e).

81% Yield; oil; IR (neat) 3320, 2950, 1740, 1695, 1550, 1440, 1380, 1318, 1275, 1210, 1170, 1105, 1180, 1035 cm^{-1} ; ^1H NMR (CDCl_3) δ 2.69 (m, 2H), 3.36 (s, 3H), 3.71 (s, 6H), 5.22 (dd, $J=4.1, 9.1$ Hz, 1H), 5.89 (bs, 1H); Anal. Calcd for $\text{C}_7\text{H}_{13}\text{NO}_5$: C, 43.97; H, 6.85; N, 7.33. Found: C, 43.83; H, 6.55; N, 7.32.

***N*-Methoxycarbonyl-1-methoxy-2-phenylethylamine (4f).**

81% Yield; mp 72-73°C (ether/hexane); IR (KBr) 3323, 2949, 1700, 1527, 1454, 1361, 1255, 1093, 1053, 1034 cm^{-1} ; ^1H NMR (CDCl_3) δ 2.93 (t, $J=5.4$ Hz, 2H), 3.34 (s, 3H), 3.66 (s, 3H), 4.92-4.97 (m, 1H), 5.09-5.19 (m, 1H), 7.36 (s, 5H); Anal. Calcd for $\text{C}_{11}\text{H}_{15}\text{NO}_3$: C, 63.14; H, 7.23; N, 6.69. Found: C, 63.53; H, 7.09; N, 6.43.

Preparation of Enecarbamates 5. The following procedure is representative.

i) Acid Catalysis Method; A solution of a mixture of **4d** (434 mg, 2 mmol) and NH_4Cl (100 mg) in toluene (15 mL) was refluxed for 5h. To the resulting solution was added brine (15 mL) and the organic layer was separated.

The organic solution was dried over MgSO_4 , and evaporated to give a residue. The residue was subjected to column chromatography on a silica gel using ethyl acetate/hexane (1/10) as an eluent to give *cis*-**5d** (84.4 mg, 0.46 mmol) and *trans*-**5d** (126.6 mg, 0.68 mmol) with a ratio of 40 to 60 in 57% yield. The *cis* isomers generally gave less polar than *trans* isomers.

ii) NaH Method; To a suspension of NaH (50% oily, 570 mg, 12 mmol) in THF (50 mL) was added a solution of **4d** (870 mg, 4.1 mmol) in THF (10 mL) at room temperature, and the reaction mixture was refluxed for 1h. The reaction mixture was quenched with an aqueous NH_4Cl solution (20 mL), and the organic portions were extracted with CH_2Cl_2 (50 mL x 2). The combined extracts were dried over MgSO_4 and evaporated to give a residue. The residue was subjected on column chromatographic purification to give *cis*-**5d** (333 mg, 1.8 mmol) and *trans*-**5d** (407 mg, 2.2 mmol) with a ratio of 45 to 55 in 97% yield. The yields of **5a-f** under the both conditions are summarized in Table 2.

***N*-Methoxycarbonyl-1-propenylamine (5a).**

cis-**5a**: oil; IR (neat) 3320, 2957, 1736, 1676, 1500, 1451, 1397, 1364, 1240, 1107, 1011, 776 cm^{-1} ; ^1H NMR (CDCl_3) δ 1.57 (d, $J=7.0$ Hz, 3H), 3.73 (s, 3H), 4.60-4.77 (m, 1H), 6.30-6.75 (m, 2H); Anal. Calcd for $\text{C}_5\text{H}_9\text{NO}_2$: C, 52.15; H, 7.88; N, 12.17. Found: C, 52.12; H, 7.78; N, 12.32.

trans-**5a**: mp 64-66 $^\circ\text{C}$ (ether); IR (KBr) 3289, 1728, 1680, 1536, 1440, 1379, 1300, 1242, 1121, 1040, 959, 749 cm^{-1} ; ^1H NMR (CDCl_3) δ 1.64 (dd, $J=1.5, 6.7$ Hz, 3H), 3.71 (s, 3H), 4.90-5.08 (m, 1H), 6.05-6.40 (bs, 1H), 6.35-6.55 (m, 1H); Anal. Calcd for $\text{C}_5\text{H}_9\text{NO}_2$: C, 52.15; H, 7.88; N, 12.17. Found: C, 52.48; H, 7.71; N, 11.98.

***N*-Methoxycarbonyl-2-methyl-1-propenylamine (5b).**

mp 52-55 $^\circ\text{C}$ (ethyl acetate/hexane); IR (KBr) 3360, 2930, 1700, 1510, 1450, 1380, 1350, 1240, 1070, 860 cm^{-1} ; ^1H NMR (CDCl_3) δ 1.59 (s, 3H), 1.68 (s, 3H), 3.71 (s, 3H), 5.80-6.25 (bs, 1H), 6.25 (d, $J=12.0$ Hz, 1H); Anal. Calcd for $\text{C}_6\text{H}_{11}\text{NO}_2$: C, 55.79; H, 8.58; N, 10.85. Found: C, 55.70; H, 8.34; N, 10.69.

***N*-Methoxycarbonyl-3-methyl-1-butenylamine (5c).**

trans-**5c**: oil; IR (neat) 3320, 2960, 2870, 1705, 1675, 1525, 1460, 1270, 1230, 1050, 950 cm^{-1} ; ^1H NMR (CDCl_3) δ 0.98 (d, $J=6.7$ Hz, 6H), 2.29 (dq, $J=6.9, 13.5$ Hz, 1H), 3.71 (s, 3H), 4.99 (dd, $J=7.1, 14.1$ Hz, 1H), 6.07-6.61 (bs, 1H), 6.41 (dd, $J=6.6, 13.9$ Hz, 1H); Anal. Calcd for $\text{C}_7\text{H}_{13}\text{NO}_2$: C, 58.72; H, 9.15; N, 9.78. Found: C, 58.55; H, 8.98; N, 9.53.

***N*-Methoxycarbonyl-1-octenylamine (5d).**

cis-**5d**: oil; IR (neat) 3320, 2925, 2860, 1710, 1675, 1515, 1460, 1350, 1240, 1060 cm^{-1} ; ^1H NMR (CDCl_3) δ 0.88 (t, $J=7.1$ Hz, 3H), 1.25-1.39 (m, 8H), 1.92 (t, $J=8.9$ Hz, 2H), 3.73 (s, 3H), 4.62 (t, $J=7.8$ Hz, 1H), 6.16-6.37 (bs, 1H), 6.43 (d, $J=9.8$ Hz, 1H); Anal. Calcd for $\text{C}_{10}\text{H}_{19}\text{NO}_2$: C, 64.82; H, 10.34; N, 7.56. Found: C, 65.16; H, 10.24; N, 7.29.

trans-5d: mp 30-31°C (ether/hexane); IR (KBr) 3300, 2920, 2850, 1695, 1535, 1460, 1300, 1240, 1055, 955 cm⁻¹; ¹H NMR (CDCl₃) δ 0.88 (t, *J*=7.0 Hz, 3H), 1.24-1.35 (m, 8H), 1.98 (t, *J*=13.6 Hz, 2H), 3.70 (s, 3H), 4.99 (dt, *J*=13.9, 7.3 Hz, 1H), 6.08-6.26 (bs, 1H), 6.44 (d, *J*=12.4 Hz, 1H); Anal. Calcd for C₁₀H₁₉NO₂: C, 64.82; H, 10.34; N, 7.56. Found: C, 65.04; H, 10.29; N, 7.46.

Methyl 3-[(methoxycarbonyl)amino]acrylate (5e).

cis-5e: oil; IR (neat) 3325, 2950, 1745, 1695, 1640, 1630, 1500, 1440, 1380, 1370, 1200, 1050, 1000, 960 cm⁻¹; ¹H NMR (CDCl₃) δ 3.62 (s, 3H), 3.69 (s, 3H), 4.96 (d, *J*=9.0 Hz, 1H), 7.16 (dd, *J*=9.0, 12.0 Hz, 1H), 9.64 (bs, 1H); Anal. Calcd for C₆H₉NO₄: C, 45.28; H, 5.70; N, 8.88. Found: C, 45.48; H, 5.62; N, 8.76.

trans-5e: mp 173-174°C (ether); IR (KBr) 3275, 1745, 1690, 1620, 1530, 1440, 1320, 1265, 1230, 1200, 1155, 1080, 1120, 1000, 950 cm⁻¹; ¹H NMR (CDCl₃) δ 3.59 (s, 3H), 3.69 (s, 3H), 5.39 (d, *J*=14.5 Hz, 1H), 7.69 (dd, *J*=14.5, 11.0 Hz, 1H), 9.72 (d, *J*=11.0 Hz, 1H); Anal. Calcd for C₆H₉NO₄: C, 45.28; H, 5.70; N, 8.88. Found: C, 45.18; H, 5.52; N, 8.77.

N-Methoxycarbonyl-2-phenylethenylamine (5f).

cis-5f: oil; IR (neat) 3302, 1734, 1701, 1670, 1599, 1545, 1446, 1336, 1325, 1309, 1288, 1259, 1055, 954 cm⁻¹; ¹H NMR (CDCl₃) δ 3.74 (s, 3H), 5.64 (d, *J*=9.3 Hz, 1H), 6.71 (t, *J*=10.3 Hz, 1H), 6.95-7.00 (bs, 1H), 7.34-7.45 (m, 5H); Anal. Calcd for C₁₀H₁₁NO₂: C, 67.78; H, 6.26; N, 7.90. Found: C, 68.28; H, 6.34; N, 7.71.

trans-5f: mp 122-124°C (ether/hexane); IR (KBr) 3298, 1736, 1701, 1677, 1545, 1336, 1325, 1288, 1259, 1055, 956 cm⁻¹; ¹H NMR (CDCl₃) δ 3.76 (s, 3H), 5.96 (d, *J*=14.6 Hz, 1H), 6.64-6.70 (bs, 1H), 7.16 (dd, *J*=3.4, 14.6 Hz, 1H), 7.27-7.40 (m, 5H); Anal. Calcd for C₁₀H₁₁NO₂: C, 67.78; H, 6.26; N, 7.90. Found: C, 67.56; H, 6.22; N, 7.85.

Hydroboration of Enecarbamats 5 with BH₃ · THF Complex. The following procedure is representative. To a solution of **5d** (370 mg, 2 mmol) in THF (2 mL) was injected BH₃ · THF complex (4 mL, 4 mmol) at 0°C. After the solution was stirred at room temperature for 1h and refluxed for 1h, the reaction mixture was oxidized with 10% NaOH (1 mL) and 30% H₂O₂ (2 mL) at room temperature, and then the reaction mixture was stirred for 12h at room temperature. THF was removed under a reduced pressure and the residue was extracted with CH₂Cl₂ (50 mL x 3). The combined extracts were dried over MgSO₄ and evaporated. The residual oil was purified by column chromatography on a silica gel using ethyl acetate/hexane (1/1) as an eluent to afford *N*-methoxycarbonyl-2-hydroxyoctylamine (**6d**) (280 mg, 1.4 mmol) in 69% yield. The yields of **6a-f** are summarized in Table 3.

N-Methoxycarbonyl-2-hydroxypropylamine (6a).

oil; IR (neat) 3400, 2936, 2880, 2361, 2342, 2150, 1723, 1538, 1450, 1380, 1264, 1196, 1115, 1017, 930 cm⁻¹; ¹H NMR (CDCl₃) δ 1.18 (d, *J*=6.3 Hz, 3H), 2.25-2.55 (bs, 1H), 2.97-3.14 (m, 1H), 3.25-3.40 (m, 1H), 3.68 (s, 3H), 3.85-4.00 (m, 1H), 5.10-5.30 (bs, 1H); Anal. Calcd for C₅H₁₁NO₃: C, 45.10; H, 8.33; N, 8.88.

10.52. Found: C, 44.83; H, 7.99; N, 10.01.

***N*-Methoxycarbonyl-2-hydroxy-2-methylpropylamine (6b).**

oil; IR (neat) 3350, 2970, 1700, 1530, 1460, 1370, 1255, 1150, 1040 cm^{-1} ; $^1\text{H NMR}$ (CDCl_3) δ 1.23 (s, 6H), 1.91-2.12 (bs, 1H), 3.18 (d, $J=6.3$ Hz, 2H), 3.69 (s, 3H), 4.93-5.24 (bs, 1H); Anal. Calcd for $\text{C}_6\text{H}_{13}\text{NO}_3$: C, 48.97; H, 8.90; N, 9.52. Found: C, 49.32; H, 8.89; N, 9.36.

***N*-Methoxycarbonyl-2-hydroxy-3-methylbutylamine (6c).**

oil; IR (neat) 3400, 2964, 1734, 1556, 1280, 1150, 1082, 958 cm^{-1} ; $^1\text{H NMR}$ (CDCl_3) δ 0.95 (d, $J=6.6$ Hz, 6H), 1.68 (q, $J=6.7$ Hz, 1H), 2.08-2.32 (bs, 1H), 3.08 (t, $J=9.6$ Hz, 1H), 3.41 (d, $J=7.1$ Hz, 1H), 3.69 (s, 3H), 4.96-5.27 (bs, 1H); Anal. Calcd for $\text{C}_7\text{H}_{15}\text{NO}_3$: C, 52.16; H, 9.38; N, 8.69. Found: C, 52.43; H, 9.15; N, 8.48.

***N*-Methoxycarbonyl-2-hydroxyoctylamine (6d).**

oil; IR (neat) 3350, 2930, 2810, 1710, 1540, 1460, 1375, 1270, 1095 cm^{-1} ; $^1\text{H NMR}$ (CDCl_3) δ 0.88 (t, $J=6.5$ Hz, 3H), 1.18-1.54 (m, 10H), 1.67 (d, $J=2.5$ Hz, 1H), 2.12-2.26 (bd, 1H), 3.37 (d, $J=3.4$ Hz, 1H), 3.04 (d, $J=5.7$ Hz, 1H), 3.68 (s, 3H), 3.62-3.80 (m, 1H); Anal. Calcd for $\text{C}_{10}\text{H}_{21}\text{NO}_3$: C, 59.09; H, 10.41; N, 6.89. Found: C, 59.10; H, 10.21; N, 6.86.

***N*-Methoxycarbonyl-2-hydroxy-2-phenylethylamine (6f).**

mp 94-95°C (ether); IR (KBr) 3350, 3070, 3040, 2950, 1700, 1550, 1453, 1250, 1196, 1154, 1094, 1067, 1001, 831 cm^{-1} ; $^1\text{H NMR}$ (CDCl_3) δ 2.68-2.78 (bs, 1H), 3.24-3.40 (m, 1H), 3.48-3.65 (m, 1H), 3.68 (s, 3H), 4.80-4.89 (m, 1H), 5.00-5.18 (bs, 1H), 7.30-7.39 (m, 5H); Anal. Calcd for $\text{C}_{10}\text{H}_{13}\text{NO}_3$: C, 61.53; H, 6.71; N, 7.17. Found: C, 61.51; H, 6.65; N, 7.18.

Hydroboration of Enecarbamates 5 with (+)-Monoisopinocampheylborane [(+)-Ipc·BH₂]. Asymmetric Synthesis of (*S*)-(+)-*N*-Methoxycarbonyl-2-hydroxy-2-phenylethylamine [(*S*)-(+)-6f]. The following procedure is representative. A solution of (+)-Ipc·BH₂ in THF (10 mL) was prepared from (Ipc·BH₂)₂·TMEDA complex (530 mg, 1.28 mmol) and BF₃·OEt₂ (0.32 mL, 2.56 mmol) according to a general procedure.¹¹ To the solution, a solution of *trans*-5f (114 mg, 0.64 mmol) in THF (5 mL) was injected at 0°C under a nitrogen atmosphere. After the resulting solution was stirred at room temperature for 1h and then refluxed for 1h, 10% NaOH (1 mL) and 30% H₂O₂ (2 mL) were successively added at room temperature. After usual workup, the chromatographic purification of the residue on a silica gel (ethyl acetate/hexane=1/5 as an eluent) gave (*S*)-(+)-*N*-methoxycarbonyl-2-hydroxy-2-phenylethylamine 6f (124 mg, 0.63 mmol) in 94% yield. The optical rotation was $[\alpha]_D^{25} +2.80^\circ$ (c 0.75 in MeOH). The ee of this 6f (70% ee) was determined on the basis of the $^1\text{H NMR}$ spectrum of Mosher's ester 11f, which showed the benzylic proton signals at δ 5.95-6.02 (m) and 6.04-6.08 (m) in an integral ratio of 85 to 15.

(*S*)-(+)-*N*-Methoxycarbonyl-2-hydroxypropylamine [(*S*)-(+)-6a]. This optically active compound

6a was obtained in 56% yield by the hydroboration of *trans*-**5a** with (+)-Ipc·BH₂ carried out under conditions similar to those described for **6f**. The optical rotation of this **6a** showed $[\alpha]^{23}_{\text{D}} +1.88^{\circ}$ (c 1.33 in MeOH). The ee of this **6a** (7% ee) was determined by comparison with authentic sample prepared from commercially available (*S*)-(+)-2-hydroxypropylamine **10** (95% ee) as described below.

(S)-(+)-*N*-Methoxycarbonyl-2-hydroxy-3-methylbutylamine [(S)-(+)-6c]. This optically active compound was obtained in 56% yield by the hydroboration of *trans*-**5c** with (+)-Ipc·BH₂ carried out under conditions similar to those described for **6f**. The optical rotation of this **6c** showed $[\alpha]^{23}_{\text{D}} +30.90^{\circ}$ (c 1.10 in MeOH). The ee of this **6c** (66% ee) was determined on the basis of the ¹H NMR spectrum of Mosher's ester **11c** in which the methoxy proton signals appeared at δ 3.53 (s) and 3.56 (s) in an integral ratio of 83 to 17.

(S)-(+)-*N*-Methoxycarbonyl-2-hydroxyoctylamine [(S)-(+)-6d]. This optically active compound was obtained in 92% yield by the hydroboration of *trans*-**5d**. The optical rotation of this **6d** showed $[\alpha]^{23}_{\text{D}} +3.35^{\circ}$ (c 1.40 in MeOH). The ee of this **6d** (60% ee) was determined on the basis of the ¹H NMR spectrum of Mosher's ester **11d** in which the methoxy proton signals appeared at δ 3.50 (s) and 3.67 (s) in an integral ratio of 80 to 20.

Hydroboration of Enecarbamates 5f with (-)-Diisopinocampheylborane [(-)-Ipc₂·BH]. A solution of (-)-Ipc₂·BH in THF (10 mL) was prepared from (+)- α -pinene (1.02 g, 6.6 mmol) and BH₃·Me₂S (0.34 mL, 3.3 mmol) according to a general method.¹¹ To the solution, a solution of *trans*-**5f** (0.33g, 1.85 mmol) in THF (5 mL) was injected at 0°C under nitrogen atmosphere. After 1h at 0°C, the solution was allowed to warm up and then the solution was refluxed for 1h. Oxidation with 10% NaOH (1 mL) and 30% H₂O₂ (1 mL) gave (*R*)-(-)-**6f** (44 mg, 0.22 mmol) in 12% yield after usual workup. The optical rotation of this **6f** was $[\alpha]^{23}_{\text{D}} -1.22^{\circ}$ (c 1.6 in MeOH). So, the ee of this **6f** was determined to be 31% by comparison of the optical rotation of (*S*)-(+)-**6f** prepared with (+)-Ipc·BH₂. Hydroboration of *cis*-**5f** was also carried out according to the method described above. The optical rotation of thus obtained **6f** was $[\alpha]^{23}_{\text{D}} -0.23^{\circ}$ (c 0.86 in MeOH), and the ee was 6%.

Synthesis of (S)-(+)-*N*-Methoxycarbonyl-2-hydroxypropylamine [(S)-(+)-6a]. To a solution of commercially available (*S*)-(+)-2-hydroxypropylamine (*S*)-(+)-**10** (95% ee, 0.15g, 2.0 mmol) in 20% aqueous NaOH (2 mL), methyl chloroformate (0.23 mL, 3.0 mmol) was added with stirring over 10min, and then the solution was stirred overnight at room temperature. The resulting solution was extracted with ether (20 mL x 2), and the combined extracts were dried over MgSO₄ and evaporated to give (*S*)-(+)-**6a** (oil, 0.23g, 1.73 mmol) in 90% yield. Optical rotation of this compound was $[\alpha]^{23}_{\text{D}} +25.98^{\circ}$ (c 3.16 in MeOH).

Deprotection of (S)-(+)-6f. The deprotection of (*S*)-(+)-**6f** was carried out according to a reported method.¹⁸

Preparation of Mosher's Esters 11c, d, f. To a solution of (*S*)-(+)-**6c, d, f** (0.41 mmol) in CH_2Cl_2 (1 mL), a solution of DCC (0.25g, 1.23 mmol), (*S*)-(-)-MTPA acid (0.19g, 0.82 mmol), and DMAP (0.1g, 0.82 mmol) in CH_2Cl_2 (1 mL) was added at room temperature. The mixture was stirred for 30min, and then allowed to stand for 8h at room temperature. Chromatographic purification on a silica gel using ethyl acetate/hexane (1/3) as an eluent gave Mosher's esters **11c, d, f** in 87%, 86%, 85% yields, respectively.

11c: oil; IR (neat) 1747 cm^{-1} ; $^1\text{H NMR}$ (CDCl_3) δ 0.90 and 0.91 (d and d, $J=6.9\text{ Hz}$, 4.98H) 0.96(t, $J=6.6\text{ Hz}$, 1.02H) 1.58-1.59 (m, 0.17H) 1.59-1.62 (m, 0.83H) 1.92-2.00 (m, 1H) 3.21-3.28 (m, 0.17H) 3.33-3.39 (m, 0.83H) 3.53 (s, 2.49H) 3.56 (s, 0.51H) 3.65-3.68 (bs, 2.49H) 3.63-3.65 (bs, 0.51H) 4.68-4.74 (bs, 0.83H) 4.52-4.57 (bs, 0.17H) 4.98-5.04 (m, 1H) 7.38-7.43 (m, 3H) 7.52-7.58 (m, 2H); Anal. Calcd for $\text{C}_{17}\text{H}_{22}\text{NO}_5\text{F}_3$: C, 54.11; H, 5.88; N, 3.71. Found: C, 54.40; H, 5.67; N, 3.50.

11d: oil; IR (neat) 1749 cm^{-1} ; $^1\text{H NMR}$ (CDCl_3) δ 1.83 (t, $J=6.9\text{ Hz}$, 3H) 1.16-1.21 (m, 8H) 1.50-1.70 (m, 2H) 3.18-3.24 (m, 0.2H) 3.30-3.38 (m, 0.8H) 3.44-3.50 (m, 1H) 3.50 (s, 2.4H) 3.67 (s, 0.6H) 3.64-3.70 (m, 3H) 4.52-4.60 (bs, 0.2H) 4.72-4.80 (bs, 0.8H) 5.10-5.20 (m, 1H) 7.38-7.45 (m, 3H) 7.53-7.58 (m, 2H); Anal. Calcd for $\text{C}_{20}\text{H}_{28}\text{NO}_5\text{F}_3$: C, 57.27; H, 6.73; N, 3.34. Found: C, 57.42; H, 6.87; N, 3.28.

11f: oil; IR (neat) 1749 cm^{-1} ; $^1\text{H NMR}$ (CDCl_3) δ 3.46 (m, 1H) 3.46 (s, 0.45H) 3.52 (s, 2.55H) 3.62-3.68 (m, 1H) 3.65 (s, 0.45H) 3.67 (s, 2.55H) 4.62-4.72 (bs, 0.15H) 4.81-4.90 (bs, 0.85H) 5.95-6.02 (m, 0.15H) 6.04-6.08 (m, 0.85H) 7.30-7.44 (m, 10H); Anal. Calcd for $\text{C}_{20}\text{H}_{20}\text{NO}_5\text{F}_3$: C, 58.39; H, 4.90; N, 3.40. Found: C, 58.14; H, 5.02; N, 3.36.

Synthesis of 5-Phenyl-2-oxazolidinone (12). To suspension of NaH (50% oily, 89 mg, 1.9 mmol) in THF (40 mL) was added a solution of (*S*)-(+)-**6f** (0.12 g, 0.62 mmol) in THF (10 mL) at room temperature and the reaction mixture was refluxed for 3h. The resulting solution was quenched with an aqueous NH_4Cl (10 mL), and the organic layer was extracted with ether (20 mL x 2). The combined extracts were dried over MgSO_4 and evaporated to give a residue. The residue was adsorbed on column of a silica gel, and the column was eluted with ethyl acetate/hexane (1/2) to afford **12** (67 mg, 0.41 mmol) in 66% yield. The optical rotation was $[\alpha]^{23}_{\text{D}} +10.0^\circ$ (c 1.0 in MeOH); mp $71-73^\circ\text{C}$ (ether/hexane); IR (KBr) 3276, 2890, 1950, 1880, 1719, 1489, 1458, 1426, 1370, 1238, 1076, 1026, 1001, 928 cm^{-1} ; $^1\text{H NMR}$ (CDCl_3) δ 3.54 (t, $J=7.8\text{ Hz}$, 1H), 3.98 (t, $J=8.4\text{ Hz}$, 1H), 5.35-5.50 (bs, 1H), 5.63 (t, $J=8.0\text{ Hz}$, 1H), 7.35-7.47 (m, 5H); Anal. Calcd for $\text{C}_9\text{H}_9\text{NO}_2$: C, 66.25; H, 5.56; N, 8.58. Found: C, 66.08; H, 5.56; N, 8.46.

Acknowledgment : This work was supported in part by a Grant-in-Aid for Scientific Research (B) (No. 07455364) from the Ministry of Education, Science and Culture, Japan.

References and Notes

† This paper is dedicated to Professor Dr. Hans J. Schäfer on the occasion of his 60th birthday.

1. *Dictionary of Antibiotics and Related Substances*, Bycroft, B. W., Ed.; Chapman and Hall: London, 1988.

2. Reviews, see: a) Ishizuka, T.; Kunieda, T. *Yuki Gosei Kagaku Kyokaishi*, **1991**, *49*, 118-127. b) Kemp, J. E. G. In *Comprehensive Organic Synthesis* Trost, B. M.; Fleming, I.; Ley S. V., Ed.; Pergamon Press: Oxford, 1991; Vol:7, p488-493. c) Ishizuka, T.; Kunieda, T. *Yakugaku Zasshi*, **1995**, *115*, 460-475. d) Ager, D. J.; Prakash, I.; Schaad, D. R. *Chem. Rev.*, **1996**, *96*, 835-875.
3. a) Sharpless, K. B.; Chong, A. O.; Oshima, K. *J. Org. Chem.*, **1976**, *41*, 177-179. b) Goralski, C. T.; Singaram, B.; Brown, H. C. *J. Org. Chem.*, **1987**, *52*, 4014-4019. c) Bartczak, A. W.; Sangaiah, R.; Kelman, D. J.; Toney, G. E.; Deterding, L. J.; Charles, J.; Marbury, G. D.; Gold, A. *Tetrahedron Lett.*, **1989**, *30*, 3251-3254.
4. Besse, P.; Veschambre, H.; Dickman, M.; Chenevert, R. *J. Org. Chem.*, **1994**, *59*, 8288-8291.
5. Hydroboration of enamines has been reported; Fisher, G. B.; Goralski, C. T.; Nicholson, L. W.; Hasha, D. L.; Zakett, D.; Singaram, B. *J. Org. Chem.*, **1995**, *60*, 2026-2034.
6. Synthetic methods of enamides and enecarbamates; a) Ruschig, H.; Fritsch, W.; Schmidt-Thome, J.; Haede, W. *Chem. Ber.*, **1955**, *88*, 883-894. b) Lenz, G. R. *Synthesis*, **1978**, 489-518, and references cited therein. c) Stille, J.K.; Becker, Y. *J. Org. Chem.*, **1980**, *45*, 2139-2145. d) Jendrzewski, S.; Steglich, W. *Chem. Ber.*, **1981**, *114*, 1337-1342. e) Slopianka, M.; Gossauer, A. *Liebigs Ann. Chem.*, **1981**, 2258-2265. f) Redeker, U.; Engel, N.; Steglich, W. *Tetrahedron Lett.*, **1981**, *22*, 4263-4264. g) Alper, H.; Mahatantila, C. P. *Organometal.* **1982**, *1*, 70-74. h) Schmidt, U.; Lieberknecht, A.; Griesser, H.; Bokens, H. *Tetraheron Lett.*, **1982**, *23*, 4911-4914. i) Tanaka, C.; Nasu, K.; Yamamoto, N.; Shibata, M. *Chem. Pharm. Bull.*, **1982**, *30*, 3374-3376. j) Schmidt, U.; Lieberknecht, A. *Angew. Chem.*, **1983**, *95*, 575. k) Schmitt, G.; Ebertz, W. *Angew. Chem.*, **1982**, *94*, 650-651. l) Brettle, R.; Shibib, S. M.; Wheeler, K. J. *J. Chem. Soc., Perkin Trans.1*, **1985**, 831-836. m) Couture, A.; Deniau, E.; Grandclaudon, P. *Tetrahedron Lett.*, **1993**, *34*, 1479-1482.
7. a) Shono, T.; Matsumura, Y.; Tsubata, K.; Sugihara, Y.; Yamane, S.; Kanazawa, T.; Aoki, T. *J. Am. Chem. Soc.*, **1982**, *104*, 6697-6703. b) Shono, T.; Matsumura, Y.; Tsubata, K. *Org. Synth.*, **1985**, *63*, 206-213.
8. Houk, K. N.; Rondan, N. G.; Wu, Y. D.; Metz, J. T.; Paddon-Row, M. N. *Tetrahedron*, **1984**, *40*, 2257-2274.
9. Starting *N*-methoxycarbonylated β -phenylethylamine was recovered.
10. Iwasaki, T.; Horikawa, H.; Matsumoto, K.; Miyoshi, M. *Bull. Chem. Soc. Jpn.*, **1979**, *52*, 826-830.
11. a) Smith, K. In *Organometallics in Synthesis*, Schlosser, M. Ed., John Wiley & Sons: Chichester, **1994**, 461-508. b) Brown, H. C.; Ramachandran, P.V. In *Advances in Asymmetric Synthesis*, Hassner, A. Ed., JAI Press, Greenwich, **1995**, Vol.1, 147-210.
12. Yahiro, N. *Chemistry Lett.*, **1982**, 1479-1480.
13. Pratesi, P.; Grassi, M. *Farmaco. Ed. sci.(Pavia)*, **1953**, *8*, 86-91; *Chem. Abstr.*, **1954**, *48*, 1995d.
14. The ee of **6f** was determined by a method described in eq 9.
15. Tadzhibaev, M. M.; Lutfullin, K. L.; Malikov, V. M.; Yunusov, S. Y. *Kim. Prir. Soedin*, **1976**, *2*, 270-271. A racemic form was described in the literature.
16. a) Kondo, K.; Sonoda, N.; Tsutsumi, S. *Chemistry Lett.*, **1972**, 373-374. b) Shono, T.; Matsumura, Y.; Yamane, S.; Kashimura, S. *Chemistry Lett.*, **1982**, 565-568. c) Shono, T.; Matsumura, Y.; Kashimura, S. *J. Org. Chem.*, **1983**, *48*, 3338-3340. d) Montagne, M. *Bull. Chem. Soc. France*, **1947**, 125-128.
17. Leuchs, H.; Geiger, W. *Chem. Ber.*, **1908**, *41*, 1721-1726.
18. Shono, T.; Matsumura, Y.; Uchida, K.; Tsubata, K.; Makino, A. *J. Org. Chem.*, **1984**, *49*, 300-304.