




with a cyclopentadienyl group (Cp) such as $\text{Me}(\text{MeO})\text{CpSi}^+$ (m/e 140, 31.9%) and $\text{Me}(\text{H})\text{-CpSi}^+$ (m/e 110, 12.3%) contain always deuterium, whereas those without a cyclopentadienyl group such as $\text{Me}_2(\text{MeO})\text{Si}^+$ (m/e 89, 100%) and Me_2SiH^+ (m/e 59, 25.8%) have no deuterium. These facts demonstrate that the retro ene reaction of 2 occurred only in the direction to form 1 which was successfully trapped by methanol.




The vacuum-flow co-pyrolysis of 2 with benzaldehyde gave 6-phenylfulvene (4)⁸ in 47% yield (after purification with TLC) together with cyclic oligomers of polydimethylsiloxane, 5 and 6. The formation of 5 and 6, evidenced by the GC-MS analysis of the pyrolysate, indicates 6-silafulvene as an intermediate in the reaction.⁹ Similarly, a benzene solution of 2 containing benzophenone was subjected to the flow pyrolysis through a quartz tube packed with quartz chips at 600°C to afford 6,6-diphenylfulvene (7)⁸ in 28% yield together with cyclic polydimethylsiloxanes, 5 and 6. It is well documented that a silaethene derivative reacts with a carbonyl compound to give a silaoxetane which decomposes to an olefin and a silanone.¹⁰ The latter is known to give 5 and 6. Therefore, these products are best explained by the following scheme of reactions.




The vacuum-flow pyrolysis of 2 at 650°C in the absence of a trapping reagent yielded a mixture of dimeric products (8a and 8b) of 1 in a 3:1 ratio in 60% yield.





Giving only one peak on both GLC and HPLC as far as we have examined, the mixture was not separated to each isomer.¹¹ However, ^1H NMR data¹² indicates clearly the structures of the abundant (8a) and less abundant (8b) isomers to be those indicated above. Thus, 8a shows two types of SiMe signals (δ -0.49 and 0.53) in an equal intensity together with the 2H peak assignable to that of the allylic hydrogen (δ 3.52), while four types of Si-Me signals (δ -1.38, 0.32, 0.47 and 0.51), accompanied by the resonance (δ 3.81) due to the corresponding allylic hydrogen, are observed for 8b. The anomalous up-field shift of one of the SiMe signals in 8b can be rationalized by diamagnetic shielding caused by the two cyclopentadienyl rings of a sandwich like arrangement. In the mass spectrum, the mixture shows a molecular ion at m/e 244 but lacks the peak at m/e 122 which corresponds to the monomeric structure.

Furthermore, the following chemical transformation supports the structures. Thus, ethanolysis of the mixture of 8a and 8b gave 11 (38%),¹³ 12 (19%),¹⁴ 13 (15%)¹⁵ and 14 (11%).¹⁶



The mechanism of the formation of 8a and 8b is not always clear. However, it is known that silaethene dimerizes rapidly to 1,3-disilacyclobutane¹⁰ and 1,5-silicon shift occurs rapidly in a silylcyclopentadiene system.¹⁷ Therefore, the possible route to the products from 1 may be one shown above in which a 1,3-disilacyclobutane (9) is a key intermediate. The stereochemical consequence of the dimers is of considerable interest and we will discuss the matter in a forthcoming paper.

Acknowledgment: We thank Toshiba Silicone Co., Ltd. for a gift of chlorosilanes. The work was supported in part by the Mitsubishi Foundation and the Ito Science Foundation.

References and Notes

- 1 Chemistry of Organosilicon Compounds, 140.
- 2 (a) T. J. Barton, D. S. Banasiak, *J. Am. Chem. Soc.*, **99**, 5199 (1977); (b) T. J. Barton, G. T. Burns, *ibid.*, **100**, 5246 (1978); (c) C. L. Kreil, O. L. Chapman, G. T. Burns, T. J. Barton, *ibid.*, **102**, 841 (1980); (d) H. Bock, R. A. Bowling, B. Solouki, T. J. Barton, G. T. Burns, *ibid.*, **102**, 429 (1980); (e) B. Solouki, P. Rosmus, H. Bock, and G. Maier, *Angew. Chem. Int. Ed. Engl.*, **19**, 51 (1980); (f) G. Mair, G. Mihm and H. P. Reisenauer, *ibid.*, **19**, 52 (1980); (g) G. Märkl and P. Hofmeister, *ibid.*, **18**, 789 (1979).
- 3 T. J. Barton and G. T. Burns, *J. Organometal. Chem.*, **179**, C17 (1979).
- 4 E. Block and L. K. Revelle, *J. Am. Chem. Soc.*, **100**, 1630 (1978).
- 5 Y. A. Ustynyuk, P. I. Zakharov, A. A. Azizov, G. A. Shchembelov and I. P. Glorizov, *J. Organometal. Chem.*, **16**, 195 (1975).
- 6 2; bp 31.5~33°C/0.6mmHg; ^1H NMR (δ in CCl_4) -0.05 (6H, s, Si- CH_3), 1.56 (2H, d, $J=8.1\text{Hz}$, Si- CH_2), 3.02 (1H, bs, SiCH) 4.79~5.02 (2H, m, Si- CH_2 -CH= $\underline{\text{CH}_2}$), 5.53~6.10 (1H, m, Si- CH_2 -CH=) 6.55 (4H, bs, Cp-H); MS m/e (%) M^+ 164 (6.7), 123 (100), 99 (10.0), 95 (22.3). Preparation of 2 has been described by L. N. Maksimova, V. I. Koshutin and V. A. Smironov, *Zh. Obshch. Khim.*, **43**, 1198 (1973); *Chem. Abst.*, **79**, 53434 (1973).
- 7 Yu. A. Ustynyuk, P. I. Zakharov, A. A. Azizov, V. K. Potapov, and I. M. Pribytkova, *J. Organometal. Chem.*, **88**, 37 (1975).
- 8 J. H. Day, *Chem. Rev.*, **53**, 167 (1953).
- 9 As a member of cyclopentadienylmetals, cyclopentadienylsilane reacts with carbonyl compounds to afford a fulvene derivative, as shown typically in the following equation.

$$\text{Me}_3\text{Si}-\text{C}_5\text{H}_4 + \text{PhCHO} \xrightarrow{\Delta} \text{PhCH}=\text{C}_5\text{H}_4 + \text{Me}_3\text{SiOH}$$
- However, in this case, no 5 nor 6 can be detected in the product.
- 10 L. E. Gusel'nikov and N. S. Nametkin, *Chem. Rev.*, **79**, 529 (1979).
- 11 8a+8b; mp 108~109°C; MS m/e (%), M^+ 244 (54), 229 (100). High resolution MS, $\text{C}_{14}\text{H}_{20}\text{Si}_2$ calcd.: 244.1104. Found: 244.1110.
- 12 8a; ^1H NMR (δ in CDCl_3) -0.49 (6H, s, SiMe), 0.53 (6H, s, SiMe), 3.52 (2H, bs, HC-), 6.56~6.98 (6H, m, HC=). 8b; ^1H NMR (δ in CDCl_3) -1.38 (3H, s, SiMe), 0.32 (3H, s, SiMe), 0.47 (3H, s, SiMe), 0.51 (3H, s, SiMe), 3.81 (2H, bs, HC-), 6.56~6.98 (6H, m, HC=).
- 13 11; purified by means of preparative VPC. ^1H nmr spectrum is rather complicated due to its fluxional structure. MS m/e (%) M^+ 290 (3.7), 261 (16.7, M-29), 225 (53.9, M-65(Cp)), 123 (100, Me_2SiCp). High resolution MS, $\text{C}_{16}\text{H}_{26}\text{OSi}$ calcd.: 290.1523. Found: 290.1523.
- 14 R. L. Schaaf, P. T. Kan, and C. T. Lenk, *J. Org. Chem.*, **26**, 1790 (1961).
- 15 K. C. Frisch, *J. Am. Chem. Soc.*, **75**, 6050 (1953).
- 16 H. J. Fletcher and M. J. Hunter, *J. Am. Chem. Soc.*, **71**, 2918 (1949).
- 17 A. J. Ashe III, *J. Am. Chem. Soc.*, **92**, 1233 (1970).

(Received June 9, 1980)