

Synthesis of Indoles via  $6\pi$ -Electrocyclic Ring Closures of Trienecarbamates

Thomas J. Greshock and Raymond L. Funk\*

Department of Chemistry, Pennsylvania State University, University Park, Pennsylvania 16802

Received January 14, 2006; E-mail: rlf@chem.psu.edu

Indoles are arguably the most important of all the privileged structures in drug discovery.<sup>1</sup> Accordingly, the development of new methods for the synthesis of indoles has been extensively investigated.<sup>2</sup> These strategies can be broadly categorized based upon the order in which the individual aromatic substructures are introduced. By far the most common approach (A, Figure 1) involves beginning the synthesis with a substituted benzene ring. The venerable Fischer indole synthesis is an example of this strategy wherein the indicated C(3)–C(3a) and N–C(2) bonds are sequentially formed. A less common approach to substituted indoles is through the annelation of pyrroles (B, Figure 1), for example, the cycloaddition of 3-vinyl pyrroles with dienophiles. The third and least explored strategy for the construction of indoles is one in which both aromatic rings of the indole are constructed in consecutive bond forming processes from acyclic precursors. One of many hypothetical disconnections is formulated in structure C. Indeed, the majority of the examples<sup>2b</sup> of this overall strategy employ the implied intramolecular cycloaddition<sup>3</sup> as the key step in the construction of the substituted benzene ring. We report herein a new approach to indoles which exploits a Stille coupling to form the C(3a)–C(7a) bond of C followed by a novel electrocyclic ring closure of the resulting trienecarbamate which effects the connection of C(4) with C(5). Finally, condensation of a C(2) enolate upon a C(3) carbonyl completes the heterocycle construction.

The feasibility of this strategic plan was easily demonstrated and is detailed in Scheme 1. Thus, Stille coupling of  $\alpha$ -(tributylstannyl)-enecarbamate **1**<sup>4</sup> with 2-iodocyclohexenone proceeded smoothly to afford amidotriene **2**. More importantly, triene **2** was found to undergo an especially facile electrocyclic ring closure<sup>5</sup> (110 °C, 1 h) to furnish cyclohexadiene **3**. This closure is most likely facilitated by a push/pull-type mechanism of the hydrogen bonded enecarbamate functionality with the proximal carbonyl group en route to the resonance stabilized vinylogous imide product.<sup>6</sup> Indeed, we found that the electrocyclic ring closure of a triene analogous to **2** that lacks the carbonyl group and possesses an *N*-methyl substituent requires higher temperatures and prolonged reaction times (120 °C, 12 h).<sup>7</sup> Not surprisingly, the electron-rich cyclohexadiene **3** can be easily oxidized to the protected aniline **4** by DDQ. This aromatization can be conveniently accomplished in the same pot once the electrocyclization is observed to be complete by TLC.

Removal of the BOC group with TFA was uneventful, and a reductive amination of the resultant aniline with glyoxylic acid provided acid **5**. It was hoped that this compound would cyclize to an indole upon subjection to conditions (Ac<sub>2</sub>O, NEt<sub>3</sub>, 130 °C) that were reported by Răileanu and co-workers<sup>8a</sup> for the cyclization of the analogously substituted *ortho*-aminobenzaldehyde to *N*-acetylindole. Indeed, this underutilized transformation proceeded smoothly to deliver the desired *N*-acetylindole **6**. This cyclization could possibly be proceeding through a Perkin-type reaction of the mixed anhydride derivative of acid **5** or, more likely, through a münchnone intermediate generated from the acetylated derivative of aniline **5**.<sup>8b,c</sup>





Figure 1. Strategies for indole synthesis.


## Scheme 1




We next directed our attention to delineating the scope and generality of this new indole annelation. Our proof-of-principle example suggested that this method would be especially useful for constructing indoles that are bridged at the C(3) and C(4) positions, a substructure that complicates the synthesis of many indole natural products. To that end, the corresponding  $\alpha$ -iodocycloalkenones were coupled with stannane **1** to afford trienecarbamates **7** and **10** (Scheme 2), and these substrates were subjected to the previously described reaction sequence to afford the heretofore unknown indole-containing ring systems **9** and **12**, respectively. The triene-carbamate **13** was also prepared from a methylated analogue of stannane **1** and transformed to indole **15** in order to demonstrate that additional substitution at C(5) is accessible by straightforward adaptation of this strategy.

It was also of interest to determine whether the Răileanu closure would tolerate the incorporation of substituents at N or C(2) of the

## Scheme 2



Scheme 3




product indole. With respect to the former question, the electrocyclic ring closure of trienecarbamate **16** (Scheme 3) proceeded as expected, although a longer reaction time (110 °C, 3 h) was required in comparison to that of triene **2**, presumably due to a lack of hydrogen bonding of the enecarbamate with the proximal carbonyl. After oxidation, removal of the BOC protecting group, and alkylation with methyl iodoacetate, the resultant ester was hydrolyzed with 4 M NaOH. To our surprise, we obtained clean conversion to the *N*-methyl indole **18** upon acidic workup with HCl. Preparation of a 2-substituted indole was accomplished through alkylation of the previously prepared aniline **19** (Scheme 1) with methyl  $\alpha$ -bromophenylacetate to afford ester **20**. Saponification of the ester, followed by subjection of the resultant acid to the Răileanu conditions afforded the desired 2-phenyl indole **21**, albeit lacking the acetyl moiety, which suggests that a münchnone intermediate may not be involved in this particular closure.


In addition to cyclic  $\alpha$ -iodoenones, we have also employed acyclic  $\alpha$ -iodoenones as the starting materials in the annelation sequence (Scheme 4). For example, the *cis*-phenyl ring did not diminish the yield (90%) of the Stille coupling reaction leading to trienecarbamate **22**. However, the electrocyclization of trienecarbamate **22** required a higher temperature, perhaps due to a more out-of-plane carbonyl in comparison to the conformationally locked cyclic trienecarbamates (Schemes 1–3). Subsequent transformation of **23** to the desired *N*-acetyl indole **24** was straightforward. Thermolysis of the  $\beta,\beta$ -disubstituted acyclic enone **25** provided the cyclohexadiene **27**. This result was not entirely unexpected since it is well-known that [1,7]-sigmatropic rearrangements are often competitive with 6 $\pi$ -electrocyclic closures, in this case providing the rearrangement product **26** followed by electrocyclic closure to **27**. Our standard protocol then furnished the 3,4,6-trisubstituted indole **29**.

We have also found that additional heterocyclic rings can be incorporated into the product indole by initiating the annelation

Scheme 4



Scheme 5



sequence with heterocyclic  $\alpha$ -iodoenones or  $\alpha$ -stannylenecarbamates (Scheme 5). Thus, electrocyclization of triene **30** proceeded smoothly with a negligible electronic/steric impact of the additional carbamate substituent to afford a cyclohexadiene that was oxidized (DDQ) in situ to the desired protected aniline **31**. Completion of the annelation afforded indole **32**, a substructure embodied in several biologically active natural products, such as the priansins.<sup>9</sup> Finally, the modularity of this indole construction method is showcased in the preparation of the unusual tetracyclic indole **35**, which originates from 4,4-dimethyl-2-iodocyclohexenone and the BOC-protected 3-vinyl-2-(trimethylstannyl)pyrroline.

In conclusion, we have shown that both of the aromatic rings of indoles can be constructed from readily available  $\alpha$ -haloenones and  $\alpha$ -(trialkylstannyl)enecarbamates using a five-step reaction sequence that features facile electrocyclic ring closures of trienecarbamates. The method may prove to be most useful for the preparation of indoles possessing complex or difficult substitution patterns. The syntheses of natural products of this type are underway.

**Acknowledgment.** We appreciate the financial support provided by the National Institutes of Health (GM28553).

**Supporting Information Available:** Spectroscopic data and experimental details for the preparation of all new compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.

## References

- (1) Horton, D. A.; Bourne, G. T.; Smythe, M. L. *Chem. Rev.* **2003**, *103*, 893.
- (2) (a) Sundberg, R. J. *Indoles*; Academic Press: San Diego, CA, 1996. (b) Gribble, G. W. *J. Chem. Soc., Perkin Trans. 1* **2000**, 1045.
- (3) Dunetz, J. R.; Danheiser, R. L. *J. Am. Chem. Soc.* **2005**, *127*, 5776 and ref 2 therein.
- (4)  $\alpha$ -(Tributylstannyl)enecarbamate **1** was prepared in three steps from the Overman diene (Overman, L. E.; Taylor, G. F.; Petty, C. B.; Jessup, P. J. *J. Org. Chem.* **1978**, *43*, 2164): (a) Protection with SEMCl, (b) metalation and stannylation with Bu<sub>3</sub>SnCl, and (c) deprotection of the SEM group. See Supporting Information for experimental procedures and spectral data.
- (5) (a) Okamura, W. H.; De Lera, A. R. In *Comprehensive Organic Synthesis*; Trost, B. M., Fleming, I., Paquette, L. A., Eds.; Pergamon Press: New York, 1991; Vol. 5, p 699. (b) For significantly slower (>200 °C) electrocyclic ring closures of 2,3-divinylpyrroles, see: Moskal, J.; van Stralen, R.; Postma, D.; van Leusen, A. M. *Tetrahedron Lett.* **1986**, *27*, 2173.
- (6) For a more pronounced acceleration of the electrocyclic closure of a related triene by this effect, see: Magomedov, N. A.; Ruggiero, P. L.; Tang, Y. *J. Am. Chem. Soc.* **2004**, *126*, 1624.
- (7) Nonetheless, this relatively facile electrocyclic ring closure indicates that an electron-donating substituent at C(3) of the triene is beneficial as has been previously suggested. See: (a) Spangler, C. W.; Jondahl, T. P.; Spangler, B. J. *J. Org. Chem.* **1973**, *38*, 2478. For the closure of a C(2) aminotriene, see: (b) Barluenga, J.; Merino, I.; Palacios, F. *Tetrahedron Lett.* **1990**, *31*, 6713. For calculations of substituent effects, see: (c) Guner, V. A.; Houk, K. N.; Davies, I. A. *J. Org. Chem.* **2004**, *69*, 8024 and references therein.
- (8) (a) Tighineanu, E.; Chiraleu, F.; Răileanu, D. *Tetrahedron* **1980**, *36*, 1385. For related ring closures leading to pyrroles, see: (b) Gabbutt, C. D.; Hepworth, J. D.; Heron, B. M.; Pugh, S. L. *J. Chem. Soc., Perkin Trans. 1* **2002**, 2799. For the generation of a similar münchnone under these conditions, see: (c) Padwa, A.; Gingrich, H. L.; Lim, R. J. *J. Org. Chem.* **1982**, *47*, 2447.
- (9) Cheng, J.; Ohizumi, Y.; Wälchli, M. R.; Hideshi, N.; Hirata, Y.; Sasaki, T.; Kobayashi, J. *J. Org. Chem.* **1988**, *53*, 4621.

JA0602820