

A Highly Regio- and Stereoselective Synthesis of α -Fluorinated Imides via Fluorination of Chiral Enamides


Yan-Shuang Xu,[†] Yu Tang,^{*,†} He-Jing Feng,[†] Ji-Tian Liu,[†] and Richard P. Hsung^{*,‡}

[†]School of Pharmaceutical Science and Technology, Key Laboratory for Modern Drug Delivery and High-Efficiency, Collaborative Innovation Center of Chemical Science and Engineering, Tianjin University, Tianjin, 300072, P. R. China

[‡]Division of Pharmaceutical Sciences, School of Pharmacy, and Department of Chemistry, University of Wisconsin, Madison, Wisconsin 53705, United States

S Supporting Information

ABSTRACT: A highly π -facial selective and regioselective fluorination of chiral enamides is described. The reaction involves an enantioselective fluorination exclusively at the electron-rich enamide olefin with N–F reagents such as Selectfluor and *N*-fluoro-benzenesulfonimide [NFSI] accompanied by trapping of the β -fluoro-iminium cationic intermediate with water. The resulting *N,O*-hemiacetal could be oxidized using Dess-Martin periodinane, leading to an asymmetric sequence for syntheses of chiral α -fluoro-imides and optically enriched α -fluoro-ketones.




The importance of organo-fluorine compounds has been abundantly validated through a broad range of applications in medicinal chemistry and drug development¹ as well as material² and agrochemical sciences.³ The incorporation of fluorine and/or fluorine-containing groups into an organic compound has often provided agents and materials with unique chemical, physical, and biological properties.⁴ One of the major synthetic challenges in fluoro-organic chemistry is to asymmetrically construct fluorinated stereogenic carbon centers. Differding⁵ demonstrated the first stoichiometric asymmetric fluorination of β -ketoester enolates with a chiral N–F (*N*-fluoroamine) reagent in 1988. Given the number of impressive examples of enantioselective fluorinations being reported over the past decade,⁶ asymmetric fluorinations represent an area of immense interest from the synthetic community. In relevance to our work, Davis⁷ reported an elegant synthesis of α -fluoro-imides using enolate derived from chiral imide **1** substituted with chiral oxazolidinone auxiliary⁸ (**1** \rightarrow **2** in Scheme 1). Our longstanding interest in the chemistry of enamides has drawn us to develop stereoselective fluorination methods using chiral enamides,⁹ which has remained elusive. Such an approach in combination with an oxidative process (**3** \rightarrow **4**) would represent a complementary approach to Davis' asymmetric approach to α -fluoro-imides.


With the nitrogen atom being substituted with an electron-withdrawing functionality, chiral enamides possess superior stability to their enamine counterparts, while maintaining excellent reactivities. This is evident by their recent emergence as another powerful chiral building block in organic synthesis.^{10,11} In particular, chiral enamides have been employed in highly regio- and/or stereoselective Diels–Alder cycloadditions,¹² [2 + 2] cycloadditions,¹³ cyclopropanation,¹⁴

Scheme 1. Stereoselective Synthesis of α -Fluoro-imides

(a) Davis' work:



(b) Our approach: Stereoselective fluorinations of chiral enamides?




epoxidation,¹⁵ and halo-etherification.¹⁶ We wish to report here a highly regio- and stereoselective fluorination of chiral enamides for the synthesis of chiral α -fluoro-imides and optically enriched α -fluoro-ketones.

Our efforts commenced with exploring the right fluorination conditions using chiral enamide **3a** as the testing enamide, and with Selectfluor,¹⁷ *N*-fluoropyridinium triflate [Py-F],¹⁸ and *N*-fluoro-benzenesulfonimide [NFSI]¹⁹ as the fluorinating agent. When reactions were carried out in CH₃CN at 40 °C, we found that it was faster with Selectfluor (entry 1 in Table 1) and much slower with Py-F (entry 2), and that NFSI was the best N–F reagent²⁰ in terms of diastereoselectivity, leading to α -fluoro-imides **4a** and **4a'** as an isomeric mixture in 57% yield after DMP oxidation (entry 3).²¹ We note here that attempts to directly

Received: December 13, 2014

Published: January 12, 2015

Table 1. Optimization of the Fluorination Conditions^a

entry	solvent	reagent ^a	time [h]	yield [%] ^b [dr] ^c
1	CH ₃ CN	Selectfluor	0.5	29 [57:43]
2	CH ₃ CN	Py-F	24	26 [61:39]
3	CH ₃ CN	NFSI	9.5	57 [89:11]
4	anhyd CH ₃ CN ^d	NFSI	8	<10 [85:15]
5	2.0 equiv H ₂ O in CH ₃ CN	NFSI	8	43 [89:11]
6	1% H ₂ O in CH ₃ CN	NFSI	12	62 [91:9]
7	2% H ₂ O in CH ₃ CN	NFSI	12	68 [91:9]
8	5% H ₂ O in CH ₃ CN	NFSI	31	29 [91:9]
9 ^e	2% H ₂ O in CH ₃ CN	NFSI	6	22 [71:29]

^aIn all reactions, 1.1 equiv of N-F reagent was added. Unless otherwise noted, reactions were run at 40 °C. DMP: Dess-Martin Periodinane; NFSI: *N*-fluoro-benzenesulfonimide; Py-F: *N*-fluoropyridinium triflate. ^bYields were determined by ¹H NMR analysis using mesitylene as the internal standard. ^cDiastereomeric ratios [dr] were determined using ¹H or/and ¹⁹F NMR spectroscopy. ^dCH₃CN distilled over CaH₂. ^eThe reaction temperature is 80 °C.

work up the crude fluorination mixture without any oxidation protocols failed to give us the desired *N,O*-hemiacetal products, or the corresponding α -fluoro-aldehydes via hydrolysis.

Most notably, H₂O plays a significant role in this reaction. The use of anhydrous CH₃CN did afford comparable diastereoselectivity, but the yield is very low (entry 4). The addition of 2.0 equiv, 1%, 2%, and 5% H₂O in the CH₃CN appeared to improve the efficiency (entries 5–8) with 2% H₂O being the most optimal (entry 7). On the other hand, when we increased the temperature from 40 to 80 °C, neither the *dr* ratio nor the yield was satisfactory.


To improve the diastereoselectivity, we examined a range of chiral auxiliaries as shown in Table 2. Enamide **3b** with the *i*-Pr-substituted Evans chiral auxiliary²² gave α -fluoro-imide **4b** in 58% yield with a 93:7 *dr* ratio. Fluorination of chiral enamide **3c** substituted with the Sibi auxiliary²³ afforded **4c** in 51% yield (entry 2). When using the Ph-substituted Evans auxiliary,

Table 2. Effect of the Chiral Auxiliary on Selectivity^a

entry	enamides	products	time [h]	yield [%] ^b [dr] ^c
1			14	58 [93:7]
2			12	51 [91:9]
3			12	72 [>95:5]
4			12	63 [>95:5]

^aReaction condition: 1.1 equiv of NFSI, 2% H₂O in CH₃CN, 40 °C; and then, DMP, NaHCO₃, CH₂Cl₂, rt. ^bIsolated yields. ^cDiastereomeric ratios [dr] were determined by ¹H or/and ¹⁹F NMR spectroscopy.

fluorinations of **3d** and *ent*-**3d** with NFSI at 40 °C in 2% H₂O in CH₃CN led to **4d** and *ent*-**4d**, respectively, in good yields and essentially as a single diastereomer (entries 3 and 4). X-ray structures of α -fluoro-imide **4b** and *ent*-**4d** allowed for unambiguous assignment of both stereochemical and structural integrity (Figure 1).

Figure 1. X-ray structures of α -fluoro-imides **4b** and *ent*-**4d**.

Success in finding chiral amides **4b** that can provide a high level of diastereoselectivity allowed us to broaden the scope of this fluorination significantly as shown in Table 3. For all examples in

Table 3. Stereoselective Fluorination of Chiral Enamides


entry	α -fluoro-imides: R =	time [h]	yield [%] ^a [dr] ^b
1	4-MeC ₆ H ₄ (6a)	10	67 [94:6]
2	4-MeOC ₆ H ₄ (6b)	12	42 [92:8]
3	4-NO ₂ C ₆ H ₄ (6c)	140	71 [>95:5]
4	4-FC ₆ H ₄ (6d)	12	61 [>95:5]
5	4-ClC ₆ H ₄ (6e)	12	65 [>95:5]
6	4-BrC ₆ H ₄ (6f)	14	69 [95:5]
7	2-ClC ₆ H ₄ (6g)	12	59 [>95:5]
8	<i>n</i> -Pent (6h)	11	53 [>95:5]

^aIsolated yields. ^bThe diastereomeric ratio [dr] was determined by ¹H or/and ¹⁹F NMR spectroscopy.


which a variety of different R substituents were evaluated, excellent diastereoselectivities of >92:8 and moderate-to-good yields were obtained. It is noteworthy that while electron-donating and -withdrawing substitution groups at the aryl does not influence diastereoselectivity, with a strong electron-withdrawing group, reactions indeed took a much longer time (see entry 3). Lastly, although predictable, these asymmetric fluorinations can indeed be highly regioselective in favor of the electron-rich enamide-olefin as demonstrated with fluorinations of chiral enamides **7** and **9** (Scheme 2).

While synthetically this method represents a stereoselective approach for constructing chiral α -fluoro-imides, mechanistically, this fluorination provides some insight into the chemistry of chiral enamides. On the basis of the observed stereochemical outcome, a proposed mechanistic model is shown in Scheme 3. We had calculated that the most stable conformation of these chiral enamides has the chiral oxazolidinone ring essentially coplanar with the alkene to maximize the delocalization of the nitrogen lone pair into the olefin.²⁴ With this conformational preference, the chiral oxazolidinone auxiliary plays a distinct role

Scheme 2. A Regioselective Fluorination



Scheme 3. A Proposed Stereochemical Model




in providing a key facial bias for the fluorine to approach from the *Si*-face as represented in *ent*-3d. Subsequent trapping of the *N*-acyl iminium ion **A** with water can in principle be stereoselective, but this stereochemical information is lost in the ensuing oxidation.

On the other hand, for *Z*-enamides such as **3e**, the predicted selectivity would be poor based on this conformational model. To alleviate the allylic strain shown in **3e**, the chiral oxazolidinone ring needs to rotate away and can no longer be coplanar with the alkene, thereby leading much less differentiated π -faces [see **C**]. To support this model, we prepared *Z*-enamide **3e** and found that not only is the yield of its fluorination inferior to those of *E*-enamides but also the diastereoselectivity is diminished significantly.

Lastly, as a useful synthetic application, we examined asymmetric fluorinations of trisubstituted enamides. As shown in Scheme 4, optically enriched 2-fluorocyclohexanone **13**²⁵

Scheme 4. Fluorinations of Trisubstituted Enamides



could be obtained directly from trisubstituted enamide **11** after a simple workup with the fluorinated *N,O*-hemiacetal intermediate **12** being insufficiently stable for isolation. Likewise, the use of enamide **14** led to α -fluorinated aryl ketone **15** in 78% yield with an *er* of $\geq 95:5$. This is in direct contrast with our earlier attempts to isolate α -fluoro-aldehydes via hydrolysis of the corresponding

N,O-hemiacetal intermediate from fluorinations of disubstituted enamides.

We have reported an asymmetric synthesis of α -fluoro-imides via fluorinations of chiral enamides. The reaction involves selectively fluorinating the electron-rich enamide olefin using Selectfluor and *N*-fluoro-benzenesulfonimide [NFSI] followed by trapping of a β -fluoro-iminium cationic intermediate with water and oxidation of the resulting *N,O*-hemiacetal. From a practical perspective, the reaction is operationally simple, requires inexpensive reagents and mild conditions, and provides chiral α -fluoro-imides and optically enriched α -fluoro-ketones in good yields and high selectivities.

■ ASSOCIATED CONTENT

S Supporting Information

Experimental procedures as well as NMR spectra, characterizations, and X-ray structural files (CIF). This material is available free of charge via the Internet at <http://pubs.acs.org>.

■ AUTHOR INFORMATION

Corresponding Authors

*E-mail: yutang@tju.edu.cn.

*E-mail: rhsung@wisc.edu.

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

R.P.H. thanks the NIH (GM66055) for funding. Y.T. thanks the National Natural Science Foundation of China (Nos. 21172169 and 21172168) and The National Basic Research Project (No. 2014CB932201) for generous funding. We also thank Dr. Hai-Bin Song of the Nankai University for providing X-ray structure and data analysis.

■ REFERENCES

- (1) Liang, T.; Neumann, C. N.; Ritter, T. *Angew. Chem., Int. Ed.* **2013**, *52*, 8214. (b) Böhm, H. J.; Banner, D.; Bendels, S.; Kansy, M.; Kuhn, B.; Müller, K.; Obst-Sander, U.; Stahl, M. *ChemBioChem* **2004**, *5*, 637. (c) Wang, J.; Sánchez-Roselló, M.; Pozo, A. D.; Sorochinsky, A. E.; Fustero, S.; Soloshonok, V. A.; Liu, H. *Chem. Rev.* **2014**, *114*, 2432. (d) Purser, S.; Moore, P. R.; Swallow, S.; Gouverneur, V. *Chem. Soc. Rev.* **2008**, *37*, 320. (e) Isanbor, C.; O'Hagan, D. *J. Fluorine Chem.* **2006**, *127*, 303. (f) Hagmann, W. K. *J. Med. Chem.* **2008**, *51*, 4359. (g) Kirk, K. L. *J. Fluorine Chem.* **2006**, *127*, 1013.
- (2) (a) Babudri, F.; Farinola, G. M.; Naso, F.; Ragni, R. *Chem. Commun.* **2007**, 1003. (b) Cametti, M.; Crousse, B.; Metrangolo, P.; Milanid, R.; Resnati, G. *Chem. Soc. Rev.* **2012**, *41*, 31. (c) Kassis, C. M.; Steehler, J. K.; Betts, D. E.; Guan, Z.; Romack, T. J.; DeSimone, J. M.; Linton, R. W. *Macromolecules* **1996**, *29*, 3247. (d) Li, Y. *Acc. Chem. Res.* **2012**, *45*, 723. (e) Kirsch, P.; Hahn, A. *Eur. J. Org. Chem.* **2005**, 3095.
- (3) (a) Jeschke, P. *ChemBioChem* **2004**, *5*, 570. (b) Maienfisch, P.; Hall, R. G. *Chimia* **2004**, *58*, 93.
- (4) Kirsch, P. *Modern Fluoroorganic Chemistry*; Wiley-VCH: Weinheim, Germany, 2004. (b) Davis, F. A.; Yi, H.; Sundarababu, G. In *Enantiocontrolled Synthesis of Fluoro-Organic Compounds*; Soloshonok, V. A., Ed.; Wiley: Chichester, U.K., 1999; p 1. (c) Taylor, S. D.; Kotoris, C. C.; Hum, G. *Tetrahedron* **1999**, *55*, 12431. (d) Smart, B. E. *J. Fluorine Chem.* **2001**, *109*, 3. (e) Schlosser, M. *Tetrahedron* **1978**, *34*, 3.
- (5) Differding, E.; Lang, R. W. *Tetrahedron Lett.* **1988**, *29*, 6087.
- (6) For reviews, see: (a) Ma, J.; Cahard, D. *Chem. Rev.* **2004**, *104*, 6119. (b) Rozen, S. *Acc. Chem. Res.* **2005**, *38*, 803. For recent development of catalytic enantioselective fluorination, see: (c) Kwiatkowski, P.; Beeson, T. D.; Conrad, J. C.; MacMillan, D. W. C. *J. Am. Chem. Soc.* **2011**, *133*, 1738. (d) Beeson, T. D.; MacMillan, D. W. C. *J. Am. Chem. Soc.* **2005**,

- 127, 8826. (e) Enders, D.; Huttel, M. R. M. *Synlett* **2005**, 6, 991. (f) Wang, F.; Li, J.; Hu, Q.; Yang, X.; Wu, X.-Y.; He, H. *Eur. J. Org. Chem.* **2014**, 3607. (g) Hamashima, Y.; Suzuki, T.; Takano, H.; Shimura, Y.; Sodeoka, M. *J. Am. Chem. Soc.* **2005**, 127, 10164. (h) Honjo, T.; Phipps, R. J.; Rauniyar, V.; Toste, F. D. *Angew. Chem., Int. Ed.* **2012**, 51, 9684. (i) Rauniyar, V.; Lackner, A. D.; Hamilton, G. L.; Toste, F. D. *Science* **2011**, 334, 1681. (j) Yang, X.; Phipps, R. J.; Toste, F. D. *J. Am. Chem. Soc.* **2014**, 136, 5225. (k) Paull, D. H.; Scerba, M. T.; Alden-Danforth, E.; Widger, L. R.; Lectka, T. *J. Am. Chem. Soc.* **2008**, 130, 17260. (l) Steiner, D. D.; Mase, N.; Barbas, C. F. *Angew. Chem., Int. Ed.* **2005**, 117, 3772. (m) Marigo, M.; Fielenbach, D.; Braunton, A.; Kjærsgaard, A.; Jørgensen, K. A. *Angew. Chem., Int. Ed.* **2005**, 117, 3769.
- (7) (a) Davis, F. A.; Han, W. *Tetrahedron Lett.* **1992**, 33, 1153. (b) Davis, F. A.; Kasu, P. V. N. *Tetrahedron Lett.* **1998**, 39, 6135.
- (8) (a) Brunet, V. A.; O'Hagan, D.; Slawin, A. M. Z. *J. Fluorine Chem.* **2007**, 128, 1271. (b) Lubin, H.; Dupuis, C.; Pytkowicz, J.; Brigaud, T. *J. Org. Chem.* **2013**, 78, 3487. (c) Nakayama, K.; Browne, D. L.; Baxendale, I. R.; Ley, S. V. *Synlett* **2013**, 24, 1298. (d) Alvarado, J.; Herrmann, A. T.; Zakarian, A. *J. Org. Chem.* **2014**, 79, 6206.
- (9) Phipps, R. J.; Hiramatsu, K.; Toste, F. D. *J. Am. Chem. Soc.* **2012**, 134, 8376.
- (10) For a recent review on chemistry of enamides, see: (a) Carbery, D. R. *Org. Biomol. Chem.* **2008**, 9, 3455. Also see: (b) Rappoport, Z. *The Chemistry of Enamines in The Chemistry of Functional Group*; John Wiley and Sons: New York, 1994.
- (11) For a review on the synthesis of enamides, see: (a) Tracey, M. R.; Hsung, R. P.; Antoline, J.; Kurtz, K. C. M.; Shen, L.; Slafer, B. W.; Zhang, Y. In *Science of Synthesis, Houben-Weyl Methods of Molecular Transformations*; Weinreb, S. M., Ed.; Georg Thieme Verlag KG: Chapter 21.4, 2005. For leading examples of enamide syntheses, see: (b) Bolshan, Y.; Batey, R. A. *Tetrahedron* **2010**, 66, 5283. (c) Zhang, X.; Zhang, Y.; Huang, J.; Hsung, R. P.; Kurtz, K. C. M.; Oppenheimer, J.; Petersen, M. E.; Sagamanova, I. K.; Tracey, M. R. *J. Org. Chem.* **2006**, 71, 4170. (d) Chechik-Lankin, H.; Livshin, S.; Marek, I. *Synlett* **2005**, 2098. (e) Han, C.; Shen, R.; Su, S.; Porco, J. A., Jr. *Org. Lett.* **2004**, 6, 27. (f) Pan, X.; Cai, Q.; Ma, D. *Org. Lett.* **2004**, 6, 1809. (g) Langner, M.; Bolm, C. *Angew. Chem., Int. Ed.* **2004**, 43, 5984. (h) Jiang, L.; Job, G. E.; Klapars, A.; Buchwald, S. L. *Org. Lett.* **2003**, 5, 3667.
- (12) For Diels–Alder cycloadditions of chiral enamides, see: (a) Crawly, S. L.; Funk, R. L. *Org. Lett.* **2006**, 8, 3995. (b) Gohier, F.; Bouhadjera, K.; Faye, D.; Gaulon, C.; Maisonneuve, V.; Dujardin, G.; Dhal, R. *Org. Lett.* **2007**, 9, 211. (c) Fang, L.-C.; Hsung, R. P.; Ma, Z.-X.; Presser, W. R. *Org. Lett.* **2013**, 15, 4842. (d) Fang, L.-C.; Hsung, R. P. *Org. Lett.* **2014**, 16, 1826.
- (13) For [2 + 2] cycloadditions of chiral enamides, see: (a) Ma, Z.-X.; Feltenberger, J. B.; Hsung, R. P. *Org. Lett.* **2012**, 14, 2742. (b) Feltenberger, J. B.; Hayashi, R.; Tang, Y.; Babiash, E. S. C.; Hsung, R. P. *Org. Lett.* **2009**, 11, 3666. (c) Riches, A. G.; Wernersbach, L. A.; Hegedus, L. S. *J. Org. Chem.* **1998**, 63, 4691. (d) Hegedus, L. S.; Bates, R. W.; Soderberg, B. C. *J. Am. Chem. Soc.* **1991**, 113, 923.
- (14) For cyclopropanations, see: (a) Lu, T.; Song, Z. L.; Hsung, R. P. *Org. Lett.* **2008**, 10, 541. (b) Song, Z. L.; Lu, T.; Hsung, R. P.; Al-Rashid, Z. F.; Ko, C. H.; Yu, T. *Angew. Chem., Int. Ed.* **2007**, 46, 4096. (c) Voigt, J.; Noltemeyer, M.; Reiser, O. *Synlett* **1997**, 202.
- (15) For epoxidations, see: (a) Xiong, H.; Hsung, R. P.; Shen, L.; Hahn, J. M. *Tetrahedron Lett.* **2002**, 43, 4449. (b) Adam, W.; Bosio, S. G.; Wolff, B. T. *Org. Lett.* **2003**, 5, 819. (c) Sivaguru, J.; Saito, H.; Poon, T.; Omonuwa, T.; Franz, R.; Jockusch, S.; Hooper, C.; Inoue, Y.; Adam, W.; Turro, N. *J. Org. Lett.* **2005**, 7, 2089. (d) Koseki, Y.; Kusano, S.; Ichi, D.; Yoshida, K.; Nagasaka, T. *Tetrahedron* **2000**, 56, 8855.
- (16) Ko, C. H.; Hsung, R. P.; Al-Rashid, Z. F.; Feltenberger, J. B.; Lu, T.; Yang, J.-H.; Wei, Y.-G.; Zifcsak, C. A. *Org. Lett.* **2007**, 9, 4459.
- (17) Banks, R. E.; Mohialdin-Khaffaf, S. N.; Lal, G. S.; Sharifa, L.; Syvret, R. G. *J. Chem. Soc., Chem. Commun.* **1992**, 595.
- (18) (a) Umamoto, T.; Kawada, K.; Tomita, K. *Tetrahedron Lett.* **1986**, 27, 4465. (b) Umamoto, T.; Fukami, S.; Tomizawa, G.; Harasawa, K.; Kawada, K.; Tomita, K. *J. Am. Chem. Soc.* **1990**, 112, 8563. (c) Umamoto, T.; Nagayoshi, M.; Adachi, K.; Tomizawa, G. *J. Org. Chem.* **1998**, 63, 3379.
- (19) Differding, E.; Ofner, H. *Synlett* **1991**, 187.
- (20) (a) Barton, D. H. R.; Ganguly, A. K.; Hesse, R. H.; Loo, S. N.; Pechet, M. M. *Chem. Commun.* **1968**, 806. (b) Barton, D. H. R.; Westcott, N. D.; Toh, H. T.; Tarzia, G.; Hesse, R. H.; Pechet, M. M. *J. Chem. Soc., Chem. Commun.* **1972**, 122. (c) Airey, J.; Barton, D. H. R.; Ganguly, A. K.; Hesse, R. H.; Pechet, M. M. *An. Quim.* **1974**, 70, 871.
- (21) See Supporting Information.
- (22) Evans, D. A. *Aldrichimica Acta* **1982**, 15, 23.
- (23) Sibi, M. P.; Porter, N. A. *Acc. Chem. Res.* **1999**, 32, 163.
- (24) (a) Lu, T.; Song, Z.; Hsung, R. P. *Org. Lett.* **2008**, 10, 541. (b) Song, Z.; Lu, T.; Hsung, R. P.; Al-Rashid, Z. F.; Ko, C.; Tang, Y. *Angew. Chem., Int. Ed.* **2007**, 46, 4069.
- (25) The absolute stereochemistry of 2-fluorocyclohexanone **13** was confirmed through comparison of its optical rotation ($[\alpha]_{\text{D}}^{23} = +45.7$ [c 0.73, CHCl₃]) with reported values (a) $[\alpha]_{\text{D}}^{23} = +54.8$ [c 1.0, C₆H₆]; Enders, D.; Faure, S.; Potthoff, M.; Runsink, J. *Synthesis* **2001**, 2307. (b) $[\alpha]_{\text{D}}^{23} = +49.6$ [c 0.9, C₆H₆]; Kwiatkowski, P.; Beeson, T. D.; Conrad, J. C.; MacMillan, D. W. C. *J. Am. Chem. Soc.* **2011**, 133, 1738. For ketone **15**, it is assigned based on a mechanistic analogy with only the racemic form having been reported and the fact that the assignment of **13** is in good agreement with our stereochemical assessment for all other α -fluorinated-imides.