Synthesis of 1,1'-Binaphthyl-2,2'-diyl Phosphoroselenoic Amides and Their Conversion to Optically Pure Phosphoramidites

Toshiaki Murai,*1 Shinsuke Inaji,1 Ken Morishita,1 Fumitoshi Shibahara,1 Makoto Tokunaga,2

Yasushi Obora,2 and Yasushi Tsuji2

¹Department of Chemistry, Faculty of Engineering, Gifu University, Yanagido, Gifu 501-1193

²Catalysis Research Center and Division of Chemistry, Graduate School of Science, Hokkaido University, Sapporo 001-0021

(Received September 27, 2006; CL-061125; E-mail: mtoshi@gifu-u.ac.jp)

Optically pure phosphoroselenoyl chloride was reacted with racemic amines to give phosphoroselenoic amides as two diastereomeric mixtures in high yields. The diastereomeric mixtures were separated by fractional recrystallization or by chromatography. Extrusion of the selenium atom from the separated amides led to diastereomerically pure phosphoramidites. The ability of the obtained amidites to act as optically active ligands was tested in the hydrogenation of an imine.

Phosphoramidites bearing a 1,1'-bi-2-naphthoyl group have been recognized as some of the most powerful ligands in catalytic asymmetric reactions¹ since Feringa and co-workers reported the first one bearing a N,N-dimethylamino group as a chiral discriminating agent in 1994.² Their synthesis generally involves the reaction of chlorophosphites with amines.³ Alternatively, the amino group substitution reaction of N,N-dimethylamino phosphoramidites with other amines is used.⁴ However, only a few phosphoramidites possessing a chiral center at the amino group⁵ have been reported except for those with chiral 1-arylalkylamino groups.⁶

We recently found that phosphoroselenoyl chloride **1** can be obtained from PCl₃, elemental selenium and 1,1'-bi-2-naphthol (Scheme 1).⁷ The chloride **1** is stable under neutral conditions but reactive under basic conditions. We report here the reaction of **1** with primary and secondary amines leading to phosphoroselenoic amides, separation of their diastereomers, and conversion of the separated diastereomers to optically pure phosphoramidites.



Scheme 1.

The reaction of chloride **1** with racemic 1-phenylethylamine (**2a**) proceeded smoothly under reflux in toluene (Scheme 2).⁸ Two diastereomers of phosphoroselenoic amide (R_{ax} ,S)-**3a** and (R_{ax} ,R)-**3a**^{r9,10} were obtained in a ratio of 39:61.^{11,15} These diastereomers were separated by fractional crystallization to give



Scheme 2	2
----------	---

Table 1. Synthesis of phosphoroselenoic amides 3 by reacting 1 with amines 2^a

п1

(<i>R</i> _{ax})-1	$\begin{array}{c} R \\ N \\ H \\ 2 \\ \hline With or without \\ DMAP \\ Toluene, reflux \end{array} $		+ (0) 	\mathbb{R}^{1} \mathbb{R}^{2} \mathbb{R}^{2}
Entry	Amine 2	Time/h	Ŋ	rield/% ^b
1	H₂N 1-naphthyl	2	(R_{ax},S) -3b (R_{ax},R) -3b'	35 ^d 57
2	H ₂ N 2c	2	(R _{ax} ,S)- 3c (R _{ax} ,R)- 3c '	46 ^e 49
3	H ₂ N 2d	2	$(R_{\rm ax},S)$ -3d $(R_{\rm ax},R)$ -3d'	44 ^e 48
4	HN 2e	2	(R _{ax} ,S)- 3e (R _{ax} ,R)- 3e'	36 ^e 58
5	2f Ph N Ph	44	3f 3f' 3f + 3f'	34 ^{f,g} 34 25
6 ^c	Ph_N_Ph	18	(R_{ax},S) -3g (R_{ax},R) -3g' 3g + 3g'	30 ^f 22 36
7	Ph N H OEt	12	3h 3h' 3h + 3h'	46 ^{f,g} 36 5
8	Ph N Ph	13	3i 3i' 3i + 3i'	41 ^{f,g} 40 14
9 ^c		2	(R _{ax} ,S)-3j (R _{ax} ,R)-3j' 3j + 3j'	14 ^f 7 54
10 ^c		2.5	(R_{ax},S) -3k (R_{ax},R) -3k' 3k + 3k'	46 ^f 29 14

^aThe chloride **1** was reacted with amine **2** (2 equiv.) under reflux in toluene unless otherwise noted. ^bIsolated yield. ^cAmine **2** (1.2 equiv.) and DMAP (1.2 equiv.) was used. ^dDiastereomers were separated by fractional recrystallization. ^eDiastereomers were separated by HPLC on silica gel. ^fDiastereomers were separated by column chromatography on silica gel. ^gThe absolute configuration of diastereomers is not determined.



diastereomerically pure **3a** and **3a'** in respective yields of 36 and 60%.

A wide variety of primary and secondary amines are subjected to the amination reaction of 1. The results are shown in Table 1. The reaction was carried out with either amines 2 (2 equiv.) (Entries 1-5, 7, and 8) or amines 2 (1.2 equiv.) in the presence of 4-dimethylaminopyridine (DMAP, 1.2 equiv.) (Entries 6, 9, and 10). In all cases, the substitution reaction of 1 with amines 2 took place selectively at the phosphorus atom to give phosphoroselenoic amides 3 as a diastereomeric mixture in high yields. The reaction rate depends on the substitution pattern around the nitrogen atom of 2. The reaction with primary amines 2b-2d was complete within 2h (Entries 1-3), whereas the reaction with acyclic secondary amines 2f-2i required a longer reaction time (Entries 5-8). The diastereomers 3b and **3b'** were separated by fractional recrystallization. Other diastereomers 3 were separated either by high performance recycle liquid chromatography on silica gel or by column chromatography on silica gel. The absolute configuration of the diastereomerically pure products 3 was determined by comparison to the ³¹PNMR spectra of authentic samples derived from optically active amines. Alternatively, stereochemistry was deduced based on X-ray molecular structure analyses of pure 3.12,15

Extrusion of the selenium atom of **3** was then performed (Scheme 3). The reaction of **3** with Bu_3P went to completion within 2 h in toluene at room temperature to give the corresponding phosphoramidites **4** along with $Bu_3P=Se$. The complete conversion of **3** to **4** was clearly confirmed by ³¹P NMR spectra, but the phosphoramidites **4** (R = H) derived from primary amines were not isolated, although their synthesis has already been reported.¹³ Phosphoramidites **4** (R = H) are sensitive toward air to give oxidized products along with several unidentified products. In contrast, phosphoramidites **4** derived from secondary amines were isolated in good yields, as exemplified by **4f'**, **4g**, and **4k**.

Finally, the ability of phosphoramidites **4** to act as optically active ligands was tested in the Ir-catalyzed hydrogenation reaction of imine 5^{14} (Scheme 4). A catalytic amount of [Ir(cod)₂]-BF₄ efficiently catalyzed the reaction, and the enantiomeric excess was influenced by the substitution pattern in the phosphoramidites **4**. The hydrogenation reaction of **5** in the presence of **4g** gave amine **6** in racemic form. The use of **4f**' effected the asymmetric hydrogenation of **5** to give (*R*)-**6** with 53% ee, whereas the reaction in the presence of **4k** led to (*S*)-**6** with 73% ee.

In summary, we have demonstrated the highly efficient synthesis of optically active phosphoroselenoic amides and their purification in diastereomerically pure form. Extrusion of the selenium atom of phosphoroselenoic amides with Bu₃P provides



new types of phosphoramidites, which are not readily accessible by known synthetic procedures, but which may be important as optically active ligands.

This work was supported in part by a Grant-in-Aid for Scientific Research on Priority Area (No. 18037024, "Advanced Molecular Transformations of Carbon Resources") from the Ministry of Education, Culture, Sports, Science and Technology, Japan. AIR WATER INC. kindly provided the optically active 1,1'-bi-2-naphthol.

References and Notes

- For recent examples: a) A. Zhang, T. V. RajanBabu, *J. Am. Chem. Soc.* 2006, *128*, 5620. b) R. B. C. Jagt, P. Y. Toullec, D. Geerdink, J. G. De Vries, B. L. Ferringa, A. J. Minnaard, *Angew. Chem., Int. Ed.* 2006, *45*, 2789. c) M. T. Reetz, G. Mehler, O. Bondarev, *Chem. Commun.* 2006, 2292.
- 2 R. Hulst, N. K. de Vries, B. L. Feringa, *Tetrahedron: Asymmetry* 1994, 5, 699.
- 3 a) A. H. M. de Vries, A. Meetsma, B. L. Feringa, Angew. Chem., Int. Ed. Engl. 1996, 35, 2374. b) L. A. Arnold, R. Imbos, A. Mandoli, A. H. M. de Vries, R. Naasz, B. L. Feringa, Tetrahedron 2000, 56, 2865.
- 4 D. Pena, A. J. Minnaard, J. G. de Vries, B. L. Feringa, J. Am. Chem. Soc. 2002, 124, 14552.
- G. Francio, F. Faraone, W. Leitner, J. Am. Chem. Soc. 2002, 124, 736. b) R. Kumareswaran, M. Nandi, T. V. RajanBabu, Org. Lett. 2003, 5, 4345. c) T. Watanabe, T. F. Knoepfel, E. M. Carreira, Org. Lett. 2003, 5, 4557. d) E. Cesarotti, S. Araneo, I. Rimoldi, S. Tassi, J. Mol. Catal. A: Chem. 2003, 204–205, 211. e) A. Duursma, J.-G. Boiteau, L. Lefort, J. A. F. Boogers, A. H. M. De Varies, J. G. De Vries, A. J. Minnaard, B. L. Feringa, J. Org. Chem. 2004, 69, 8045.
- For recent examples: a) Q.-H. Zeng, X.-P. Hu, Z.-C. Duan, X.-M. Liang, Z. Zheng, J. Org. Chem. 2006, 71, 393. b) E. P. Kündig, P. D. Chaudhuri, D. House, G. Bernardinelli, Angew. Chem., Int. Ed. 2006, 45, 1092. c) G. P. Howell, A. J. Minnaard, B. L. Feringa, Org. Biomol. Chem. 2006, 4, 1278. d) D. Polet, A. Alexakis, K. Tissot-Croset, C. Corminboeuf, K. Ditrich, Chem. Eur. J. 2006, 12, 3596.
- 7 T. Murai, D. Matsuoka, K. Morishita, J. Am. Chem. Soc. 2006, 128, 4584.
- 8 A precedent for the reaction of achiral phosphoroselenoyl chlorides with allylamine has been reported: Y. I. Mel'nik, D. I. Prots, Y. I. Kolodii, Ukr. Khim. Zh. 1987, 53, 638.
- 9 The oxygen isologue of **3a** was prepared at room temperature for 36 h in a manner similar to that in Scheme 1: B.-Q. Gong, W.-Y. Chen, B.-F. Hu, *J. Org. Chem.* **1991**, *56*, 423.
- 10 The sulfur isologue of **3a** was prepared by reacting phosphoramidothioic dichloride with 1,1'-bi-2-naphthol: D. Fabbri, G. Delogu, O. De Lucchi, *J. Org. Chem.* **1993**, *58*, 1748.
- 11 For spectroscopic properties of the new compounds, see the Supporting Information.
- 12 For details of X-ray molecular structure analyses, see the Supporting Information.
- 13 B. Bartels, C. Garcia-Yebra, G. Helmchen, *Eur. J. Org. Chem.* 2003, 1097.
- 14 For recent examples of the asymmetric catalytic hydrogenation of imines: a) C. Moessner, C. Bolm, *Angew. Chem., Int. Ed.* 2005, 44, 7564. b) T. Imamoto, N. Iwadate, K. Yoshida, *Org. Lett.* 2006, 8, 2289.
- 15 Supporting Information is available electronically on the CSJ Journal WEB site: http://www.csj.jp/journals/chem-lett/index.html.