This article was downloaded by: [University of Kiel] On: 22 October 2014, At: 18:33 Publisher: Taylor & Francis Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



# Synthetic Communications: An International Journal for Rapid Communication of Synthetic Organic Chemistry

Publication details, including instructions for authors and subscription information: http://www.tandfonline.com/loi/lsyc20

# RuCl<sub>2</sub>(PPh<sub>3</sub>)<sub>3</sub> Catalyzed Isomerization of the Baylis-Hillman Adducts

Deevi Basavaiah <sup>a</sup> & Kannan Muthukumaran <sup>a</sup> <sup>a</sup> School of Chemistry, University of Hyderabad , Hyderabad, 500 046, India Published online: 17 Sep 2007.

To cite this article: Deevi Basavaiah & Kannan Muthukumaran (1999) RuCl<sub>2</sub>(PPh<sub>3</sub>)<sub>3</sub> Catalyzed Isomerization of the Baylis-Hillman Adducts, Synthetic Communications: An International Journal for Rapid Communication of Synthetic Organic Chemistry, 29:4, 713-719, DOI: <u>10.1080/00397919908085821</u>

To link to this article: http://dx.doi.org/10.1080/00397919908085821

## PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages,

and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at http://www.tandfonline.com/page/terms-and-conditions

### RuCl<sub>2</sub>(PPh<sub>3</sub>)<sub>3</sub> CATALYZED ISOMERIZATION OF THE BAYLIS-HILLMAN ADDUCTS

Deevi Basavaiah\* and Kannan Muthukumaran

School of Chemistry, University of Hyderabad, Hyderabad-500 046, India

Abstract: RuCl<sub>2</sub>(PPh<sub>3</sub>)<sub>3</sub> catalyzed isomerization of the Baylis-Hillman adducts *i.e.* methyl 3-aryl-3-hydroxy-2-methylenepropanoates to methyl 3-aryl-2-methyl-3-oxopropanoates has been described.

The transition metal catalyzed isomerization of allyl alcohols to saturated carbonyl molecules is an important and useful operation in synthetic organic chemistry and is of current interest.<sup>1-9</sup> In continuation of our interest in the Baylis-Hillman reaction,<sup>10-13</sup> we herein report the ruthenium catalyzed isomerization of the Baylis-Hillman adducts *i.e.* methyl 3-aryl-3-hydroxy-2-methylenepropanoates leading to the formation of methyl 3-aryl-2-methyl-3-oxopropanoates, thus describing a new aspect of Baylis-Hillman chemistry.

The Baylis-Hillman reaction provides functionalized allyl alcohols (eq. 1) which have been successfully employed in various useful organic transformations.<sup>10-19</sup> However, these allyl alcohols have not been employed as substrates for transition metal catalyzed isomerization to obtain the corresponding saturated carbonyl

Copyright © 1999 by Marcel Dekker, Inc.

<sup>\*</sup> To whom correspondence should be addressed

molecules. We have therefore examined the possible isomerization of these allyl alcohols *i.e.* methyl 3-hydroxy-2-methylenealkanoates (1) derived from methyl acrylate under the catalytic influence of RuCl<sub>2</sub>(PPh<sub>3</sub>)<sub>3</sub> with a view to expand the scope of the application of the Baylis-Hillman adducts in organic synthesis. We have first selected methyl 3-hydroxy-2-methylene-3-phenylpropanoate (1a) as a substrate for the isomerization reaction. The best result was obtained when this molecule *i.e.* methyl 3-hydroxy-2-methylene-3-phenylpropanoate (1a) (1 mM) was treated with a catalytic amount of RuCl<sub>2</sub>(PPh<sub>3</sub>)<sub>3</sub> (4 mol%, 0.04 mM) and K<sub>2</sub>CO<sub>3</sub> (1 mM) in toluene (2 mL) at reflux temperature for 12 hours to provide the desired methyl 2-methyl-3-oxo-3-phenylpropanoate (2a) in 53% yield.



Encouraged by this result, we have subjected a representative class of Baylis-Hillman adducts (1a-f) to  $RuCl_2(PPh_3)_3$  catalyzed isomerization (eq. 2) (Table) which provided the desired methyl 3-aryl-2-methyl-3-oxopropanoates (2a-f) in 42-61% yields.<sup>20</sup>



Substrate	R	Product	Yield <sup>c</sup>
1a	C <sub>6</sub> H <sub>5</sub>	2a	53
1b	4-MeC <sub>6</sub> H₄	2b	51
1c	4-EtC <sub>6</sub> H₄	2c	42
1d	4-i-PrC <sub>6</sub> H₄	2d	57
1e	4-MeOC <sub>6</sub> H <sub>4</sub>	2e	61
1f	2-MeOC <sub>6</sub> H₄	2f	50

Table : Isomerization of the Baylis-Hillman Adducts<sup>a, b</sup>

a) All reactions were carried out in 1 mM scale in toluene at reflux temperature for 12 hours.

b) All products were obtained as colorless liquids and gave satisfactory IR, <sup>1</sup>H NMR, <sup>13</sup>C NMR, mass spectral and elemental analyses.

c) Yields of the products after column chromatography (silica gel, 5% EtOAc in hexane).

In conclusion, we have successfully carried out the RuCl<sub>2</sub>(PPh<sub>3</sub>)<sub>3</sub> catalyzed isomerization of methyl 3-aryl-3-hydroxy-2-methylenepropanoates to methyl 3-aryl-2-methyl-3-oxopropanoates. Though these kind of  $\beta$ -keto esters can be obtained in better yields using other conventional methods, the present methodology represents a new direction for Baylis-Hillman chemistry.

#### **EXPERIMENTAL**

IR spectra were recorded on Jasco-FT-IR model 5300 spectrometer using samples as neat liquids. <sup>1</sup>H NMR (200 MHz) and <sup>13</sup>C NMR (50 MHz) spectra were recorded on Bruker-AC-200 spectrometer using CDCl<sub>3</sub> as solvent and

tetramethylsilane (TMS,  $\delta = 0$  ppm) as internal standard. Elemental analyses were performed on Perkin-Elmer 240C-CHN analyser. Mass spectra were recorded on Hewlett Packard (model HP 5989A) or Finnigon MAT (1020C) mass spectrometer.

## General procedure for the isomerization of methyl 3-aryl-3-hydroxy-2methylenepropanoates:

To a solution of the allyl alcohol (1) (1 mM) in toluene (2 mL), was added RuCl<sub>2</sub>(PPh<sub>3</sub>)<sub>3</sub> (4 mol%, 0.04 mM, 38 mg) followed by K<sub>2</sub>CO<sub>3</sub> (1 mM, 138 mg) at room temperature and this suspension was refluxed for 12 hours. Solvent was evaporated under vacuum. The residue obtained was purified by column chromatography (silica gel, 5% EtOAc in hexane) under N<sub>2</sub> pressure to obtain the pure  $\beta$ -keto ester (2).

#### Methyl 2-methyl-3-oxo-3-phenylpropanoate (2a):

Yield: 53%, IR (neat)/cm<sup>-1</sup> : 1687, 1743; <sup>1</sup>H NMR :  $\delta$  1.50 (d, 3H, J= 6.8 Hz), 3.68 (s, 3H), 4.40 (q, 1H, J= 6.8 Hz), 7.40-7.67 (m, 3H), 7.98 (d, 2H, J= 7.2 Hz); <sup>13</sup>C NMR :  $\delta$  13.81, 48.09, 52.38, 128.59, 128.76, 133.47, 135.88, 171.26, 195.79; MS (m/z) : 192 (M<sup>+</sup>); Analysis calculated for C<sub>11</sub>H<sub>12</sub>O<sub>3</sub>: C, 68.73; H, 6.29; found: C, 68.80; H, 6.32.

#### Methyl 2-methyl-3-(4-methylphenyl)-3-oxopropanoate (2b):

Yield: 51%; IR (neat)/cm<sup>-1</sup>: 1682, 1743; <sup>1</sup>H NMR :  $\delta$  1.49 (d, 3H, J= 7.0 Hz), 2.42 (s, 3H), 3.68 (s, 3H), 4.38 (q, 1H, J= 7.0 Hz), 7.27 (d, 2H, J= 8.2 Hz), 7.88 (d, 2H, J= 8.2 Hz); <sup>13</sup>C NMR :  $\delta$  13.69, 21.38, 47.76, 52.11, 128.58, 129.30,

133.25, 144.23, 171.19, 195.23; MS (m/z): 206 (M<sup>+</sup>); Analysis calculated for  $C_{12}H_{14}O_3$ : C, 69.88; H, 6.84; found: C, 70.11; H, 6.83.

#### Methyl 3-(4-ethylphenyl)-2-methyl-3-oxopropanoate (2c):

Yield: 42%; IR (neat)/cm<sup>-1</sup>: 1684, 1745; <sup>1</sup>H NMR :  $\delta$  1.26 (t, 3H, J= 7.6 Hz), 1.48 (d, 3H, J= 6.8 Hz), 2.72 (q, 2H, J= 7.6 Hz), 3.68 (s, 3H), 4.39 (q, 1H, J= 6.8 Hz), 7.30 (d, 2H, J= 8.4 Hz), 7.90 (d, 2H, J= 8.4 Hz); <sup>13</sup>C NMR :  $\delta$  13.86, 15.00, 28.91, 47.97, 52.33, 128.25, 128.84, 133.56, 150.53, 171.38, 195.38; MS (m/z): 220 (M<sup>+</sup>); Analysis calculated for C<sub>13</sub>H<sub>16</sub>O<sub>3</sub>: C, 70.88; H, 7.32; found: C, 70.57; H, 7.35.

#### Methyl 3-(4-isopropylphenyl)-2-methyl-3-oxopropanoate (2d):

Yield: 57%; IR (neat)/cm<sup>-1</sup> : 1684, 1743; <sup>1</sup>H NMR :  $\delta$  1.27 (d, 6H, J= 7.2 Hz), 1.48 (d, 3H, J=7.0 Hz), 2.95 (q, 1H, J= 7.2 Hz), 3.69 (s, 3H), 4.39 (q, 1H, J= 7.0 Hz), 7.32 (d, 2H, J= 8.0 Hz), 7.93 (d, 2H, J= 8.0 Hz); <sup>13</sup>C NMR :  $\delta$  13.80, 23.49, 34.16, 47.84, 52.23, 126.78, 128.83, 133.60, 155.00, 171.30, 195.30; MS (m/z): 234 (M<sup>+</sup>); Analysis calculated for C<sub>14</sub>H<sub>18</sub>O<sub>3</sub>: C, 71.77; H, 7.74; found: C, 71.80; H, 7.76.

#### Methyl 3-(4-methoxyphenyl)-2-methyl-3-oxopropanoate (2e):

Yield: 61%; IR (neat)/cm<sup>-1</sup>: 1678, 1743; <sup>1</sup>H NMR :  $\delta$  1.48 (d, 3H, J=7.0 Hz), 3.68 (s, 3H), 3.87 (s, 3H), 4.35 (q, 1H, J= 7.0 Hz), 6.95 (d, 2H, J= 8.8 Hz), 7.96 (d, 2H, J= 8.8 Hz); <sup>13</sup>C NMR :  $\delta$  13.78, 47.59, 52.16, 55.35, 113.86, 128.66, 130.83, 163.79, 171.33, 194.16; MS (m/z): 222 (M<sup>T</sup>); Analysis calculated for C<sub>12</sub>H<sub>14</sub>O<sub>4</sub>: C, 64.85; H, 6.34; found: C, 64.62; H, 6.32.

#### Methyl 3-(2-methoxyphenyl)-2-methyl-3-oxopropanoate (2f):

Yield: 50%; IR (neat)/cm<sup>-1</sup>: 1678, 1741; <sup>1</sup>H NMR :  $\delta$  1.43 (d, 3H, J=6.8 Hz), 3.66 (s, 3H), 3.87 (s, 3H), 4.33 (q, 1H, J= 6.8 Hz), 7.00 (m, 2H), 7.48 (m, 1H), 7.79 (d,

1H, J= 7.8 Hz); <sup>13</sup>C NMR :  $\delta$  13.41, 51.94, 52.54, 55.26, 111.54, 120.88, 126.69, 131.07, 134.10, 158.47, 171.97, 197.08; MS (m/z): 222 (M<sup>+</sup>); Analysis calculated for C<sub>12</sub>H<sub>14</sub>O<sub>4</sub>: C, 64.85; H, 6.34; found: C, 65.00; H, 6.31.

Acknowledgements: We thank DST (New Delhi) for funding this project. KMK thanks UGC (New Delhi) for his research fellowship. We also thank UGC (New Delhi) for COSIST and Special Assistance Programs in Organic Synthesis in the School of Chemistry, University of Hyderabad.

### **REFERENCES AND NOTES**

- 1. Trost, B. M.; Kulawiec, R. J.; J. Am. Chem. Soc., 1993, 115, 2027.
- 2. Goetz, R. W.; Orchin, M.; J. Am. Chem. Soc., 1963, 85, 1549.
- 3. McGrath, D. V.; Grubbs, R. H.; Organometallics, 1994, 13, 224.
- 4. Damico, R.; Logan, T. J.; J. Org. Chem., 1967, 32, 2356.
- Smadja, W.; Ville, G.; Georgoulis, C.; J. Chem. Soc., Chem. Commun., 1980, 594.
- 6. Li, C.-J.; Wang, D.; Chen, D.-L.; J. Am. Chem. Soc., 1995, 117, 12867.
- 7. Backvall, J.-E.; Andreasson, U.; Tetrahedron Lett., 1993, 34, 5459.
- 8. Sasson, Y.; Rempel, G. L.; Tetrahedron Lett., 1974, 4133.
- 9. Tani, K.; Pure & Appl. Chem., 1985, 57, 1845.
- 10. Basavaiah, D.; Gowriswari, V. V. L.; Tetrahedron Lett., 1986, 27, 2031.
- 11. Basavaiah, D.; Sarma, P. K. S.; J. Chem. Soc., Chem. commun., 1994, 955.
- 12. Basavaiah, D.; Pandiaraju, S.; Padmaja, K.; Synlett, 1996, 393.
- Basavaiah, D.; Krishnamacharyulu, M.; Suguna Hyma, R.; Pandiaraju, S.; Tetrahedron Lett., 1997, 38, 2141.
- 14. Drewes, S. E.; Roos, G. H. P.; Tetrahedron., 1988, 44, 4653.
- Basavaiah, D.; Dharma Rao, P.; Suguna Hyma, R.; *Tetrahedron.*, 1996, 52, 8001 and references cited therein.

- 16. Brzezinski, L. J.; Rafel, S.; Leahy, J. W.; J. Am. Chem. Soc., 1997, 119, 4317.
- 17.Brown, J. M.; Cutting, I.; J. Chem. Soc., Chem. commun., 1985, 578.
- 18. Hoffmann, H. M. R.; Rabe, J.; J. Org. Chem., 1985, 50, 3849.
- Bailey, M.; Marko, I. E.; Ollis, W. D.; Rasmussen, P. R.; *Tetrahedron Lett.*, 1990, 31, 4509.
- We have found that isomeriztation of methyl 3-hydroxy-2-methylenehexanoate (R= n-propyl) was not clean.

(Received in the USA 18 August 1998)