

Refinement

Refinement on F^2	$(\Delta/\sigma)_{\max} < 0.001$
$R[F^2 > 2\sigma(F^2)] = 0.061$	$\Delta\rho_{\max} = 0.29 \text{ e } \text{\AA}^{-3}$
$wR(F^2) = 0.150$	$\Delta\rho_{\min} = -0.16 \text{ e } \text{\AA}^{-3}$
$S = 1.079$	Extinction correction:
8434 reflections	<i>SHELXTL</i> (Sheldrick, 1996)
528 parameters	Extinction coefficient:
All H-atom parameters refined	0.0196 (16)
$w = 1/[\sigma^2(F_o^2) + (0.0382P)^2 + 0.5786P]$	Scattering factors from
where $P = (F_o^2 + 2F_c^2)/3$	<i>International Tables for Crystallography</i> (Vol. C)

Table 1. Selected geometric parameters ($\text{\AA}$, $^\circ$)

O1A—C1A	1.340 (2)	O1B—C1B	1.346 (2)
O2A—C4A	1.335 (2)	O2B—C4B	1.343 (2)
O3A—C13A	1.244 (2)	O3B—C13B	1.242 (2)
O4A—C6A	1.240 (2)	O4B—C6B	1.247 (2)
C9A—C10A	1.316 (3)	C9B—C10B	1.316 (3)
C8A—C15A—C11A	94.33 (15)	C8B—C15B—C11B	93.2 (2)
C12A—C7A—C8A—C9A	66.7 (2)		
C12A—C7A—C8A—C15A	−37.2 (2)		
C15A—C8A—C9A—C10A	32.1 (2)		
C7A—C8A—C9A—C10A	−71.1 (2)		
C8A—C9A—C10A—C11A	0.6 (2)		
C9A—C10A—C11A—C15A	−32.9 (2)		
C9A—C10A—C11A—C12A	70.4 (2)		
C8A—C7A—C12A—C11A	0.7 (2)		
C10A—C11A—C12A—C7A	−67.8 (2)		
C15A—C11A—C12A—C7A	36.0 (2)		
C9A—C8A—C15A—C11A	−48.6 (2)		
C7A—C8A—C15A—C11A	59.0 (2)		
C10A—C11A—C15A—C8A	49.0 (2)		
C12A—C11A—C15A—C8A	−58.6 (2)		
C12B—C7B—C8B—C9B	69.8 (2)		
C12B—C7B—C8B—C15B	−34.7 (2)		
C15B—C8B—C9B—C10B	34.1 (3)		
C7B—C8B—C9B—C10B	−70.2 (2)		
C8B—C9B—C10B—C11B	−0.7 (3)		
C9B—C10B—C11B—C15B	−32.6 (3)		
C9B—C10B—C11B—C12B	71.1 (2)		
C8B—C7B—C12B—C11B	−3.2 (2)		
C10B—C11B—C12B—C7B	−64.6 (2)		
C15B—C11B—C12B—C7B	39.8 (2)		
C9B—C8B—C15B—C11B	−49.9 (2)		
C7B—C8B—C15B—C11B	58.3 (2)		
C10B—C11B—C15B—C8B	49.4 (2)		
C12B—C11B—C15B—C8B	−60.5 (2)		

Table 2. Hydrogen-bonding geometry ($\text{\AA}$, $^\circ$)

D—H...A	D—H	H...A	D...A	D—H...A
O1A—H10A...O3A	0.99 (3)	1.63 (3)	2.526 (2)	148 (2)
O1B—H10B...O3B	1.10 (4)	1.52 (4)	2.530 (2)	149 (4)
O2A—H20A...O4A	0.96 (3)	1.64 (3)	2.514 (3)	148 (2)
O2B—H20B...O4B	0.96 (4)	1.65 (3)	2.529 (3)	150 (3)

The data collection covered a hemisphere of reciprocal space by a combination of three sets of exposures; each set had a different φ angle (0, 88 and 180°) for the crystal and each exposure of 30 s covered 0.3° in ω . The crystal-to-detector distance was 4 cm ($2\theta_{\max} = 66.37^\circ$) and the detector swing angle was -35° . Coverage of the unique set is over 99% complete. Crystal decay was monitored by repeating 30 initial frames at the end of data collection and analysing the duplicate reflections, and was found to be negligible. Only reflections with $2\theta < 62^\circ$ were used for the refinement.

The structure was solved by direct methods and refined by full-matrix least-squares techniques. All H atoms were located from a difference Fourier map and refined isotropically. C—H

distances range from 0.93 (2) to 1.02 (3) $\text{\AA}$ and U_{iso} values for H atoms range from 0.49 (6) to 0.15 (2) $\text{\AA}^2$.

Data collection: *SMART* (Siemens, 1996a). Cell refinement: *SAINT* (Siemens, 1996b). Data reduction: *SAINT*. Program(s) used to solve structure: *SHELXTL* (Sheldrick, 1996). Program(s) used to refine structure: *SHELXTL*. Molecular graphics: *SHELXTL*. Software used to prepare material for publication: *SHELXTL* and *PARST* (Nardelli, 1995).

DM would like to thank DST, New Delhi, for financial assistance. The authors would also like to thank the Malaysian Government for the research grant R&D No. 190-9609-2801. KC thanks the Universiti Sains Malaysia for a Visiting Post-Doctoral Fellowship.

Supplementary data for this paper are available from the IUCr electronic archives (Reference: FR1163). Services for accessing these data are described at the back of the journal.

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
- Cheng, R. K. Y. & Cheng, C. C. (1978). *J. Med. Chem.* **21**, 291–294.
- Majumdar, G., Murty, K. V. S. N. & Mal, D. (1994). *Tetrahedron Lett.* **35**, 6139–6140.
- Nardelli, M. (1995). *J. Appl. Cryst.* **28**, 659.
- Sheldrick, G. M. (1996). *SHELXTL Reference Manual*. Version 5.0. University of Göttingen, Germany.
- Siemens (1996a). *SMART Software Reference Manual*. Siemens Analytical X-ray Instruments Inc., Madison, Wisconsin, USA.
- Siemens (1996b). *SAINT Software Reference Manual*. Version 4. Siemens Analytical X-ray Instruments Inc., Madison, Wisconsin, USA.

Acta Cryst. (1999). **C55**, 778–781

5-Benzyloxy-1-phenyltetrazole: catalytic transfer hydrogenolysis of benzyl ethers

AMADEU F. BRIGAS,^a WILLIAM CLEGG,^b ROBERT A. W. JOHNSTONE^c AND RICARDO F. MENDONCA^a

^aUCEH, Universidade do Algarve, Campus de Gambelas, 8000 Faro, Portugal, ^bDepartment of Chemistry, University of Newcastle upon Tyne, Newcastle upon Tyne NE1 7RU, England, and ^cDepartment of Chemistry, University of Liverpool, Liverpool L69 3BX, England. E-mail: w.clegg@ncl.ac.uk

(Received 4 November 1998; accepted 4 January 1999)



Abstract

In contrast to the earlier reported inertness of 5-benzyloxy-1-phenyltetrazole, (1a), and 3-benzyloxy-1,2-benzisothiazole 1,1-dioxide, (2a), towards catalytic

transfer hydrogenolysis in toluene/water solvent systems, it is now shown that simply changing the solvent to tetrahydrofuran/water leads to their rapid hydrogenolysis with formation of toluene. Compound (1a), C₁₄H₁₂N₄O, has a short C—O ether bond to the tetrazole ring [1.3261(15) Å] and a particularly long C—O bond to the benzyl group [1.4676(16) Å], consistent with an activation of this longer bond by the strongly electronegative tetrazole unit.

Comment

It has been proposed, and demonstrated, that bond lengths are related to chemical reactivity (Bürgi & Dunitz, 1987; Edwards *et al.*, 1986). Structures of aryl, alkyl and allyl ethers, (1b)–(1d) and (2b)–(2d), have



been determined by X-ray crystallography (Alves *et al.*, 1997; Brigas & Johnstone, 1996; Barkley *et al.*, 1997; Brigas *et al.*, 1998, 1997). Generally, it is found that the ether linkage in aryl compounds (1b) and (2b) has a very short C—O bond (*p*, approximately 1.32 Å) to the heteroaromatic group and a very long C—O ether bond (*q*, approximately 1.46 Å) to the aryl group. The combined length of the C—O—C bonds (*p* + *q*) in ethers (1b)–(1d) and (2b)–(2d) is remarkably constant at about 2.78 Å, the two bonds being strongly negatively correlated. On the basis of these results, it was suggested that bond lengths did reflect chemical reactivity in the series of ethers (1) and (2) (Alves *et al.*, 1997).

Considered as 'alkyl' ethers, and by comparison with (1c) and (2c), the benzylic compounds (1a) and (2a) might be expected to be inert to hydrogenolysis (Johnstone *et al.*, 1985; Alves *et al.*, 1997). Alternatively, benzylic compounds are known to add oxidatively to transition metals (Chaloner, 1986). Whereas benzyloxy carbonyl esters are easily hydrogenolysed catalytically by use of hydrogen itself or by transfer methods, benzyl ethers are much more resistant to transfer reduction, typically requiring several hours at elevated tempera-

tures or the additional use of Lewis acids (Johnstone *et al.*, 1985). Therefore the benzylic ethers (1a) and (2a) might be considered to be anomalous whether they react or not, when subjected to hydrogenolysis, especially as benzyl carbamates can be readily hydrogenolysed (Jackson & Johnstone, 1976).

Heterogeneous catalytic transfer hydrogenolyses are commonly carried out in biphasic benzene/water or toluene/water solvent systems. Under such conditions, the ethers (1a) and (2a) remained unchanged over a period of 1 h, but formed some benzyl alcohol by hydrolysis. A similar result has been recorded earlier (Price, 1985). In contrast, the same ethers in a refluxing tetrahydrofuran/water biphasic solvent system with the same Pd/C catalyst and sodium phosphinate as hydrogen donor give quantitative formation of toluene in 10 min at lower temperatures (336 K), with no hydrolysis to benzyl alcohol [reactions (I) and (II); compounds (1a) and (2a)]. This effect of simply changing from one biphasic solvent system to another is remarkable in heterogeneous catalytic transfer hydrogenolysis.

The crystal structure for benzyl ether (2a) has been reported recently (Brigas *et al.*, 1997); the molecule has a very short bond *p* and a very long bond *q*, like others in the series of ethers (2b)–(2d), with *p* + *q* equal to 2.769(6) Å, very similar to the aryl ethers (2b). Structural results for benzyl ether (1a) are presented here (Fig. 1, Table 1). Particularly notable are the short O—C1 and long O—C2 bonds. Indeed, although O—C1 (bond *p*) is typical of those in the series of compounds (1) and (2), O—C2 (bond *q*) is the longest such bond so far observed. The sum of these two bond lengths is 2.794(3) Å, in line with the other compounds, (1b)–(1d).




Fig. 1. The molecular structure of (1a) with atom labels and 50% probability ellipsoids for non-H atoms.

The geometry of the tetrazole ring is entirely typical of those previously reported, with all bond lengths and angles well within the ranges found for 245 such rings in 152 structures generated by a search of the Cambridge

Structural Database (October 1998 version; Allen & Kennard, 1993). This heterocyclic ring is essentially planar and the ether linkage lies almost in the same plane, but both phenyl groups are rotated significantly out of this plane (see torsion angles in Table 1).

Experimental

Synthesis of 5-benzyloxy-1-phenyltetrazole (1a): potassium *t*-butoxide (1.1 g, 9.8 mmol) was added to a stirred solution of benzyl alcohol (1.1 g, 10.17 mmol) in dry dimethyl sulfoxide (30 ml) under argon. When all the base had dissolved (10 min), 5-chloro-1-phenyltetrazole (2.0 g, 11.07 mmol) was added. The mixture was stirred at 348 K for 1 h and then poured into a large excess of ice/water. The resulting colourless solid was filtered off and dried in air. This solid was then dissolved in dichloromethane (50 ml), washed with saturated NaHCO₃ (3 × 20 ml) and HCl (1 M, 3 × 20 ml) and then dried (Na₂SO₄). After evaporation of the solvent, the residual solid was recrystallized from ethanol to afford compound (1a) as colourless needles (2.3 g, 88% yield; m.p. 421–422 K). Found: C 66.7, H 4.8, N 22.4%. C₁₄H₁₂N₄O requires C 66.7, H 4.8, N 22.2%. ¹H NMR (300 MHz, CDCl₃): δ 5.18 (2H, s), 7.26–7.52 (8H, m), 7.92–7.96 (2H, d, *J* = 6.6 Hz); IR ν_{max}: 1592, 1562 cm⁻¹; MS (*m/z*): 252 (*M*⁺).

Transfer reduction of 5-benzyloxy-1-phenyltetrazole: 10% palladium-on-charcoal (36 mg) was added to a stirred solution of 5-benzyloxy-1-phenyltetrazole, (1a) (50 mg, 0.19 mmol), and an internal standard (xylene, 101 mg) in tetrahydrofuran (THF) (20 ml) and the mixture was heated to reflux (336 K). Sodium phosphinate (161 mg, 1.52 mmol) in distilled water (3 ml) was added and the progress of the reaction was monitored for formation of toluene by gas chromatography and for the disappearance of the starting material by thin-layer chromatography. Transfer hydrogenolysis of the starting material to form only toluene and 5-phenyl-1-tetrazolone was complete in 10 min.

Transfer reduction of 3-benzyloxy-1,2-benzisothiazole 1,1-dioxide: this was carried out as described above for 5-benzyloxy-1-phenyltetrazole but using 44 mg (0.16 mmol) of the starting thiazole 1,1-dioxide (2a). Conversion to toluene and saccharin was complete in 15 min at 336 K. For either compound, (1a) or (2a), carrying out the reaction in exactly the same way but omitting the sodium phosphinate gave no hydrogenolysis, thereby showing that THF itself is not a hydrogen donor in this reaction.

Crystal data

C₁₄H₁₂N₄O
M_r = 252.28
 Monoclinic
*P*2₁/*c*
a = 9.5461 (15) Å
b = 6.2510 (10) Å
c = 20.773 (3) Å
 β = 98.008 (4)°
V = 1227.5 (3) Å³
Z = 4
D_x = 1.365 Mg m⁻³
D_m not measured

Mo *K*α radiation
 λ = 0.71073 Å
 Cell parameters from 2880 reflections
 θ = 2.15–28.12°
 μ = 0.091 mm⁻¹
T = 160 (2) K
 Plate
 0.50 × 0.45 × 0.15 mm
 Colourless

Data collection

Bruker AXS SMART CCD diffractometer
 ω rotation with narrow frames
 Absorption correction: none
 7286 measured reflections
 2911 independent reflections

1713 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.047$
 $\theta_{\text{max}} = 28.71^\circ$
 $h = -12 \rightarrow 12$
 $k = -8 \rightarrow 8$
 $l = -11 \rightarrow 27$
 Intensity decay: none

Refinement

Refinement on F^2
 $R[F^2 > 2\sigma(F^2)] = 0.040$
 $wR(F^2) = 0.090$
 $S = 0.910$
 2911 reflections
 173 parameters
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0403P)^2]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\text{max}} = 0.001$

$\Delta\rho_{\text{max}} = 0.212 \text{ e \AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.212 \text{ e \AA}^{-3}$
 Extinction correction: *SHELXTL* (Sheldrick, 1994)
 Extinction coefficient: 0.0089 (13)
 Scattering factors from *International Tables for Crystallography* (Vol. C)

Table 1. Selected geometric parameters (Å, °)

O—C1	1.3261 (15)	N2—N3	1.2904 (17)
O—C2	1.4676 (16)	N3—N4	1.3665 (15)
N1—C1	1.3159 (17)	N4—C1	1.3412 (18)
N1—N2	1.3791 (17)	C2—C3	1.498 (2)
C1—O—C2	114.28 (10)	O—C1—N4	120.73 (12)
N1—C1—O	128.72 (13)	O—C2—C3	106.99 (11)
N1—C1—N4	110.55 (12)		
C2—O—C1—N1	2.04 (19)	O—C2—C3—C8	102.69 (15)
C2—O—C1—N4	−177.99 (12)	C1—N4—C9—C14	−63.31 (19)
C1—O—C2—C3	−170.36 (11)	N3—N4—C9—C10	−62.34 (17)
O—C2—C3—C4	−77.53 (16)		

The data collection nominally covered over a hemisphere of reciprocal space, by a combination of three sets of exposures; each set had a different φ angle for the crystal and each exposure covered 0.3° in ω . Coverage was essentially complete for $\theta < 26^\circ$.

H atoms were placed geometrically and refined with a riding model, and with U_{iso} constrained to be $1.2 \times U_{\text{eq}}$ of the carrier atom.

Data collection: *SMART* (Siemens, 1995). Cell refinement: local programs. Data reduction: *SAINT* (Siemens, 1995). Program(s) used to solve structure: *SHELXTL* (Sheldrick, 1994). Program(s) used to refine structure: *SHELXTL*. Molecular graphics: *SHELXTL*. Software used to prepare material for publication: *SHELXTL* and local programs.

The authors gratefully acknowledge the financial support of EPSRC (UK), FCT (Portugal), the Eschenmoser Trust (UK) and the Royal Society (to AFB). The voluntary assistance of Sra Nuna C. P. Araujo for some of the work is greatly appreciated.

Supplementary data for this paper are available from the IUCr electronic archives (Reference: LN1073). Services for accessing these data are described at the back of the journal.

References

- Allen, F. H. & Kennard, O. (1993). *Chem. Des. Autom. News*, **8**, 31–37.
- Alves, J. A. C., Barkley, J. V., Brigas, A. F. & Johnstone, R. A. W. (1997). *J. Chem. Soc. Perkin Trans. 2*, pp. 669–677.
- Barkley, J. V., Cristiano, M. L. S., Johnstone, R. A. W. & Loureiro, R. M. S. (1997). *Acta Cryst. C53*, 383–386.
- Brigas, A. F., Goncalves, P. M. & Johnstone, R. A. W. (1997). *J. Chem. Soc. Perkin Trans. 1*, pp. 2603–2605.
- Brigas, A. F., Goncalves, P. M. & Johnstone, R. A. W. (1998). *Acta Cryst. C54*, 251–253.
- Brigas, A. F. & Johnstone, R. A. W. (1996). *Acta Cryst. C52*, 1293–1296.
- Bürgi, H.-B. & Dunitz, J. D. (1987). *J. Am. Chem. Soc.* **109**, 2924–2926.
- Chaloner, P. A. (1986). *Handbook of Coordination Catalysis in Organic Chemistry*, pp. 266–268; 851. London: Butterworths.
- Edwards, M. R., Jones, P. G. & Kirby, A. J. (1986). *J. Am. Chem. Soc.* **108**, 7067–7073.
- Jackson, A. E. & Johnstone, R. A. W. (1976). *Synthesis*, pp. 685–687.
- Johnstone, R. A. W., Wilby, A. H. & Entwistle, I. D. (1985). *Chem. Rev.* **85**, 129–170.
- Price, P. J. (1985). PhD thesis, University of Liverpool, England.
- Sheldrick, G. M. (1994). *SHELXTL*. Version 5. Bruker AXS Inc., Madison, Wisconsin, USA.
- Siemens (1995). *SMART and SAINT. Area Detector Control and Integration Software*. Bruker AXS Inc., Madison, Wisconsin, USA.

Acta Cryst. (1999). **C55**, 781–783

12-Formyl-9,10-dimethyl-9,10-dihydro-11-thia-9,10-ethanoanthracene

FRÉDÉRIC MINASSIAN, NADIA PELLOUX-LÉON, ANDRÉ DURIF AND YANNICK VALLÉE

LEDSS, UMR CNRS 5616, Université Joseph Fourier, BP 53, 38041 Grenoble CEDEX 9, France. E-mail: andre.durif@ujf-grenoble.fr

(Received 10 November 1998; accepted 17 December 1998)


Abstract

The structure of the title compound, $C_{18}H_{16}OS$, is described. Two conformers co-exist in the crystal structure with a very near spatial arrangement.

Comment

Thiaethanoanthracenes are convenient precursors for thioaldehydes (Pelloux-Léon & Vallée, 1997). Upon heating, they undergo cyclo-reversion and give anthracene and a compound containing a carbon–sulfur double bond. The title compound, (1), is a precursor for monothioglyoxal and can also be used as an intermediate

in the synthesis of various thiaethanoanthracenes (Arnaud *et al.*, 1994; Pelloux-Léon *et al.*, 1997). We have recently reported the nucleophilic attack of an enolate on the monothioglyoxal precursor (Minassian *et al.*, 1999). To understand the stereoselectivity observed in such additions, knowledge of the preferred conformation of (1) would be of interest. We have thus determined its atomic arrangement.



The present determination shows the existence of two conformers (Fig. 1) with very similar spatial conformations. Owing to this similarity, we think that both would give the same diastereoselectivity results. If one supposes that these conformers are also predominantly present in solution, then, according to Fig. 2, and assuming that the attack would occur from the less hindered side, the preferred face would be the *S* one. If the entering nucleophilic (Nu) group ranks third in the CIP priority rule (Cahn *et al.*, 1996), the relative configuration of the obtained product will be *R**,*S**. In order to simplify the text, we note the $C_{18}H_{16}OS$ conformer as *C* and the $C'_{18}H'_{16}O'S'$ conformer as *C'* hereafter.




Fig. 1. An ORTEP (Johnson, 1976) plot of the two conformers of (1) shown with 40% probability displacement ellipsoids.