Preliminary Note

Tetraphenylphosphonium bromide-catalyzed 'Halex' fluorination of chloroaryl sulfonyl chlorides

Hiroshi Suzuki, Hiroyuki Kageyama, Yasuo Yoshida and Yoshikazu Kimura*

Research and Development Department, Ihara Chemical Industry Co., Ltd. Fujikawacho, Ihara-gun, Shizuoka 421-33 (Japan)

(Received August 10, 1991; accepted October 22, 1991)

Abstract

Halogen-exchange of chloroaryl sulfonylchloride derivatives with spray-dried potassium fluoride was found to proceed efficiently by employing tetraphenylphosphonium bromide as a catalyst. Subsequent desulfonylation of the fluoroaryl sulfonyl derivatives readily afforded fluoroaromatics under acidic condition.

Recent studies in our laboratory have been directed towards the development of methodologies suitable for the preparation of aryl fluorides via the 'Halex' reaction [1]. Halex fluorination of aryl chlorides is well documented and can generally be a high-yielding process if there are strong electronwithdrawing groups situated *ortho* and/or *para* to the chlorine substituents of the aryl chlorides.

A very limited number of examples, however, are found in the literature for the reaction of chloro-substituted aryl sulfonyl chlorides (1) with potassium fluoride (KF). The only reported methods for the Halex fluorination of 1 are (a) a reaction in an autoclave at 275-290 °C without solvent [2] and (b) a reaction in the presence of tris(3,6,9-trioxadecyl)amine in an aprotic solvent [3].



These reactions seem not to be applicable to the large-scale preparation of fluoroaryl sulfonyl fluoride (2) due to the rather low yield (40-60%) and/ or the use of tedious reaction conditions.

We and others have recently reported that tetraphenylphosphonium bromide (Ph₄PBr) is a superior catalyst for Halex fluorinations [1, 4, 5].

^{*}Author to whom correspondence should be addressed.

During the course of these studies, we became interested in the Halex fluorination of 1 catalyzed by Ph_4PBr , because this reaction followed by desulfonylation would seem to be a new route to fluoroaromatics.

In a typical experiment, a mixture of 4-chlorobenzene sulfonyl chloride (105 g, 0.5 mol), spray-dried KF (116 g, 2 mol), Ph₄PBr (21 g, 0.05 mol) and anhydrous sulfolane (30 g) was stirred for 2 h at 215 °C after azeotropic dehydration by toluene. After cooling, the mixture was diluted with dichloromethane (0.5 l) and then filtered to remove inorganic matter. The residue was concentrated and distilled to give pure 4-fluorobenzene sulfonyl fluoride (78 g, 88%) having a boiling point of 63–64 °C/4 mmHg and ¹H NMR, IR and mass spectra identical with those of an authentic sample. The preparation of several varieties of **2** are listed in Table 1. Simple work-up of the reaction mixture gave virtually pure product.

Next, transformation of 2 to fluoroaromatics through desulfonylation was examined. Thus, 2,4-difluorobenzene sulfonyl fluoride was hydrolyzed

TABLE 1

Preparation	of	fluoroaryl	sulfonyl	fluoride	derivatives
-------------	----	------------	----------	----------	-------------

Run	1 ^a	KF	Catalyst	0.1	Temp, Time			(GC Yield/%)	D D
		(equiv)	(equiv)	Solvent	°C	h	2	[Isolated yield/%]b	B.P.
1	CISO2CI	4.0	Ph ₄ PBr (0.1)	Sulfolane	215	2	F-{SO₂F	(93) [88]	63-64°C/4mmHg
2	"	4.0	-	n	215	2	"	(29) ^c	
3	"	3.0	DMAPBr ^d (0.1)	"	210	5	"	(74)	
4	"	4.0	Ph ₄ PBr (0.1)	3,4-Dichloro- toluene	reflux	7	"	(59)	
5	ci–€–so₂ci	4.0	Ph ₄ PBr (0.1)	Sulfolane	180	4	F-√SO₂F	[74]	58-59°C/4mmHg
6	CI SO₂CI CI	4.0	Ph ₄ PBr (0.1)	"	180	5	CI F_SO₂F	[62]	96-97°C/7mmHg
7	CH₃_CI CI→_SO₂CI	4.0	Ph ₄ PBr (0.1)	"	200	1.5	CH ₃ _F F-{_}-SO₂F	[74]	110ºC/10mmHg
8	"	4.0	Ph ₄ PBr (0.1)	1,3-Dimethyl- imidazolidinone	200	1.5	"	[66]	
9	CI CI→ CH₃ CH₃	4.0	Ph ₄ PBr (0.1)	Sulfolane	180	4	F- CH₃ F-SO₂F	[77]	103-105°C /4mmHg
10	CI→CI CI→SO₂CI	4.0	Ph ₄ PBr (0.1)	Sulfolane	180	4	F-€-SO₂F	[32] ^e	69-70°C/2mmHg
11		2.5	Ph ₄ PBr (0.05)	"	130	16	F	[90]	115°C/0.3mmHg

^a Starting substrates were prepared from the corresponding any chlorides and chlorosulfonic acid.

^b All products gave satisfactory NMR, IR, and MS spectra.

^c 70% of chlorobenzene sulfonyl fluoride was given.

d

^e Reaction is not optimized.

to form sodium 2,4-difluorosulfonate, which was treated in 80% sulfuric acid at 200 °C. 1,3-Difluorobenzene, having a low boiling point (82 °C), was readily collected, as it distilled from the reaction mixture during the reaction

$$F \xrightarrow{F} SO_2F \xrightarrow{aq.NaOH} F \xrightarrow{F} SO_3Na \xrightarrow{80\%H_2SO_4} F \xrightarrow{F}$$

(62% yield) and was identified by GC-MS analysis. Similarly, 2,6-difluorotoluene was produced from 2,4-difluoro-3-methylbenzene sulfonyl fluoride in 79% yield.

As described above, we have succeeded in developing a novel catalyzed reaction which is useful in the synthesis of fluoroaryl sulfonyl fluoride derivatives. The method may offer a new synthetic route to some fluoroaromatics.

References

- 1 H. Suzuki, N. Yazawa, Y. Yoshida, O. Furusawa and Y. Kimura, Bull. Chem. Soc. Jpn., 63 (1990) 2010, and references cited therein.
- 2 (a) G. G. Yakobson, V. M. Vlasov and N. N. Voroshtsov, Jr., Zh. Vses. Khim. Ova., 10 (1965) 466 [Chem. Abstr., 63 (1965) 14 740e]; (b) idem, Zh. Obshch. Khim., 37 (1967) 163 [Chem. Abstr., 66 (1967) 94 741e].
- 3 G. Soula (to Rhone Poulenc), Eur. Pat. Appl. 32 077 (1981) [Chem. Abstr., 95 (1981) 168 793w].
- 4 J. H. Clark and D. J. Macquarrie, Tetrahedron Lett., 28 (1987) 111.
- 5 S. Kumai, Y. Furukawa, M. Sasabe and M. Matsuo, Rep. Res. Asahi Glass Co., 39 (1989) 317.