

Monitoring Carbonyl–Amine Reaction and Enolization of 1-Hydroxy-2-propanone (Acetol) by FTIR Spectroscopy

Varoujan A. Yaylayan,* Susan Harty-Majors, and Ashraf A. Ismail

Department of Food Science and Agricultural Chemistry, McGill University, 21,111 Lakeshore, Ste. Anne de Bellevue, Quebec, Canada H9X 3V9

Infrared absorption bands characteristic of the *aldehyde*, *keto*, and *enediol* forms of 1-hydroxy-2-propanone (acetol) were identified and used to study the effect of solvent on the absorption frequencies and the effect of temperature and acid/base catalysis on the enolization reactions. The data indicated that, in addition to water, acids and bases can catalyze the enolization of 1-hydroxy-2-propanone and that the temperature inversely effects the rate of enolization under basic conditions. However, under acidic conditions, increasing the temperature favors the enolization process. In addition, the reaction of 1-hydroxy-2-propanone with a primary and a secondary amine was also monitored by Fourier transform infrared spectroscopy. The data indicated that at room temperature the rate of amine reaction was faster than the rate of its catalysis of enolization; however, below room temperature, the rate of base-catalyzed enolization became comparable with the rate of carbonyl–amine reaction forming both Heyns and Amadori adducts.

Keywords: FTIR; α -hydroxycarbonyl moiety; enediol; enolization; 1-hydroxy-2-propanone (acetol); carbonyl–amine reactions; Heyns and Amadori product formation

INTRODUCTION

Many chemical transformations of reducing sugars, such as cyclizations, enolizations, and isomerizations, are initiated by the presence of the α -hydroxycarbonyl moiety. The stabilization of the initial imine adduct formed during Maillard reaction, through rearrangement reactions (Amadori and Heyns), is also due to the presence of the α -hydroxyl groups (Yaylayan and Huyghues-Despointes, 1994). The physical and chemical properties of reducing sugars in solution depend on the relative concentrations of different forms originating from the α -hydroxycarbonyl moiety. Their biological properties can also have similar dependence. It appears that the enediol forms of certain sugar derivatives in biological systems are the active forms with which enzymes react. The conversion of cytotoxic methylglyoxal into lactate, for example, by the enzyme glyoxylase I (thioester hydrolase glyoxylase II, EC 3.1.2.6), involves an enediolate intermediate of the methylglyoxal derivative (Hamilton and Creighton, 1992). Rubisco (ribulose-1,5-bisphosphate carboxylase, EC 4.1.1.39) adds carbon dioxide to the 2,3-enediolate form of ribulose-1,5-bisphosphate (Lorimer et al., 1993). Enediol forms are known to play an important role in metal-catalyzed oxidative degradation of reducing sugars specially in biological systems (Thornalley et al., 1984). The complexity in the population of reducing sugars in solution makes their study a difficult task, especially in hexoses and pentoses, where the presence of more stable furanose and pyranose forms renders the α -hydroxycarbonyl moiety undetectable at room temperature due to their lower concentrations (<1%). Solvent interference, especially water, makes the detection of aldehyde forms even more difficult due to hydration. Fourier transform

infrared spectroscopy (FTIR) has been used to study the effect of temperature (Yaylayan and Ismail, 1992) on acyclic forms of D-fructose, mutarotation of D-glucose and D-fructose (Back et al., 1984), and enolization and carbonyl group migration in selected sugars (Yaylayan and Ismail, 1995). Kobayashi et al. (1976) studied dimeric structures of D,L-glyceraldehyde and dihydroxyacetone by infrared and Raman spectroscopy. FTIR spectroscopy is ideally suited to study the different forms of α -hydroxycarbonyl moiety. Recently, the simplest α -hydroxyaldehyde (glycolaldehyde) was investigated by FTIR (Yaylayan et al., 1998). The data indicated that the glycolaldehyde cyclic dimer (2,5-dihydroxy-1,4-dioxane) undergoes a ring opening to form the acyclic dimer, which can recycize into the 2-hydroxymethyl-4-hydroxy-1,3-dioxolane structure. The acyclic dimer can dissociate into monomeric glycolaldehyde in equilibrium with the enediol form. In this paper, the enolization and its effect on the carbonyl–amine reactions of the simplest α -hydroxyketone (acetol) is investigated.

MATERIALS AND METHODS

All reagents and chemicals were purchased from Aldrich Chemical Co. (Milwaukee, WI) and used without further purification. All solvents used were of HPLC grade; D₂O was purchased from MSD Isotopes (Montreal, Quebec, Canada).


FTIR Analysis. Infrared spectra were recorded on a Nicolet 8210 Fourier transform infrared spectrometer (Madison, WI) purged with dry air and equipped with a deuterated triglycine sulfate (DTGS) detector. The spectra were acquired on a CaF₂ IR cell with a 25- μ m Teflon spacer at room temperature unless otherwise specified. A total of 128 scans at 4 cm⁻¹ resolution were co-added. Processing of the FTIR data was performed using GRAMS/386 version 3.01. Second-order derivatization was performed using the Savitsky–Golay function (9 points) to enhance closely absorbing peaks.

Temperature Studies. Sample solutions were placed in a CaF₂ IR cell with a 25- μ m Teflon spacer. The temperature of

* Corresponding author. Tel: 514-398-7918. Fax 514-398-7977. E-mail: yaylayan@agradm.lan.mcgill.ca.

Table 1. Carbonyl Absorption Frequencies of Relevant 2-Propanone (Acetone) Derivatives


acetone derivative ^a	carbonyl absorption frequency (cm ⁻¹)
acetone (g)	1740
acetone (D ₂ O)	1698
acetone (l)	1717
1-chloroacetone (g)	1743
1-chloroacetone (l)	1736
1,3-dichloroacetone (l)	1743
1,3-diaminoacetone (l)	1743
1-hydroxyacetone (D ₂ O)	1720
1-hydroxyacetone (g)	1750
1-hydroxyacetone (l)	1726

^a g, gas phase; l, liquid.**Figure 1.** Effect of temperature on the intensity of carbonyl absorption bands (1680–1740 cm⁻¹) of 1-hydroxy-2-propanone (2% D₂O).

2-propanone absorbs at 1726 cm⁻¹. In 5% NaOD, due to the formation of a bulky sodium alkoxide at the α -carbon, prevented 1-hydroxy-2-propanone to attain eclipsed conformation, and consequently, the absorption frequency was lowered to 1712 cm⁻¹.

A solution of 1-hydroxy-2-propanone in D₂O, also exhibited a weak absorption band in the carbonyl region centered at 1702 cm⁻¹ (see Figure 1) very close to the absorption frequency of acetone in D₂O (1698 cm⁻¹). Tentatively, this peak was assigned to the 1-hydroxy-2-propanone rotomer in which the hydroxyl group is near-staggered conformation (**1'**). By increasing the temperature of the 1-hydroxy-2-propanone solution, the intensity of non-H-bonded rotomer (**1'**) should increase on the expense of H-bonded rotomer (**1**). The initial temperature (30 °C) of the cell was raised by 1 °C/min, and every 5 min the temperature was kept constant for 15 min to record the spectra. During the heating cycle, the intensity of the band at 1702 cm⁻¹ increased during the heating cycle and decreased during the cooling cycle, indicating reversibility of the process.

Effect of Acid/Base Catalysis and Temperature on the Enolization of 1-Hydroxy-2-propanone. Enolization of 1-hydroxy-2-propanone (**1**, acetol) can be initiated by both α -carbons to produce the enediol **1b** and the enol **1a**. This process can be catalyzed by acids and bases. Bases in general are more efficient catalysts than acids. Under basic conditions, enolization initiated from the more acidic α -hydrogens will predominate (Rappe and Sachs, 1967). In acidic and basic conditions, 1-hydroxy-2-propanone exhibited peaks in the alkene region; however, unlike reducing sugars (Yaylayan and Ismail, 1995), no evidence of enolization was observed

**Figure 2.** Second-derivative spectra (1600–1800 cm⁻¹) of a 2% solution of 1-hydroxy-2-propanone in 5% DCl in D₂O at 80 °C (dotted line) and in 5% NaOD in D₂O at 30 °C (solid line).

in D₂O due to the presence of low concentrations of enolized species. Enolization in D₂O can be verified however by observing deuterium exchange at the α -hydrogens by GC/MS analysis. A solution of 1-hydroxy-2-propanone in D₂O was equilibrated overnight and analyzed by GC/MS. The data indicated the occurrence of deuterium exchange of all the six hydrogens of 1-hydroxy-2-propanone in the following percentages: M, 2%; M + 1, 10%; M + 2, 22%; M + 3, 28%; M + 4, 22%; M + 5, 12%; M + 6, 4%. When 1-hydroxy-2-propanone was equilibrated in 5% NaOD solution at 30 °C, the carbonyl peak was shifted to 1712 cm⁻¹, and two new bands appeared in the alkene region, a major band at 1694 cm⁻¹ and a weaker band at 1667 cm⁻¹ (Figure 2). Under basic conditions, the negatively charged alkoxide group should destabilize the developing partial negative charge on C-1 and make it less acidic relative to C-3. Accordingly, the stronger band at 1694 cm⁻¹ was assigned to the enol **1a**, and the weaker band at 1667 cm⁻¹ was assigned to the enediol **1b**. Under acidic conditions, the reverse trend should be observed where C-1 protons will be more acidic relative to C-3 protons due to the formation of alkyl oxonium ions. 1-Hydroxy-2-propanone was equilibrated in a 5% DCl solution at 30 °C, no absorption was observed in the alkene region, only the carbonyl band was observed at 1719 cm⁻¹. At temperatures higher than 65 °C, a new band appeared at 1667 cm⁻¹ and increased with temperature with corresponding decrease in the intensity of carbonyl band. This observation is consistent with the assignment of the enol and enediol bands under basic conditions. The process was not reversible during cooling cycle due to the formation of stable intramolecular H-bonding in **1b**. To prevent hydration and consequently to observe the carbonyl band due to the formation of 2-hydroxypropionaldehyde (**1c**), 1-hydroxy-2-propanone was analyzed in 100% triethylamine solution. The appearance of two bands in the carbonyl region (see Figure 3) indicated the occurrence of enolization. The addition of a small amount of D₂O to this solution caused the *aldehyde* band (1735 cm⁻¹) to disappear and the *keto* (1726 cm⁻¹) band to shift to 1720 cm⁻¹. However, neither enol nor enediol bands were observed in the alkene region, indicating the presence of very low concentrations of enol or enediol forms in the equilibrium mixture under basic conditions. This might be due to the enhanced acidity (~ 4–5 pK_a units) of enediol protons relative to α -hydrogens of their corresponding




Figure 3. Absorption of the carbonyl and enediol region (1660–1760 cm^{-1}) of a 2% solution of 1-hydroxy-2-propanone in (a) D_2O (solid line), (b) triethylamine (dotted line), and (c) NaOD (dashed line).

carbonyl forms (Bender and Williams, 1966). Although enol and enediol forms were not detected, they can be trapped as stable trimethylsilyl enol ethers using trimethylchlorosilane (TMS), a reagent commonly used for this purpose under alkaline conditions. A solution of 1-hydroxy-2-propanone in excess trimethylchlorosilane dissolved in acetonitrile was monitored at 30 $^{\circ}\text{C}$ for 6 h, after addition of a catalytic amount of triethylamine to initiate enolization. As expected, no aldehydic peak was detected due to trapping by TMS of the enediol intermediate (**1b**) leading to it. By the time of the first scan, a new band had appeared at 1698 cm^{-1} , in addition to the *keto* band at 1724 cm^{-1} , indicating the formation of trimethylsilyl ether of **1b** instead of aldehyde **1c**. However, over time, the *keto* band at 1724 cm^{-1} decreased in intensity with an increase of both 1698 cm^{-1} and a new band at 1714 cm^{-1} . The latter was assigned to the trimethylsilyl ether of **1a**.

To study the effect of temperature on enolization, 1-hydroxy-2-propanone solutions in TEA were equilibrated separately for 20 min at 30 and 60 $^{\circ}\text{C}$, and the relative percentages of *keto* and *aldehyde* forms were estimated based on the peak heights of the second-derivative spectra. At 30 $^{\circ}\text{C}$, the 1-hydroxy-2-propanone solution exhibited $\sim 40\%$ *keto* form and $\sim 60\%$ *aldehyde* form. However, at 60 $^{\circ}\text{C}$, the percent of *keto* form increased to $\sim 55\%$, indicating a slower rate of enolization at higher temperatures. At higher temperatures, the rate of enolization was also slowed in reducing sugars (Yaylayan and Ismail, 1995). It is interesting to note that under acid catalysis the rate of enolization increased with increasing temperature due to different mechanisms of catalysis (Lienhard and Wang, 1969).

Monitoring Carbonyl–Amine Reactions of 1-Hydroxy-2-propanone. To provide further evidence for the formation of 2-hydroxypropionaldehyde (**1c**) in the presence of basic amines, 1-hydroxy-2-propanone was reacted with amylamine (a reactive 1 $^{\circ}$ amine) and pyrrolidine (a reactive 2 $^{\circ}$ amine) in separate time-run experiments. These reactive amines, in addition to their ability to catalyze the enolization of 1-hydroxy-2-propanone (Bender and Williams, 1966), can also react with carbonyl compounds and form stable Amadori and Heyns rearrangement products (Scheme 2). Consequently, the carbonyl peaks due to **1** and **1c** should decrease over time with concomitant formation and an increase of new peaks in the carbonyl region during




Figure 4. Time-dependent second-derivative spectra (1660–1760 cm^{-1}) of a equimolar mixture of amylamine and 1-hydroxy-2-propanone at room temperature.




Figure 5. Time-dependent second-derivative spectra (1660–1760 cm^{-1}) of a equimolar mixture of pyrrolidine and 1-hydroxy-2-propanone at room temperature.

time-run experiments. When equimolar amount of amylamine was added to neat 1-hydroxy-2-propanone, directly on the IR cell, the *keto* band of 1-hydroxy-2-propanone centered at 1726 cm^{-1} decreased rapidly over time with the formation and increase of a new band at 1711 cm^{-1} as shown in Figure 4. The reaction was completed within 35 min. The same experiment was repeated with pyrrolidine. Again, the *keto* band of 1-hydroxy-2-propanone decreased with the formation of a new band at 1712 cm^{-1} . However, at the end of 35 min, a new band at 1724 cm^{-1} started to form as shown in Figure 5. In both systems, the *aldehyde* band expected at 1734 cm^{-1} was not detected. This could mean either enolization did not occur fast enough or the aldehyde formed reacted very fast. To study this interaction in more detail, the reaction of pyrrolidine with 1-hydroxy-2-propanone was further investigated at a lower initial temperature, and the reactants were mixed prior to FTIR analysis to avoid diffusion control of the reaction rates. Both reactants were cooled separately to -2°C and vortexed outside of the IR cell for 20 s, and a sample was removed for time-run analysis. The first scan, obtained 5 min after mixing, indicated the formation of a new peak at 1712 cm^{-1} and the presence of an unreacted *keto* peak of the 1-hydroxy-2-propanone at 1726 cm^{-1} (see Figure 6). The second scan, obtained 10 min after mixing, indicated the complete disappearance of the *keto* peak and the formation of a new band centered at 1724 cm^{-1} . For the next 50 min both new peaks increased continuously as shown in Figure 6. Monitoring the reaction for another hour indicated that




Figure 6. Time-dependent second-derivative spectra (1660–1760 cm^{-1}) of a equimolar mixture of pyrrolidine and 1-hydroxy-2-propanone mixed at -2°C and vortexed before monitoring at room temperature.

the new peaks reached their optimum concentrations after around 80 min. A small amount of D_2O was added to the IR cell after the final spectrum had been collected to identify the presence of aldehyde bands. The band at 1712 cm^{-1} disappeared due to hydration, and the band at 1724 cm^{-1} was shifted to 1719 cm^{-1} due to hydrogen bonding with D_2O . A similar shift of the keto band and disappearance of the aldehyde peak was also observed when D_2O was added to a solution of 1-hydroxy-2-propanone in TEA. Consequently, the band centered at 1712 cm^{-1} was assigned to the Heyns product (**3**).

When 1-hydroxy-2-propanone was equilibrated in TEA, a nonreactive tertiary amine, FTIR analysis indicated the presence of an aldehyde (**1c**) and a ketone (**1**) in the equilibrium mixture (Figure 3). Decreasing the temperature favored the enolization process and the formation of the aldehyde (**1c**). When a reactive primary amine (amylamine) was added to the neat 1-hydroxy-2-propanone, on the IR cell, and the reaction mixture was immediately scanned, only one carbonyl product was formed on the expense of 1-hydroxy-2-propanone (see Figure 4). Disappearance of this band after the addition of D_2O to the IR cell at the end of the last scan indicated the formation of an aldehyde (Heyns product **3**) rather than a ketone. Evidently, the rate of nucleophilic addition of the primary amine reaction with 1-hydroxy-2-propanone was much faster than the rate of its catalysis of enolization. As a result, only a Heyns product was formed. When the primary amine was replaced by a less nucleophilic but a more basic secondary amine (pyrrolidine), enolization was catalyzed to a small extent to generate toward the end of the reaction a new peak centered at 1724 cm^{-1} in addition to the major peak at 1712 cm^{-1} . The rate of formation of the new peak (1724 cm^{-1}) increased, as expected, by lowering the initial temperature of the reaction due to the decreased rate of the nucleophilic addition and the increased rate of enolization. Lowering the initial temperature allowed the enolization to proceed to a greater extent relative to the rate of nucleophilic addition and establish higher equilibrium concentrations of 2-hydroxypropionaldehyde (**1c**). This enabled the amine to react and produce the Amadori product **5**. The data indicated that within 5 min all the ketone **1** and the aldehyde **1c** have already been converted into their respective tetrahedral intermediates **2** and **4** and that the rate of conversion of tetrahedral intermediate into




Figure 7. Mass spectra of two products extracted from the reaction mixture of pyrrolidine and 1-hydroxy-2-propanone. A, Amadori product (**5**); B, Heyns product (**3**).

iminium ion and subsequently into **3** and **5** is much slower. Consequently, the rate of increase of the bands centered at 1712 and 1724 cm^{-1} over time, shown in Figure 6, corresponds to the rate of conversion of the tetrahedral intermediates into their corresponding Heyns (**3**) and Amadori (**5**) products. This observation is consistent with the literature data (Williams and Bender, 1966), which indicate that the conversion of the tetrahedral intermediate into imine is the rate-limiting step during the reaction of acetone with reactive amines. To provide direct evidence for the formation of Heyns (**3**) and Amadori products (**5**), the reaction solution was extracted with chloroform and analyzed by GC/MS. The analysis indicated the presence of two products with molecular weights of 127 amu and with mass spectral fragmentation patterns consistent with the assigned structures (see Figure 7).

ACKNOWLEDGMENT

V.Y. acknowledges funding for this research by the Natural Sciences and Engineering Research Council of Canada (NSERC).

LITERATURE CITED

- Back, D. M.; Michalska, D. F.; Polavarpu, P. L. Fourier Transform Infrared spectroscopy as a potential tool for the study of carbohydrates in aqueous solutions. *Appl. Spectrosc.* **1984**, *38*, 173–178.
- Bender, M. L.; Williams, A. Ketimine intermediates in amine-catalyzed enolization of acetone. *J. Am. Chem. Soc.* **1966**, *88*, 2502–2508.

- Coleman, W. M., III; Gordon B. M. Examination of the matrix isolation infrared spectra of organic compounds: part XIII. *Appl. Spectrosc.* **1989**, 43, 305–310.
- Davis, L. The structure of dihydroxyacetone in solution. *Bioorg. Chem.* **1973**, 2, 197–201.
- Hamilton, D. S.; Creighton, D. J. Inhibition of glyoxalase I by the enediol mimic S-(N-hydroxy-N-methylcarbamoyl)glutathione. The possible basis of a tumor-selective anticancer strategy. *J. Biol. Chem.* **1992**, 267, 24933–36.
- Kobayashi, Y.; Igarashi, T.; Takahashi, H.; Higasi, K. Infrared and Raman Studies of the Dimeric Structure of 1,3-Dihydroxyacetone, D(+)- and DL-Glycerldehyde. *J. Mol. Struct.* **1976**, 35, 85–99.
- Leinhard, G. E.; Wang, T.-C. On the mechanism of acid-catalyzed enolization of ketones. *J. Am. Chem. Soc.* **1969**, 91, 1146–1153.
- Lorimer, G. H.; Chen, Y.-R.; Hartman, F. C. A role for the -amino group of lysine-334 of ribulose-1,5-bisphosphate carboxylase in the addition of carbon dioxide to the 2,3-enediol(ate) of ribulose-1,5-bisphosphate. *Biochemistry* **1993**, 32, 9018–24.
- Rappe, C.; Sachs, W. H. Enolization of ketones III. Rate and orientation of base-catalyzed deuteration of some methylketones. *J. Org. Chem.* **1967**, 32, 4127–4128.
- Thornalley, P.; Wolf, S.; Crabe, J.; Stern, A. The autoxidation of glyceraldehyde and other simple monosaccharides under physiological conditions catalyzed by buffer ions. *Biochim. Biophys. Acta.* **1984**, 797, 276–287.
- Williams, A.; Bender, M. L. Studies on the mechanism of oxime and ketimine formation. *J. Am. Chem. Soc.* **1966**, 88, 2508–2513.
- Yasuda, T.; Okuyama, M.; Tanimoto, N.; Sekiguchi, S.; Ikawa, S. Intramolecular hydrogen bonding and molecular conformation. *J. Chem. Soc., Faraday Trans.* **1995**, 91, 3379–3383.
- Yaylayan, V.; Huygues-Despointes, A. Chemistry of Amadori Rearrangement products. *C. R. C. Crit. Rev. Food Sci. Nutr.* **1994**, 34, 321–369.
- Yaylayan, V. A.; Ismail, A. A. Determination of the effect of temperature on the concentration of keto form of D-fructose by FT-IR spectroscopy. *J. Carbohydr. Chem.* **1992**, 11 (2), 149–158.
- Yaylayan, V. A.; Ismail, A. A. Investigation of the enolization and carbonyl group migration in reducing sugars by FTIR spectroscopy. *Carbohydr. Res.* **1995**, 276, 253–265.
- Yaylayan, V. A.; Harty-Majors, S.; Ismail, A. A. Investigation of the mechanism of dissociation of glycolaldehyde dimer (2,5-dihydroxy-1,4-dioxane) by FTIR spectroscopy. *Carbohydr. Res.* **1998**, 309, 31–38.

Received for review November 23, 1998. Revised manuscript received March 24, 1999. Accepted April 2, 1999.

JF9812836