


This article was downloaded by: [McMaster University]

On: 18 February 2015, At: 12:25

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number:
1072954 Registered office: Mortimer House, 37-41 Mortimer Street,
London W1T 3JH, UK



Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/gpss20>

Preparation Of Tetraethyl-4-Hydroxyphenylmethylene-1,1-Bisphosphonate By Hydroxy-De-Diazonation Of The Corresponding Diazonium Salt Of Tetraethyl-4-Aminophenylmethylene-1,1-Bisphosphonate

T. Pieper^a & B. K. Keppler^a

^a Institute of Inorganic Chemistry, University of Vienna, Waehringerstr, 42, A-1090, Vienna, Austria

Published online: 19 Dec 2006.

To cite this article: T. Pieper & B. K. Keppler (2000) Preparation Of Tetraethyl-4-Hydroxyphenylmethylene-1,1-Bisphosphonate By Hydroxy-De-Diazonation of The Corresponding Diazonium Salt Of Tetraethyl-4-Aminophenylmethylene-1,1-Bisphosphonate, Phosphorus, Sulfur, and Silicon and the Related Elements, 165:1, 77-82, DOI: [10.1080/10426500008076326](https://doi.org/10.1080/10426500008076326)

To link to this article: <http://dx.doi.org/10.1080/10426500008076326>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our

platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

PREPARATION OF TETRAETHYL-4-HYDROXYPHENYLMETHYLENE-1,1-BISPHOSPHONATE BY HYDROXY-DE-DIAZONIATION OF THE CORRESPONDING DIAZONIUM SALT OF TETRAETHYL-4-AMINOPHENYLMETHYLENE-1,1-BISPHOSPHONATE

T. PIEPER and B.K. KEPPLER*

*Institute of Inorganic Chemistry, University of Vienna, Waehringerstr. 42,
A-1090 Vienna, Austria*

(Received May 02, 2000)

Bisphosphonates are widely used in diagnosis and therapy of different bone diseases. They are useful agents in osteotic vectorization of antitumor and antiinflammatory drugs because of their potential to accumulate in the inorganic bone matrix hydroxylapatite. Hence, there is great interest in alkyl-bisphosphonates, containing functional groups advantageous for further synthetic modification, as structural units for coupling with the drug. We report on the synthesis of tetraethyl-4-hydroxyphenylmethyle-1,1-bisphosphonate **8** by hydroxy-de-diazonation of the diazonium salt **7**, prepared by diazotation of the corresponding amine tetraethyl-4-aminophenylmethyle-1,1-bisphosphonate **6**.

Keywords: bisphosphonates; hydroxy-de-diazonation; drug targeting; bone malignancies

INTRODUCTION


Bisphosphonates, especially *gem*-bisphosphonates, show high affinity towards bone and other calcified tissues^[1]. They are used in diagnosis and therapy of osteoporosis^[2,3], tumor-induced hypercalcemia^[4,5] and Paget's disease^[6,7], due to their influence on bone metabolism and bone-seeking

* To whom correspondence should be sent.

properties^[8]. Their great potential to accumulate in the inorganic bone matrix hydroxylapatite makes them useful as carriers for cytotoxic or antibiotic substances, increasing the concentration of the drug in bone tissue. This drug targeting with bisphosphonates should result in an improved therapy of several bone malignancies. According to this concept, alkyl-bisphosphonates containing functional groups are useful structural units for coupling with the drug.

RESULTS AND DISCUSSION


Several alkyl-bisphosphonates containing an amino- or hydroxy-function have been synthesized^[9-11]. Nevertheless, unsubstituted 4-hydroxy- or 4-aminophenylmethylenetetraalkyl-1,1-bisphosphonates can't be found in the literature. The synthesis of the free bisphosphonic acid **1** (see Scheme 1) was described by Gross and Oszegowski^[12], but they obtained 4-hydroxyphenylmethylenetetraalkyl-1,1-bisphosphonate **1** by silylation and subsequent hydrolysis of 3,5-di-*tert*-butyl-4-hydroxyphenylmethylenetetraalkyl-1,1-bisphosphonate **2**, derived from 3,5-di-*tert*-butyl-4-hydroxybenzaldehyde. The *tert*-butyl groups are necessary for conversion of the aldehyde into the bisphosphonate. They are removed during hydrolysis. A direct synthesis of the corresponding alkylester of **1** was not possible and esterification of **1** with triethylorthoformate led to the 4-ethoxyphenylmethylenetetraalkyl derivative.



SCHEME 1

A convenient, direct synthesis of the corresponding ethylester of **1**, the title compound **8**, was found by hydroxy-de-diazonium of the diazonium

salt **7**, derived from the new tetraethyl-4-amino-phenylmethylene-1,1-bisphosphonate **6** (see Scheme 2). Compound **6** was synthesized by radical bromination of the diethyl-4-(trifluoracetyl-amino)phenylmethyl-phosphonate **3**, using *N*-bromosuccinimide (NBS), and subsequent Michaelis-Arbuzov reaction of the resulting diethyl-bromo-[4-(trifluoracetyl-amino)phenyl]methylene-phosphonate **4** with triethylorthophosphite (TEP), leading to the trifluoracetyl-protected amine **5**. Basic hydrolysis affords **6**, which can be used for further synthesis.



SCHEME 2

6 is a white solid that changes color to brown during storage. The trifluoracetyl-group is advantageous for storage, purification and also for protection of the amino-function during synthesis.

Thus, the title compounds **6** and **8**, as well as the diazonium salt **7**, are useful agents for further synthesis, especially in the field of osteotic vectorization of diverse drugs.

EXPERIMENTAL

The solvents used were dried following the usual methods. Diethyl-4-aminophenylmethyl-phosphonate was prepared by Arbuzov reaction of TEP with 4-nitrophenylmethylbromide (purchased from Merck) and subsequent catalytic hydrogenation.

All NMR spectra (^1H , ^{13}C , ^{31}P) were recorded on a Bruker WH 200 MHz spectrometer in $\text{dms}\text{-d}_6$. TMS and H_3PO_4 were used as internal standards.

Elemental analysis were performed using a Hearaeus C,H,N-Rapid system. P was determined as magnesium pyrophosphate.

Diethyl-4-(trifluoroacetyl-amino)phenylmethyl-phosphonate **3**. Diethyl-4-aminophenylmethyl-phosphonate (5.0 g, 0.02 mol) was dissolved in 25 ml of dry tetrahydrofuran. Trifluoroacetic anhydride (4.2 g, 0.02 mol) was added dropwise. After stirring for one hour at room temperature 30 ml of water were poured in and the mixture was concentrated at the rotavapor to about half of the volume. On standing, the product precipitated as clear, colorless crystals (m.p. 141°C).

Yield: 5.5 g (81%); NMR (in $\text{dms}\text{-d}_6$): ^{31}P : 26.6; ^1H : 1.15 [t, $^3J = 7.1$ Hz, 6 H], 3.21 [d, $^2J = 21.5$ Hz, 2 H], 3.92 [m, 4 H], 7.29 [d, $^3J = 8.4$ Hz, 2 H], 7.60 [d, $^3J = 8.4$ Hz, 2 H], 11.25 [s, 1 H]; ^{13}C : 16.18 [d, $^3J_{\text{PC}} = 5.7$ Hz], 46.70 [d, $^1J_{\text{PG}} = 135.2$ Hz], 61.42 [d, $^2J_{\text{PC}} = 6.5$ Hz], 115.85 [q, $^1J_{\text{FC}} = 288.6$ Hz], 120.97 [d, $^4J_{\text{PC}} = 2.5$ Hz], 129.71 [d, $^2J_{\text{PC}} = 9.1$ Hz], 130.26 [d, $^3J_{\text{PC}} = 6.7$ Hz], 134.84 [m, $^4J_{\text{FC}} = 3.7$ Hz], 154.45 [q, $^2J_{\text{FC}} = 36.8$ Hz].

Tetraethyl-4-(trifluoroacetyl-amino)phenylmethylene-1,1-bisphosphonate **5**. To a hot solution of 3.0 g (0.009 mol) of **3** in dry CCl_4 (60 ml) N-bromosuccinimide (1.76 g, 0.009 mol) was added. The mixture was refluxed and irradiated with a 500 W light bulb for three hours. After filtration of succinimide the filtrate was concentrated and 3.0 g of crude diethyl-bromo-[4-(trifluoroacetyl-amino)phenyl]methylene-phosphonate **4** precipitates. **4** was dissolved, without further purification, in 20 ml of dry tetrahydrofuran. After addition of 1.2 g (0.007 mol) of triethylphosphite (TEP) the mixture was refluxed for eight hours. Subsequently, all volatiles were removed in the vacuum and the resulting residue was treated with ether and filtered to give **5** as a white solid (m.p. 165°C).

Yield: 1.9 g (57%); NMR (in $\text{dms}\text{-d}_6$): ^{31}P : 19.3; ^1H : 1.03 [t, $^3J = 7.1$ Hz, 6 H], 1.18 [d, $^3J = 7.1$ Hz, 6 H], 3.85 [m, 4 H], 4.05 [m, 4 H],

4.34 [t, $^2J_{PH} = 24.8$ Hz, 1 H], 7.52 [d, $^3J = 8.6$ Hz, 2 H], 7.61 [d, $^3J = 8.6$ Hz, 2 H]; ^{13}C : 12.83 [m], 42.71 [t, $^1J_{PC} = 126.8$ Hz], 62.29 [m], 115.76 [q, $^1J_{FC} = 288.5$ Hz], 120.73, 128.03 [t, $^2J_{PC} = 7.8$ Hz], 130.95 [t, $^3J_{PC} = 6.2$ Hz], 135.39 [m, $^4J_{FC} = 2.6$ Hz], 154.44 [q, $^2J_{FC} = 36.9$ Hz].

$\text{C}_{15}\text{H}_{26}\text{F}_3\text{NO}_7\text{P}_2$ (475.34) calc.: C 42.96, H 5.51, N 2.95, P 13.03, found: C 43.12, H 5.60, N 3.08, P 13.29.

Tetraethyl-4-aminophenylmethylene-1,1-bisphosphonate **6**. 1.9 g [0.004 mol] of **5** were added to 50 ml of 0.1 M aqueous NaOH. After stirring at 100°C for three hours, the solution was extracted several times with CH_2Cl_2 . The combined organic layers were dried with Na_2SO_4 and the solvent was removed. The resulting oil crystallized and 1.3 g of a white solid were obtained (m.p. 92°C).

Yield: 1.3 g (88%); NMR (in dms $\text{O}-d_6$): ^{31}P : 18.8; ^1H : 1.01 [t, $^3J = 7.0$ Hz, 6 H], 1.18 [d, $^3J = 7.0$ Hz, 6 H], 3.81 [m, 4 H], 3.93 [t, $^2J_{PH} = 24.8$ Hz, 1 H], 3.99 [m, 4 H], 5.06 [s, 2H], 6.47 [d, $^3J = 8.3$ Hz, 2 H], 7.10 [dt, $^3J = 8.3$ Hz, $^4J_{PH} = 2.0$ Hz, 2 H]; ^{13}C : 15.91 [m], 42.51 [t, $^1J_{PC} = 131.7$ Hz], 61.93 [m], 113.49, 116.42 [t, $^2J_{PC} = 7.9$ Hz], 130.92 [t, $^3J_{PC} = 6.2$ Hz], 147.63.

$\text{C}_{15}\text{H}_{27}\text{NO}_6\text{P}_2$ (379.37) calc.: C 47.50, H 7.17, N 3.69, P 16.33, found: C 47.47, H 7.16, N 3.62, P 16.17.

Tetraethyl-4-hydroxyphenylmethylene-1,1-bisphosphonate **8**. 1.0 g (0.0026 mol) of **6** was dissolved in 2.5 ml of semi-concentrated hydrochloric acid. NaNO_2 (0.18 g, 0.0026 mol) in 1 ml of water was added within 10 minutes, while the temperature was kept below -10°C. The resulting diazonium salt **7** solution was diluted with 5 ml of water and heated for two hours at 100°C under generation of nitrogen. Then the yellow solution was extracted several times with CH_2Cl_2 . The combined organic layers were washed with water and dried with Na_2SO_4 . The solvent was removed and the resulting oil crystallized slowly. The product was recrystallized from ether, yielding a light yellow solid (m.p. 89°C).

Yield: 0.5 g (50%); NMR (in dms $\text{O}-d_6$): ^{31}P : 19.2; ^1H : 1.01 [t, $^3J = 7.0$ Hz, 6 H], 1.18 [d, $^3J = 7.0$ Hz, 6 H], 3.81 [m, 4 H], 4.00 [m, 4 H], 4.08 [t, $^2J_{PH} = 25.1$ Hz, 1 H], 6.68 [d, $^3J = 8.5$ Hz, 2 H], 7.26 [dt, $^3J = 8.5$ Hz, $^4J_{PH} = 2.0$ Hz, 2 H], 9.37 [s, 1 H]; ^{13}C : 16.00 [m], 42.34 [t, $^1J_{PC} = 131.2$ Hz], 62.11 [m], 114.89, 120.30 [t, $^2J_{PC} = 7.9$ Hz], 131.51 [t, $^3J_{PC} = 6.2$ Hz], 156.57.

$\text{C}_{15}\text{H}_{26}\text{O}_7\text{P}_2$ (380.31) calc.: C 47.37, H 6.89, P 16.29, found: C 47.24, H 6.93, P 16.04.

References

- [1] H. Fleisch, W.F. Naumann, *Am. J. Physiol.*, **200**, 1296 (1961).
- [2] R. P. Heany, P. D. Saville, *Clin. Pharmacol. Ther.*, **20**, 593 (1976).
- [3] S. E. Papapioulos, et al., *Bone*, **13**, S41 (1992).
- [4] A. Jung, H. Fleisch, *Schweiz. Med. Wochenschr.*, **111**, 1878 (1981).
- [5] H. Fleisch, *Drugs*, **42**, 919 (1991).
- [6] R. L. Hilderbrand, *The Role of Phosphonates in Living Systems*, (CRC-Press, Boca Raton, Florida, 1983).
- [7] D. K. J. Hosking, *Drugs*, **40**, 829 (1990).
- [8] H. Blum, *J. Prakt. Chem.*, **336**, 492 (1994).
- [9] W. Winckler, T. Pieper, B. K. Keppler, *Phosphorus, Sulfur and Silicon*, **112**, 137 (1996). [10] G. Sturtz, J. Guervenou, *Synthesis*, **8**, 661 (1991).
- [11] G. Sturtz, H. Couthon, *C. R. Acad. Sci. Paris*, **t. 316**, 181 (1993).
- [12] H. Gross, S. Oszegowski, *Phosphorus, Sulfur and Silicon*, **47**, 1 (1990).