


Radiochemical purity of this material was determined to be  $\geq 98\%$  by TLC and HPLC [HPLC system: Varian 5500 instrument, Whatman Partisil 5,  $4.6 \times 250$  mm column, mobile phase 13% 80:20:05  $\text{CHCl}_3/\text{MeOH}/\text{Et}_3\text{N}$  in  $\text{CHCl}_3$ , flow rate 1.0 mL  $\text{min}^{-1}$ , Varian UV-200 UV detector, Radiomatic HP30 Flo-One


Cell suspension (150  $\mu$ L) was incubated at 37 °C for 15 min with 0.1–10 nM of radioligand in 20 mM Tris-HCl, 154 mM NaCl, pH 7.5 solution in a final volume of 250  $\mu$ L. Nonspecific binding was determined by 10  $\mu$ M zacopride. Incubations were terminated by rapid filtration through 0.2% PEI-soaked borosilicate filtermats and washed with ice-cold Tris-HCl-NaCl buffer (3  $\times$  5 mL). Filters were soaked overnight in 4 mL of Ecolume, and the retained radioactivity was determined with an LKB liquid-scintillation counter. Data were analyzed with the Lunden 1 software program.

## 0022-2623/90/1833-3023\$02.50/0 © 1990 American Chemical Society

**Table I.** *trans*-3,4-Dihydro-4-(2-oxopyrrolidin-1-yl)-2*H*-pyranopyridin-3-ols and Analogues

compd	R <sub>1</sub>	N-posn	R <sub>2</sub>	R <sub>3</sub>	R <sub>4</sub>	R <sub>5</sub>	% yield	mp, °C	recryst solv <sup>a</sup>	formula	anal. <sup>b</sup>	dose, mg/kg po	% max fall in <sup>c</sup> bp ± SEM
1 <sup>d</sup>	CN		-(CH <sub>2</sub> ) <sub>3</sub> -		OH	H						0.1	13 ± 5
												0.3	39 ± 4
												1.0	47 ± 1
6	H	5	-(CH <sub>2</sub> ) <sub>3</sub> -		OH	H	31	214–215	E-P	C <sub>14</sub> H <sub>18</sub> N <sub>2</sub> O <sub>3</sub>	C, H, N	10.0	15 ± 3
7		6	-(CH <sub>2</sub> ) <sub>3</sub> -		OH	H	12	253	E	C <sub>14</sub> H <sub>18</sub> N <sub>2</sub> O <sub>3</sub>	C, H, N	1.0	39 ± 7
												3.0	41 ± 4
8	H	7	-(CH <sub>2</sub> ) <sub>3</sub> -		OH	H	21	221–225	E-M	C <sub>14</sub> H <sub>18</sub> N <sub>2</sub> O <sub>3</sub>	H, N; C <sup>e</sup>	1.0	7 ± 3
												10.0	41 (2)
9	H	8	-(CH <sub>2</sub> ) <sub>3</sub> -		OH	H	30	202–203	E-P	C <sub>14</sub> H <sub>18</sub> N <sub>2</sub> O <sub>3</sub>	C, H, N	10.0	5 ± 3
10		6	-(CH <sub>2</sub> ) <sub>4</sub> -		OH	H	16	241–243	E	C <sub>15</sub> H <sub>20</sub> N <sub>2</sub> O <sub>3</sub>	C, H, N	0.1	24 ± 4
												0.3	55 ± 4
												1.0	56 ± 4
11	O	6	-(CH <sub>2</sub> ) <sub>4</sub> -		OH	H	75	290–291	E-M	C <sub>15</sub> H <sub>20</sub> N <sub>2</sub> O <sub>4</sub>	C, H, N	0.3	39 ± 0
												1.0	68 ± 1
12		6	-(CH <sub>2</sub> ) <sub>4</sub> -		bond		31	108–113	E-P	C <sub>15</sub> H <sub>18</sub> N <sub>2</sub> O <sub>2</sub>	C, H, N	1.0	14 ± 4
												3.0	37 ± 3
13		6	H	Me	OH	H	32	208–210	E	C <sub>12</sub> H <sub>16</sub> N <sub>2</sub> O <sub>3</sub>	C, H, N	0.3	14 ± 2
												1.0	43 ± 5

<sup>a</sup>E = EtOAc, M = MeOH, P = pentane. <sup>b</sup>Analyses for the elements indicated were within ±0.4% of the theoretical values. <sup>c</sup>Systolic blood pressure was measured indirectly at intervals from 1 to 6 h in groups of six SH rats per dose level. On occasion pulses were determined from only (*n*) SH rats. <sup>d</sup>See ref 6. <sup>e</sup>C: calcd, 64.11; found, 63.67.

**Table II.** *trans*-6-Alkyl-3,4-dihydro-4-(2-oxopyrrolidin-1-yl)-2*H*-1-benzopyran-3-ols and Analogues

compd	R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>	% yield	mp, °C	recryst solv <sup>a</sup>	formula	anal. <sup>b</sup>	dose, mg/kg po	% max fall in <sup>c</sup> bp ± SEM
14	CF <sub>3</sub>	-(CH <sub>2</sub> ) <sub>3</sub> -		44	181–182	E	C <sub>16</sub> H <sub>18</sub> NO <sub>3</sub> F <sub>3</sub>	H, N; C <sup>d</sup>	0.1	17 ± 2
									0.3	26 ± 4
									1.0	43 ± 5
15 <sup>e</sup>	H	-(CH <sub>2</sub> ) <sub>3</sub> -							10.0	9 ± 4
16	Me	-(CH <sub>2</sub> ) <sub>3</sub> -		35	187–188	E-P	C <sub>16</sub> H <sub>21</sub> NO <sub>3</sub>	C, H, N	3.0	12 ± 6
									10.0	49 ± 7 (4)
17	Me	-(CH <sub>2</sub> ) <sub>4</sub> -		15	162–164	E	C <sub>17</sub> H <sub>23</sub> NO <sub>3</sub>	C, H, N	0.3	16 ± 5
									1.0	33 ± 2
									3.0	56 ± 2
18	Et	-(CH <sub>2</sub> ) <sub>3</sub> -		49	154–163	E-PE	C <sub>17</sub> H <sub>23</sub> NO <sub>3</sub>	C, H, N	0.3	11 ± 4
									1.0	54 ± 2
19	Me(CH <sub>2</sub> ) <sub>2</sub>	-(CH <sub>2</sub> ) <sub>3</sub> -		15	139–140	E	C <sub>18</sub> H <sub>25</sub> NO <sub>3</sub>	C, H, N	1.0	17 ± 7
									3.0	42 ± 7 (5)
20	Me <sub>2</sub> CH	-(CH <sub>2</sub> ) <sub>3</sub> -		32	167–173	E	C <sub>18</sub> H <sub>25</sub> NO <sub>3</sub>	C, H, N	0.3	17 ± 4
									1.0	36 ± 5
									3.0	60 (2)
21	Me <sub>3</sub> C	-(CH <sub>2</sub> ) <sub>3</sub> -		16	227–228	E-PE	C <sub>19</sub> H <sub>27</sub> NO <sub>3</sub>	C, H, N	0.3	9 ± 5
									1.0	39 ± 9
									3.0	49 ± 6 (4)
22	<i>c</i> -C <sub>5</sub> H <sub>9</sub>	-(CH <sub>2</sub> ) <sub>3</sub> -		27	179–184	E	C <sub>20</sub> H <sub>27</sub> NO <sub>3</sub>	C, H, N	1.0	24 ± 2
									10.0	66 ± 2 (3)
23	Ph	-(CH <sub>2</sub> ) <sub>3</sub> -		10	217–220	E-P	C <sub>21</sub> H <sub>23</sub> NO <sub>3</sub>	C, H, N	3.0	14 ± 5
									10.0	20 ± 2 (4)
24	Me	H	Me	21	188–190	E	C <sub>14</sub> H <sub>19</sub> NO <sub>3</sub> ·0.25H <sub>2</sub> O	C, H, N	0.3	9 ± 4
									3.0	20 ± 6


<sup>a</sup>E = EtOAc, P = pentane, PE = 60–80 °C petroleum ether. <sup>b,c</sup>See footnotes *b* and *c* of Table I. <sup>d</sup>C: calcd, 58.36; found, 59.63. <sup>e</sup>See ref 3.

In accordance with the requirement for a strong electron-withdrawing group, 6-CF<sub>3</sub> compound 14 showed good

activity (Table II), while the unsubstituted compound 15 did not lower blood pressure at the top dose investigated (10 mg/kg).<sup>3</sup> However, in stark contrast to these results, the 6-methyl compound 16 had surprisingly good activity at 10 mg/kg, producing a fall of approximately 50%. This result prompted an investigation of a series of 6-alkyl

(5) Evans, J. M.; Cassidy, F.; Stemp, G. Partly presented at the 197th ACS National Meeting, Dallas, Texas, 1989, Symposium on Potassium Channels.

Scheme I




analogues of compound 16 (see Table II).<sup>5</sup>

### Chemistry

Synthesis of pyranopyridines 6–10 (see Table I) required as starting materials the appropriate 2,2-dimethyl-2H-pyranopyridines 25, the syntheses of which have been reported recently.<sup>6</sup> Thus by the established route for compound 1, compounds 6, 7, 9, and 10 were prepared (Scheme I) via bromohydrin 26 and epoxide 27. Compounds 6, 7, and 9 were obtained by the action of pyrrolidone and compound 10 was obtained by the action of piperidone on the appropriate epoxide 27 in the presence of NaH base. The synthesis of compound 8 required changes in strategy as treatment of pyrano[2,3-c]pyridine 25 (7-N) with moist NBS failed to give a bromohydrin. However, conversion of pyrano[2,3-c]pyridine 25 to its *N*-oxide with *m*-CPBA followed by treatment with moist NBS gave bromohydrin 26 (7-N<sup>+</sup>-O<sup>-</sup>). Conversion of bromohydrin 26 (7-N<sup>+</sup>-O<sup>-</sup>) to epoxide 27 (7-N<sup>+</sup>-O<sup>-</sup>) was followed by treatment with ammonia and amino alcohol 28 acylated with chlorobutyryl chloride to give (chlorobutyryl)amino compound 29, as epoxide 27 (7-N<sup>+</sup>-O<sup>-</sup>) failed to react with pyrrolidone under the usual conditions described for the preparation of compounds 6, 7, and 9. Ring closure under basic conditions yielded pyrrolidone 30. Deoxygenation of 30 with hexachlorodisilane<sup>7</sup> provided the required compound 8. Treatment of compound 10 with *m*-CPBA gave the *N*-oxide 11, and dehydration of 10 with NaH gave the dihydro compound 12 (Scheme II). Finally, the pyrano[3,2-c]pyridine epoxide 27 (6-N) was treated with ammonia, and amino alcohol 31 was acylated with acetyl chloride to give compound 13 (Scheme II).

Scheme II



The synthesis of the 6-CF<sub>3</sub> compound 14, the 6-alkyl compounds 16–22, and the 6-phenyl compound 23 also followed the established synthetic route (see Scheme III) devised for compound 1. Thus the appropriate phenols were alkylated with 3-chloro-3-methylbut-1-yne under phase-transfer catalytic conditions to furnish 3-methyl-3-(4-substituted-phenoxy)-but-1-ynes 32. These were thermally cyclized in boiling *o*-dichlorobenzene to give 6-substituted chromenes 33 as oils, which were characterized by NMR spectroscopy before being converted directly into bromohydrins 34 by treatment with aqueous NBS. Action of base yielded epoxides 35 which were treated with the appropriate cyclic amide and 1 equiv of NaH to give compounds 14 and 16–23 (Table II). The 6-methyl epoxide 35 (R = Me)<sup>8</sup> was converted to amino alcohol 36 and reacted with acetyl chloride to give compound 24.

### Results and Discussion

Compounds were evaluated for oral antihypertensive activity in the SHR. Systolic blood pressure, recorded from the tail, was determined before dosing and at various time intervals during the ensuing 6 h. Maximum falls in blood pressure obtained for all the compounds (Tables I and II) occurred at 1–4 h postdose with some recovery to the predose level of blood pressure being observed at 6 h.

In the four isomeric pyranopyridine analogues 6–9, structure–activity relationships are in line with previous results obtained<sup>3</sup> when variation of the position of substituents in the aromatic ring was studied and in agreement with the predictions from the modeling of two-dimensional electrostatic potentials. Thus the most potent compound (Table I) is pyrano[3,2-c]pyridine compound 7, while the pyrano[2,3-c]pyridine compound 8 only possessed reasonable activity at the highest dose. The pyrano[3,2-b]- and -[2,3-b]pyridines compounds 6 and 9, respectively, showed little activity at the highest dose employed. In the potent pyrano[3,2-c]pyridine series the homologous piperidone compound 10 showed enhanced activity over pyrrolidone 7, a differential in activity first observed<sup>3</sup> for the piperidone analogue of compound 1.


Preparation of the *N*-oxide of compound 10 resulted in an analogue, 11, of similar potency, but the unsaturated variant 12 of compound 10 was found to be less active, thus illustrating again the variability in comparative activities

(6) Evans, J. M.; Stemp, G. *Synth. Commun.* 1988, 18, 1111.

(7) Naumann, K.; Zon, G.; Mislow, K. *J. Am. Chem. Soc.* 1969, 91, 2788.

(8) Evans, J. M.; Fake, C. S.; Hamilton, T. C.; Poyser, R. H.; Watts, E. A. *J. Med. Chem.* 1983, 26, 1582.

## Scheme III



of 3-hydroxy compounds and dehydro compounds that we have reported previously.<sup>3</sup>

In accord with the earlier work, compound 14 (Table II) which contains the strong electron-withdrawing CF<sub>3</sub> group at C(6) showed good activity, being approximately equiactive with compound 1. Unsubstituted compound 15 was inactive at the dose tested, but 6-methyl compound 16, which is similar in shape but not in electronic effect to 6-CF<sub>3</sub> compound 14, had surprisingly good activity at 10 mg/kg. The possibility arises that the activity of compound 16 might be ascribed to the shape and/or lipophilicity of the C(6) substituent, and that the enhanced activity of compound 14 may be due to the additional electron-withdrawing properties of the CF<sub>3</sub> substituent when compared with a methyl substituent. Again piperidone compound 17 showed a greater potency than pyrrolidone compound 16, as was noted in the pyrano[3,2-*c*]pyridine series above.

Extension of the methyl group to ethyl as in compound 18 substantially enhanced the activity, but the *n*-propyl compound 19 was only about 1/3 as potent as 6-ethyl compound 18. However, introduction of branched alkyl groups gave isopropyl (20) and *tert*-butyl (21) compounds of similar potency to the ethyl compound 18, while cyclopentyl analogue 22 was at least 3-fold less active than compound 18. Replacement of the alkyl group by a phenyl ring (compound 23) reduced activity to modest levels.

Representative acyclic amido analogues, 4-acetylamino compounds 13 (Table I) and 24 (Table II), have been prepared in the pyrano[3,2-*c*]pyridine and the 6-alkyl series, respectively. These compounds compare very favorably in potency with the equivalent pyrrolidones 7 and 16, respectively. Such equivalence has been reported previously<sup>4</sup> for the 4-(acetylamino) analogue of compound 1.

## Experimental Section

Melting points were determined with a Büchi capillary melting point apparatus and are uncorrected. NMR spectra were recorded on a Varian EM 360A at 60 MHz, a Varian CFT-20 at 80 MHz, or a JEOL GX 270, and mass spectra were recorded on a VG 70-70 or 70 ZAB or a JEOL DX 303 at 70 eV, respectively, and were in agreement with the structures cited. HF<sub>254</sub> silica gel plates (2 mm) were used for Chromatotron chromatography (radial chromatography). Petroleum ether refers to that fraction boiling between 60 and 80 °C. Two dimensional electrostatic potential maps were generated with a modified VSS procedure<sup>9</sup> from partial charges calculated by the semiempirical CNDO method.<sup>10</sup>

**3-Methyl-3-(4-substituted-phenoxy)but-1-yne (32).** The appropriate phenol (0.06 mol) and NaOH (0.1 mol) were added to a stirred mixture of H<sub>2</sub>O (60 mL) and CH<sub>2</sub>Cl<sub>2</sub> (60 mL), followed by benzyltrimethylammonium hydroxide (40% in MeOH, 0.06 mol) and 3-chloro-3-methyl-but-1-yne (0.1 mol). The mixture was

stirred at room temperature for 4–11 days, and the layers were separated. The aqueous layer was extracted with CHCl<sub>3</sub>, and the combined organic layer was concentrated in vacuo and taken up in Et<sub>2</sub>O. The solution was washed with H<sub>2</sub>O, 10% NaOH, and brine and dried over anhydrous MgSO<sub>4</sub>. Filtration and evaporation gave compounds 32 in yields of 19–50%.

**2,2-Dimethyl-6-substituted-2H-1-benzopyrans (33).** Phenoxybutynes 32 (0.05 mol) were heated in *o*-dichlorobenzene (2 mL/g of 32) under N<sub>2</sub> for 2–12 h. Solvent was removed under vacuum, and crude 33 was used directly or, more usually, purified by chromatography on silica gel with EtOAc as eluant in yields of 20–90%. Compound 33, R = C(Me)<sub>3</sub>: NMR (CDCl<sub>3</sub>) δ 1.27 (s, C(Me)<sub>3</sub>), 1.40 (s, C(Me)<sub>2</sub>), 5.50 (d, *J* = 9 Hz, H-3), 6.25 (d, *J* = 9 Hz, H-4), 6.60 (d, *J* = 8 Hz, H-8), 7.10 (m, H-7 and H-5).

**trans-3-Bromo-3,4-dihydro-2,2-dimethyl-6-substituted-2H-1-benzopyran-4-ols (34).** To a solution of the appropriate 2,2-dimethylbenzopyran 33 (10 mmol) in DMSO (35 mL) and H<sub>2</sub>O (20 mL) was added NBS (20 mmol) and the mixture was stirred for 1 h. The solution was diluted with H<sub>2</sub>O and extracted with EtOAc. The organic layer was washed with H<sub>2</sub>O and brine and dried over MgSO<sub>4</sub>. Filtration and evaporation gave bromohydrins 34, usually as oils (70–90%) which were characterized by NMR and high-resolution MS before conversion to epoxides 35. Compound 34 (R = Me<sub>2</sub>CH): NMR (CDCl<sub>3</sub>) δ 1.20 (d, *J* = 7 Hz, Me<sub>2</sub>CH), 1.37 (s, C(Me)<sub>2</sub>), 1.55 (s, C(Me)<sub>2</sub>), 2.80 (sept, *J* = 7 Hz, Me<sub>2</sub>CH), 4.03 (d, *J* = 9 Hz, H-3), 4.80 (m, H-4), 6.55 (d, *J* = 8 Hz, H-8), 6.80–7.20 (m, H-5, H-7); mass spectrum, observed M<sup>+</sup> at *m/z* 298.0563, 300.0553; C<sub>14</sub>H<sub>18</sub>O<sub>2</sub>Br requires 298.0568, 300.0548. Compound 34 (R = Ph): 69%; mp 135–139 °C (EtOAc–petroleum ether). Anal. (C<sub>17</sub>H<sub>17</sub>O<sub>2</sub>Br) C, H, Br. Compound 34 (R = *c*-C<sub>5</sub>H<sub>9</sub>): 92%; mp 121–122 °C (EtOAc–petroleum ether). Anal. (C<sub>16</sub>H<sub>21</sub>O<sub>2</sub>Br) C, H. Compound 34 (R = *t*-Bu): 18%; mp 122–123 °C (EtOAc–petroleum ether). Anal. (C<sub>15</sub>H<sub>21</sub>O<sub>2</sub>Br·0.5H<sub>2</sub>O) C, H.

**3,4-Epoxy-3,4-dihydro-2,2-dimethyl-6-substituted-2H-1-benzopyrans (35).** Bromohydrins 34 (5 mmol) were stirred with KOH pellets (5 mmol) in Et<sub>2</sub>O (250 mL) at room temperature for 1–4 days. Filtration and evaporation gave epoxides 35, which were used directly. Compound 35 (R = Me<sub>2</sub>CH): NMR (CDCl<sub>3</sub>) δ 1.20 (d, *J* = 7 Hz, Me<sub>2</sub>CH) overlapping 1.20 (s, C(Me)<sub>2</sub>), 1.50 (s, C(Me)<sub>2</sub>), 2.80 (sept, *J* = 7 Hz, Me<sub>2</sub>CH), 3.40 (d, *J* = 4 Hz, H-3), 3.80 (d, *J* = 4 Hz, H-4), 6.60 (d, *J* = 8 Hz, H-8), 6.80–7.30 (m, H-5, H-7).

**trans-4-Amido-3,4-dihydro-2,2-dimethyl-6-substituted-2H-1-benzopyrans (14, 16–23).** The requisite cyclic amide (5 mmol) was treated with 80% NaH (5 mmol) and the appropriate epoxide (5 mmol) in Me<sub>2</sub>SO (40 mL) with stirring under N<sub>2</sub> for 2–6 h. The reaction mixture was diluted with H<sub>2</sub>O and extracted with EtOAc. The organic layer was washed with H<sub>2</sub>O and brine and dried over anhydrous MgSO<sub>4</sub>. Filtration and evaporation gave the title compounds as crude solids which were purified by chromatography and recrystallization (see Table II). Compound 14: NMR (CDCl<sub>3</sub>) δ 1.30 (s, C(Me)<sub>2</sub>), 1.50 (s, C(Me)<sub>2</sub>), 2.15 (m, NCH<sub>2</sub>CH<sub>2</sub>), 2.50 (m, NCOCH<sub>2</sub>), 3.20 (m, NCH<sub>2</sub>), 3.75 (d, *J* = 10 Hz, H-3), 4.60 (br s, OH), 5.25 (d, *J* = 10 Hz, H-4), 6.80 (d, *J* = 8 Hz, H-8), 7.10 (d, *J* = 2 Hz, H-5), 7.30 (dd, *J* = 8, 2 Hz, H-7).

**2,2-Dimethyl-2H-pyrano[2,3-*c*]pyridine N-Oxide (25 N-Oxide).** To a stirred solution of 2,2-dimethyl-2H-pyrano[2,3-*c*]pyridine<sup>6</sup> (25, 8.64 g, 54 mmol) in CHCl<sub>3</sub> was added *m*-CPBA (9.15 g, 53 mmol) portionwise during 15 min. The solution was then boiled for 1 h, cooled, and evaporated, and the residue was

(9) Giessner-Prettre, C. *QCPE* 1974, No. 249.

(10) Dobosh, P. A. *QCPE* 1968, No. 141.

chromatographed on silica gel (500 g). Elution with 5% MeOH-CHCl<sub>3</sub> gave **25** (*N*-oxide) (5.96 g, 61%) as a colorless gum: NMR (CDCl<sub>3</sub>)  $\delta$  1.45 (s, C(Me)<sub>2</sub>), 5.73 (d, *J* = 10 Hz, H-3), 6.27 (d, *J* = 10 Hz, H-4), 6.87 (d, *J* = 6 Hz, H-5), 7.75 (m, H-6, H-8).

**trans-3-Bromo-3,4-dihydro-2,2-dimethyl-2H-pyranopyridin-3-ols (26).** 2,2-Dimethyl-2H-pyranopyridines **25** were treated in a manner identical with that for 2H-1-benzopyrans **33** with aqueous NBS. The crude solids were purified by recrystallization, after optional chromatography. Compound **26** (3,2-*b*): 59%; mp 133–135 °C (radial chromatography eluted with 30% ethyl acetate–petroleum ether). Anal. (C<sub>10</sub>H<sub>12</sub>NO<sub>2</sub>Br) C, H, N. Compound **26** (3,2-*c*): 34%; mp 140–141 °C (EtOAc–pentane). Anal. (C<sub>10</sub>H<sub>12</sub>NO<sub>2</sub>Br) C, H, N. Compound **26** (2,3-*c* *N*-oxide): 68%; mp 185–187 °C (EtOAc–MeOH). Anal. (C<sub>10</sub>H<sub>12</sub>NO<sub>3</sub>Br) C, H, N. Compound **26** (2,3-*b*): 89%; mp 163–165 °C (EtOAc). Anal. (C<sub>10</sub>H<sub>12</sub>NO<sub>2</sub>Br) C, H, N.

**3,4-Epoxy-3,4-dihydro-2,2-dimethyl-2H-pyranopyridines (27).** Bromohydrins **26** were treated as bromohydrins **34** with KOH/Et<sub>2</sub>O to give epoxides **27**, which were used directly. Bromohydrin **26** (2,3-*c* *N*-oxide, 1.56 g, 5.5 mmol) and powdered KOH pellets (2.0 g, 36 mmol) were stirred and heated under reflux in THF (150 mL) for 4 h. The solution was cooled, filtered, and evaporated to give epoxide **27** (2,3-*c* *N*-oxide, 1.01 g, 91%) as a glass: NMR (CDCl<sub>3</sub>)  $\delta$  1.31 (s, C(Me)<sub>2</sub>), 1.58 (s, C(Me)<sub>2</sub>), 3.51 (d, *J* = 4 Hz, H-3), 3.88 (d, *J* = 4 Hz, H-4), 7.20 (d, *J* = 7 Hz, H-8), 7.75 (m, H-5, H-7), which was used directly in the preparation of compound **30**.

**trans-3,4-Dihydro-2,2-dimethyl-4-(2-oxopyrrolidin-1-yl)-2H-pyran[3,2-*c*]pyridin-3-ol (7).** [3,2-*c*]Epoxide **27** (0.43 g, 2.4 mmol) was added to a stirred solution of 2-pyrrolidone (0.22 mL, 2.9 mmol) in dry DMSO (10 mL) containing NaH (80 mg of 80% dispersion in oil, 2.7 mmol) under N<sub>2</sub> at room temperature. The reaction mixture was stirred for an additional 24 h. Water (100 mL) was added cautiously and the aqueous layer was extracted into EtOAc. The organic layer was washed with H<sub>2</sub>O (at pH 7) and brine and then dried over anhydrous MgSO<sub>4</sub>. Filtration and evaporation, followed by radial chromatography (silica gel, gradient elution CHCl<sub>3</sub> to 20% MeOH–CHCl<sub>3</sub>) and recrystallization, gave compound **3** (74 mg). Similarly prepared were compounds **6**, **9**, and **10** (using 2-piperidinone). See Table I.

**trans-3,4-Dihydro-2,2-dimethyl-4-(2-oxopyrrolidinyl)-2H-pyran[2,3-*c*]pyridin-3-ol *N*-Oxide (30).** Epoxide **27** (2,3-*c* *N*-oxide, 1.01 g, 5.2 mmol) was dissolved in dry ethanolic ammonia (30 mL) and maintained at room temperature for 7 days. Evaporation and trituration with EtOAc gave amino alcohol **28** (0.62 g, 56%) as a pale yellow solid. A sample recrystallized from EtOAc–MeOH had mp 206–208 °C. Anal. (C<sub>10</sub>H<sub>14</sub>N<sub>2</sub>O<sub>3</sub>) H; C, N: calcd C, 57.13, N, 13.33; found C, 56.69, N, 12.88.

To an ultrasonicated, stirred solution of amino alcohol **28** (1.54 g, 5.0 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (600 mL) and Et<sub>3</sub>N (0.64 mL, 4.6 mmol) was added chlorobutyl chloride (0.91 mL, 8.34 mmol). The ultrasonication and stirring were continued for 1 h. The solution was evaporated and the residue was chromatographed on silica gel (200 g) and eluted with a gradient of CHCl<sub>3</sub> to 10% MeOH–CHCl<sub>3</sub> to give compound **29** (1.0 g, 43%). To the (chlorobutyl)amino compound **29** (487 mg, 1.55 mmol) dissolved in dry DMSO (10 mL) and dry THF (10 mL) was added 80 NaH (50 mg, 1.7 mmol), and the mixture was stirred for 16 h at room temperature. Water was added cautiously, and the resulting solution was extracted with CHCl<sub>3</sub>. The organic layer was dried over anhydrous MgSO<sub>4</sub>, filtered, and evaporated to give a solid which was recrystallized from EtOAc–MeOH to give compound **30** (135 mg, 31%); mp 268–271 °C. Anal. (C<sub>14</sub>H<sub>18</sub>N<sub>2</sub>O<sub>4</sub>) H, N; C: calcd, 60.42; found, 59.85.

**trans-3,4-Dihydro-2,2-dimethyl-4-(2-oxopyrrolidinyl)-2H-pyran[2,3-*c*]pyridin-3-ol (8).** To a stirred solution of compound **30** (0.24 g, 0.9 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (200 mL) was added Si<sub>2</sub>Cl<sub>6</sub><sup>7</sup> (0.4 mL, 2.3 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (20 mL) dropwise under N<sub>2</sub>. The mixture was stirred for 1 h before addition of a 10% aqueous NaOH solution (0.5 mL). The mixture was stirred for 10 min and evaporated to dryness. Extraction with CHCl<sub>3</sub> and with MeOH and evaporation of the mixture gave a gum which was purified with radial chromatography (gradient elution CHCl<sub>3</sub> to 10% MeOH–CHCl<sub>3</sub>) to give title compound **8** (0.11 g) (see Table I).

**trans-3,4-Dihydro-2,2-dimethyl-4-(2-oxopiperidinyl)-2H-pyran[3,2-*c*]pyridin-3-ol *N*-Oxide (11).** A solution of compound **10** (0.1 g, 0.36 mmol) and *m*-CPBA (125 mg, 0.72 mmol) in CHCl<sub>3</sub> was heated under reflux for 1.5 h. The reaction mixture was cooled and evaporated and the residue was radially chromatographed with CHCl<sub>3</sub> to 10% MeOH–CHCl<sub>3</sub> in a gradient elution to give a gum which was recrystallized to give compound **11** (70 mg) (see Table I).

**2,2-Dimethyl-4-(2-oxopiperidinyl)-2H-pyran[3,2-*c*]pyridine (12).** Compound **10** (0.4 g, 1.5 mmol) and NaH (88 mg of 80% dispersion in oil, 2.9 mmol) were boiled in dry xylene (35 mL) under N<sub>2</sub> for 2.5 h. The reaction mixture was cooled, and H<sub>2</sub>O was added cautiously and the solution was evaporated to give a gum which was radially chromatographed with CHCl<sub>3</sub> to 25% MeOH–CHCl<sub>3</sub> in a gradient elution to give a solid which was recrystallized to give compound **12** (86 mg) (see Table I).

**trans-4-(Acetylamino)-3,4-dihydro-2,2,6-trimethyl-2H-1-benzopyran-3-ol (24).** Epoxide **35** (R = Me, 7.0 g, 0.037 mol) was dissolved in 0.88 M NH<sub>4</sub>OH solution (90 mL, 2.3 mmol) and EtOH (160 mL) and stirred at room temperature for 5 days. Evaporation gave crude amino alcohol **36** (7.0 g, 92%). To this amino alcohol (**36**; 1.0 g, 4.8 mmol) and Et<sub>3</sub>N (0.67 mL, 4.8 mmol) dissolved in CH<sub>2</sub>Cl<sub>2</sub> (60 mL) was added acetyl chloride (0.34 mL, 4.8 mmol) dropwise with stirring at room temperature. The reaction mixture was stirred for 3 h and then washed with H<sub>2</sub>O and brine and dried over anhydrous MgSO<sub>4</sub>. The solution was filtered and evaporated and the residue was radially chromatographed with a 30% pentane–EtOAc to EtOAc gradient elution, and the crude product was recrystallized to give compound **24** (0.25 g) (see Table II).

In a similar manner, **trans-4-(acetylamino)-3,4-dihydro-2,2-dimethyl-2H-pyran[3,2-*c*]pyridin-3-ol (13)** was prepared from amino alcohol **31** (see Table I).

**Pharmacological Testing. Hypertensive Rats.** All of the test compounds, and the standard drug, were evaluated for antihypertensive activity in conscious, spontaneously hypertensive rats (14–24 weeks old), derived from the Japanese (Okamoto) strain. Animals with systolic blood pressure >180 mmHg (1 mmHg = 133 Pa) were considered to be hypertensive.

Systolic blood pressure was recorded by the tail-cuff method by using a W+W blood pressure recorder, Model no. 8005; each determination was the mean of at least six recordings. Blood pressure measurements were made prior to the oral administration of test compound and at intervals for up to 6 h postdose.

All compounds were administered (via an oral dosing needle placed in the esophagus) as a solution or suspension in 1% w/v methylcellulose solution.

With the use of the above procedure, vehicle alone typically has little or no effect on blood pressure apart from a slight reduction (by 5–10%) at 6 h postdose.

**Acknowledgment.** We thank Carol Doe, Frances Hicks, and Graham Moore for their skilled and enthusiastic technical assistance.