ACS Medicinal Chemistry Letters

Letter

A Click-addressable Cassette For Photoaffinity Labeling

Bosheng Zhao, and Kevin Burgess

ACS Med. Chem. Lett., Just Accepted Manuscript • DOI: 10.1021/acsmedchemlett.7b00516 • Publication Date (Web): 24 Jan 2018 Downloaded from http://pubs.acs.org on January 25, 2018

Just Accepted

"Just Accepted" manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides "Just Accepted" as a free service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. "Just Accepted" manuscripts appear in full in PDF format accompanied by an HTML abstract. "Just Accepted" manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are accessible to all readers and citable by the Digital Object Identifier (DOI®). "Just Accepted" is an optional service offered to authors. Therefore, the "Just Accepted" Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the "Just Accepted" Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these "Just Accepted" manuscripts.



ACS Medicinal Chemistry Letters is published by the American Chemical Society. 1155 Sixteenth Street N.W., Washington, DC 20036

Published by American Chemical Society. Copyright © American Chemical Society. However, no copyright claim is made to original U.S. Government works, or works produced by employees of any Commonwealth realm Crown government in the course of their duties.

7 8

9 10

11

12

13

14

15

16

17

18

19

20 21

22 23

24

25

26

27

28

A Click-addressable Cassette For Photoaffinity Labeling

Bosheng Zhao, and Kevin Burgess*

Department of Chemistry, Texas A & M University, Box 30012, College Station, TX 77842

ABSTRACT: A small molecule 1 was designed to contain an alkyne, a trifluoromethyl-substituted diazirine, and a free piperidine-NH for facile conjugation to protein binding ligands. This "cassette" 1 was

synthesized via a relatively direct route involving only routine steps. In this proof-of-concept study, putative ligands for carbonic anhydrase IX and for TrkC were conjugated to 1. Photoaffinity labeling was performed using purified extracellular regions of both these protein-receptors, and using cells that express these receptors (isolation via a pull-down procedure); labeling of the protein was observed in all four experiments.



KEYWORDS: photoaffinity labeling, click chemistry, carbonic anhydrase IX, TrkC, targeting.

Photoaffinity labeling (PAL) is a valuable tool in target identification.¹ Preferred embodiments of this method include photolabile groups that decompose to carbenes or nitrenes, which undergo insertion reactions at reactive centers of proteins drawn into the environment by an attached ligand (Figure 1a). Difficulties with PAL tend to be encountered in detecting the proteins targeted by the reactive species; this usually requires labeling with a fluor or another entity to make the conjugates conspicuous.² However, fluors incorporated before the photodecomposition step can impact binding of the targeting ligand to the protein. Indeed, even without fluors, addition of PAL-groups to targeting ligands can impact how they bind proteins.3 For these reasons, it is often advantageous to "click" 4,5 on a fluor after photodecomposition so that fluor-protein interactions have no impact on the coupling step, but after click the covalent conjugates can be readily detected. This Letter describes a simple, conveniently accessible, "clickable cassette" 1 (Figure 1b) that may be readily conjugated to targeting ligands, and includes an alkyne for click-labeling with azides. Two applications of cassette 1 are also presented, ie labeling of carbonic anhydrase IX (CAIX) and of tropomyosin kinase receptor C (TrkC).

Several criteria were prioritized in the design of PAL-cassette 1, 44 specifically: (i) ease of synthesis; (ii) a trifluoromethyl-substituted 45 diazirine because that substituent is often preferred for PAL;⁶ (iii) a 46 highly nucleophilic secondary amine for coupling to small molecule 47 ligands; (iv) an alkyne for click-capture; and, (v) small structure 48 overall to minimize non-specific binding of the cassette fragment to 49 proteins. The presence of a tertiary amine in this system also 50 means that it could be protonated to increase water solubilities of 51 ligand-conjugates. 52

Scheme 1a describes the synthesis of cassette 1 from the known diazirine precursor **A**.⁷ Nucleophilic displacement with the amine 2 (prepared via a reductive amination of propargyl amine with the corresponding ketone, see supporting) gave the Boc-protected cassette 3; the deprotected cassette 1 was obtained through treatment with acid.

As a test application of PAL-1, this material was coupled with cyanuric chloride to give the pivotal intermediate 4 (Scheme 1b). A second S_NAr displacement on the triazine enabled incorporation of a carbonic anhydrase IX ligand in the conjugate 5.



Figure 1. a Typical PAL strategy. b Cassette featured in this work.

Carbonic anhydrase (CA) IX is a transmembrane protein, one of a family of enzymes that mediate reversible hydration of CO₂.⁸ This enzyme tends to be overexpressed in hypoxic tumors but not significantly expressed in most normal tissues.9 Small molecules that ACS Paragon Plus Environment



Scheme 1. Synthesis of: **a** cassette **1**; and, **b 5** comprising **1** conjugated with a CAIX ligand.

In a photoaffinity labeling experiment, the conjugate **5** with recombinant CAIX in pH 7.4 HEPES buffer was illuminated at 365 nm (light emitting diode) for 30 min. Fifty-fold excesses of azide-fluor-488 (compared to ligand **5**), copper (2+) salts, and a reducing agent (*tris*(2-carboxyethyl)phosphine, TCEP) were added and the reaction mixture was held at 25 °C for 3 – 4 h in the dark to allow the click cycloaddition to proceed. Constituents of the sample were separated on an SDS-PAGE gel; after fluorescence imaging, a band at 55 kDa was observed corresponding to labeled CAIX. Staining all proteins on the same gel with a Coomassie blue dye revealed the protein is present in all lanes, but comparison with the in-gel fluorescent image showed that 50-fold excess of the conjugate *without a diazirine* **6** blocked the photoaffinity labeling (lanes 2 and 4). When the sample was not illuminated (lane 5), no photoaffinity labeling occurred.



Figure 2. a *Experiments with isolated protein*. Photoaffinity labeling of recombinant CAIX after 30 min illumination with excess ligand **5** and click with azide-fluor-488; the top gel (black background) shows the fluorescence at 488 nm, and the one below depicts protein staining with Coomassie Brilliant Blue (CBB) G250. Competition with **6**, which lacks a diazirine, suppresses labeling with **5** (lanes 2 and 4). **b** *Experiments with MDA-MB-231 cell lysates*. MDA-MB-231 cell lysate was treated with PAL probe **5** under 365 nm illumination for 30 min, then clicked with biotin-azide. NeutrAvidin agarose was used to pull down the biotinylated proteins, and the material captured was run on a SDS-PAGE gel. Lane 1 shows staining of CAIX on the Western blot image with a CAIX Ab, lane 2 shows sample pretreated with blocking ligand **6**. Lane 3 shows no ligands and lane 4 depicts the lysate without PAL.

The next set of experiments were undertaken to confirm that the cassette could be used on cell lysates, rather than purified recombinant protein, and that the conjugate could be observed via biotin-labeling and a pull-down procedure (Figure 2b). "Triple negative" breast cancer cells, MDA-MB-231 were used since these are known to express CAIX.¹³ Thus, the overall procedure was photoaffinity labeling, click with biotin azide, pull-down on streptavidin coated beads, then SDS-PAGE gel and Western blot. This sequence revealed the presence of CAIX (lane 1), but not when excess blocking ligand **6** was present (lane 2) or when no photoaffinity label at all was used (lane 3). Lane 4 is a positive control showing that CAIX was present in the cell lysate before pull-down.



Figure 3. a *Experiments with solubilized TrkC extracellular domain*.
Solubilized TrkC extracellular domain was incubated with 7 for 1 h, illuminated for 30 min at 365 nm, clicked with azide-fluor-488, then subjected to SDS-PAGE. Top gel: fluorescence at 488 nm.
Bottom gel: protein staining with CBB G250. Competition with 8, which lacks a diazirine, suppresses labeling with 7 (lanes 2 and 4).
b *Experiments with NIH3T3 TrkC+ cell lysates*. Cell lysate was treated with PAL probe 7 under 365 nm illumination for 30 min, then clicked with biotin azide. NeutrAvidin agarose was used to pull down the biotinylated proteins, and the material captured was run on a SDS-PAGE gel. Lane 1 shows staining of TrkC on the Western blot image with anti-TrkC mAb, lane 2 shows sample pretreated with blocking ligand 8.

Several studies from our laboratories have featured a bivalent system with dipeptide fragments (blue in structures 7 and **8**) that putatively bind the cell surface receptor TrkC.¹⁴¹⁹ Thus conjugates 7 and **8** were formed to test this hypothesis via photoaffinity labeling.

Figure 3a shows in-gel fluorescence after a photoaffinity labeling, click, and SDS-PAGE sequence for solubilized TrkC receptor and the conjugate 7. A band at around 100 kDa was observed (lanes 1 and 3) but not when 80x of the blocking ligand **8** was added (lanes 2 and 4), or if the illumination step was excluded (lane 5). Figure 3b shows a Western blot after illumination of 7 with NIH3T3 cells stably transfected with TrkC, "click" with biotin azide, capture onto NeutrAvidin agarose beads, then SDS-PAGE. Two TrkC-Ab bands were observed (whereas the parent NIH3T3 cells give none, see supporting); these are at similar, but not the same, molecular mass as the solubilized extracellular domain of the TrkC receptor in Figure 3a.

Other PAL ligands functionalized with alkynes have been reported prior to our work,^{20,21} but the synthesis of cassette 1 appears to be more convenient than most, and provides another type of system. This is because, unlike some other probes, synthesis of cassette 1 does not require reactive organometallics, except in the first step wherein the starting material was made on scale. Cassette 1 has a secondary amine that can be easily coupled to activated carboxylic acids; this feature makes the probe more portable. Both the small-molecule conjugates prepared in this work labeled their anticipated targets with sufficient efficiencies for observation in SDS-PAGE experiments.

ASSOCIATED CONTENT

Supporting Information. The Supporting Information is available free of charge on the ACS Publications website at DOI: XXX. synthetic procedures for compound **1-8**, procedures for photoaffinity labeling, full gel images

AUTHOR INFORMATION

Corresponding Author

* FAX: +1 979 845 8839. E-mail: burgess@tamu.edu.

ACKNOWLEDGMENT

We thank DoD BCRP Breakthrough Award (BC141561), CPRIT (RP150559 and RP170144), The Robert A. Welch Foundation (A-1121). The NMR instrumentation at Texas A&M University was supported by a grant from the National Science Foundation (DBI-9970232) and the Texas A&M University System.

REFERENCES

- (1) Sumranjit, J.; Chung, S. J. Recent advances in target characterization and identification by photoaffinity probes. *Molecules* **2013**, *18*, 10425-10451.
- (2) Lapinsky, D. J. Tandem photoaffinity labeling-bioorthogonal conjugation in medicinal chemistry. *Bioorg. Med. Chem.* 2012, 20, 6237-6247.
- (3) Park, J.; Koh, M.; Koo, J. Y.; Lee, S.; Park, S. B. Investigation of Specific Binding Proteins to Photoaffinity Linkers for Efficient Deconvolution of Target Protein. ACS Chem. Biol. 2016, 11, 44-52.
- (4) Meldal, M.; Tornoe, C. W. Cu-Catalyzed Azide-Alkyne Cycloaddition. *Chem. Rev.* 2008, 108, 2952-3015.
- (5) Kolb, H. C.; Finn, M. G.; Sharpless, K. B. Click Chemistry: Diverse Chemical Function from a Few Good Reactions. *Angew. Chem. Int. Ed.* 2001, 40, 2004-2021.
- (6) Dubinsky, L.; Krom, B. P.; Meijler, M. M. Diazirine based photoaffinity labeling. *Bioorg. Med. Chem.* 2012, 20, 554-570.
- (7) Stromgaard, K.; Saito, D. R.; Shindou, H.; Ishii, S.; Shimizu, T.; Nakanishi, K. Ginkgolide Derivatives for Photolabeling Studies:

Preparation and Pharmacological Evaluation. J. Med. Chem. 2002, 45, 4038-4046.

- (8) Krishnamurthy, V. M.; Kaufman, G. K.; Urbach, A. R.; Gitlin, I.; Gudiksen, K. L.; Weibel, D. B.; Whitesides, G. M. Carbonic Anhydrase as a Model for Biophysical and Physical-Organic Studies of Proteins and Protein-Ligand Binding. *Chem. Rev.* 2008, 108, 946-1051.
- (9) Swietach, P.; Vaughan-Jones, R. D.; Harris, A. L. Regulation of tumor pH and the role of carbonic anhydrase 9. *Cancer and Metastasis Reviews* 2007, 26, 299-310.
- (10) Thiry, A.; Dogne, J.-M.; Masereel, B.; Supuran, C. T. Targeting tumorassociated carbonic anhydrase IX in cancer therapy. *Trends Pharmacol. Sci.* 2006, 27, 566-573.
- (11) Krall, N.; Pretto, F.; Neri, D. A bivalent small molecule-drug conjugate directed against carbonic anhydrase IX can elicit complete tumour regression in mice. *Chem. Sci.* **2014**, 5, 3640-3644.
- (12) Krall, N.; Pretto, F.; Decurtins, W.; Bernardes, G. J. L.; Supuran, C. T.; Neri, D. A small-molecule drug conjugate for the treatment of carbonic anhydrase IX expressing tumors. *Angew. Chem., Int. Ed.* **2014**, 53, 4231-4235.
- (13) Li, Y.; Wang, H.; Oosterwijk, E.; Tu, C.; Shiverick, K. T.; Silverman, D. N.; Frost, S. C. Expression and Activity of Carbonic Anhydrase IX Is Associated With Metabolic Dysfunction in MDA-MB-231 Breast Cancer Cells. *Cancer Invest.* 2009, *27*, 613-623.
- (14) Chen, D.; Brahimi, F.; Angell, Y.; Li, Y.-C.; Moscowicz, J.; Saragovi, H. U.; Burgess, K. Bivalent Peptidomimetic Ligands of TrkC are Biased Agonists, Selectively Induce Neuritogenesis, or Potentiate Neurotrophin-3 Trophic Signals. ACS Chem. Biol. 2009, 4, 769-781.
- (15) Ko, E.; Kamkaew, A.; Burgess, K. Small Molecule Ligands for Active Targeting of TrkC-expressing Tumor Cells. ACS Med. Chem. Lett. 2012, 3, 1008-1012.

- (16) Kamkaew, A.; Burgess, K. Double-targeting Using a TrkC-Ligand Conjugated to Dipyrrometheneboron Difluoride (BODIPY) Based Photodynamic Therapy (PDT) Agent. J. Med. Chem. 2013, 56, 7608-7614.
- (17) Brahimi, F.; Ko, E.; Malakhov, A.; Burgess, K.; Saragovi, H. U. Combinatorial assembly of small molecules into bivalent antagonists of TrkC or TrkA receptors. *PLoS One* **2014**, *9*, e89617/89611e89617/89612.
- (18) Kue, S. C., Anyanee Kamkaew, Hong Boon Lee, Lip Long Chung, Lik Voon Kiew, and Kevin Burgess. Targeted PDT Agent Eradicates TrkC Expressing Tumors Via Photodynamic Therapy (PDT). *Mol. Pharmacetics* 2015, 12, 212-222.
- (19) Kamkaew, A.; Li, F.; Li, Z.; Burgess, K. An agent for optical imaging of TrkC-expressing, breast cancer. *MedChemComm* **2017**, *8*, 1946-1952.
- (20) Kumar, N. S.; Young, R. N. Design and synthesis of an all-in-one 3-(1,1difluoroprop-2-ynyl)-3H-diazirin-3-yl functional group for photo-affinity labeling. *Bioorg. Med. Chem.* **2009**, *17*, 5388-5395.
- (21) Li, Z.; Hao, P.; Li, L.; Tan, C. Y. J.; Cheng, X.; Chen, G. Y. J.; Sze, S. K.; Shen, H.-M.; Yao, S. Q. Design and Synthesis of Minimalist Terminal Alkyne-Containing Diazirine Photo-Crosslinkers and Their Incorporation into Kinase Inhibitors for Cell- and Tissue-Based Proteome Profiling. *Angew. Chem., Int. Ed.* **2013**, *52*, 8551-8556.

60