

Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information:
<http://www.tandfonline.com/loi/gpss20>

Synthesis, Spectral Studies and C-S, C-N Bond Fissions of Some Novel N-[Alkyl (4'- Substituted Phenylthio)] Phthalimides

Ali A. El-Bardan

^a Chemistry Department, Faculty of Science, Alexandria University, P.O. 426 Ibrahimia, Alexandria 21321, Egypt
Published online: 17 Mar 2008.

To cite this article: Ali A. El-Bardan (1998) Synthesis, Spectral Studies and C-S, C-N Bond Fissions of Some Novel N-[Alkyl (4'- Substituted Phenylthio)] Phthalimides, *Phosphorus, Sulfur, and Silicon and the Related Elements*, 134:1, 511-519, DOI: [10.1080/10426509808545492](https://doi.org/10.1080/10426509808545492)

To link to this article: <http://dx.doi.org/10.1080/10426509808545492>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the

Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

SYNTHESIS, SPECTRAL STUDIES AND C-S, C-N BOND FISSIONS OF SOME NOVEL N-[ALKYL (4'- SUBSTITUTED PHENYLTHIO)] PHTHALIMIDES*

ALI A. EL-BARDAN

*Chemistry Department, Faculty of Science, Alexandria University, P.O. 426 Ibrahimia,
Alexandria 21321, Egypt*

(Received 9 January, 1998; In final form 30 April, 1998)

N-[(4'-Substituted phenylthio)ethyl] phthalimide 1a-g and their corresponding n-propyl 2a-g, n-butyl 3a-g derivatives have been synthesised. The structure of these compounds were proved by UV, IR, ^1H NMR, ^{13}C MR and mass spectra. The ^1H and ^{13}C chemical shifts of the adjacent methylene protons and carbons of both the sulfur atom and phthalimido nitrogen group were reasonably correlated using σ^p values. The carbon-nitrogen and the carbon-sulfur bond fissions in alkaline and acidic media were discussed.

Keywords: Arylthio alkyl phthalimides; ^1H NMR; ^{13}C MR; mass spectra; C-S and C-N bond fissions

INTRODUCTION

Previously we have described the synthesis of different series of diaryl, aryl benzyl, aryl picryl, aryl heterocyclic sulfides, sulfoxides and sulfones as well as aryl mercapto acetates and propionates¹⁻⁸. The mechanisms of the alkaline hydrolysis of the C-S bonds of some of these compounds were reported^{9,10}. Furthermore, the effect of various substituents of some of these sulfides and sulfones on toxicity¹¹, chemical shifts of methylene and aromatic protons¹²⁻¹⁴, mass spectra¹⁵, pK_a ¹⁶ and nucleophilic substitutions¹⁷⁻¹⁹ were examined. The previous work has generated an interest in synthesizing new derivatives of the basic

* This paper was presented at the 6th Ibn Sina International Conference on pure and Applied Heterocyclic Chemistry, December 13-16, 1997, Cairo, Egypt.

structure involving the N-alkyl phthalimido nucleus for chemical and biological studies. These compounds are N-[2-(4'-substituted phenylthio)ethyl] phthalimide 1a-g, N-[3-(4'-substituted phenylthio)propyl] phthalimide 2a-g and N-[4-(4'-substituted phenylthio)butyl] phthalimide 3a-g.

Generally compounds containing the $\text{CH}_2\text{-S}$ moiety show low acidity of the α -hydrogen under basic conditions, while those containing more than one sulfide group enhances it²⁰. This could be attributed to the formation of a carbanion intermediate which is stabilized by the 3d-orbital resonance.

RESULTS AND DISCUSSION

The structure of the title compounds were confirmed by UV, ^1H NMR and ^{13}CMR , Tables I-III. The IR spectra²¹⁻²³ (KBr pellets) for all compounds showed a sharp band at $1690\text{--}1705\text{ cm}^{-1}$ (C=O) and at $716\text{--}723\text{ cm}^{-1}$ (C-S). The mass spectra^{23,24} of compounds 2a,b,e were studied. The assignments of the major fragment ion peaks of these compounds are compiled in Table IV and Scheme 1. The major role of the aliphatic n-propyl moiety is evident in directing the electron impact mass spectroscopic fragmentation pathways and this favors the formation of two successive ion series. The first includes the phthalimido ion peak at m/z 146 as well as the other ions at m/z 160, 174, 188 and 220. The second series of ions is characterized by the presence of the aryl part of the compound at m/z (M-220), (M-188), (M-174), (M-160) and (M-146) ions. The general behavior in the mass spectra of compounds 2a,b,e is characterized by the base peak formation in all compounds at m/z 160 which is formed via β -bond cleavage to the nitrogen atom giving rise to the formation of $[\text{phthalimido-CH}_2]^+$ ion, which is more favourable due to charge stabilization by resonance with the nitrogen atom.

Plots of the ^1H and ^{13}CMR δ ppm chemical shifts of the adjacent methylene protons and carbons to both sulfur and nitrogen atoms in all series against σ° constants²⁵ give good straight lines, Table V. The diminution in the transmission inductive effects of N-CH_2 relative to S-CH_2 is attributed to the proximity effect of the latter.

TABLE I Properties, UV, ^1H NMR and ^{13}C NMR of N-[2-(4'-substituted phenylthio)ethyl] phthalimide 1a-g

Cpd. I	R	MP °C	λ^a nm	ϵ/dm^3 mol^{-1} cm^{-1}	S % Found	^1H NMR (^{13}C NMR) δ ppm in CDCl_3							
						S-CH ₂ (t, 2H)	N-CH ₂ (t, 2H)	C _{3',6'} (d, 2H)	C _{4',5'} (t, 2H)	C _{2',6'} (d, 2H)	C _{3,5'} (d, 2H)	C _{4'} (t, 1H)	C=O
a ^b	H	138	224.5	17170	10.98	3.45 (32.18)	3.58 40.09	7.88 134.11	7.58 129.19	7.46 131.03	7.33 128.58	7.26 125.5	---
b ^c	CH ₃	145	252.5	5860	10.52	3.06 (31.78)	3.53 39.97	7.80 134.37	7.58 129.41	7.36 131.28	7.18 127.56	---	---
c ^d	OCH ₃	132	255	9130	9.92	3.00 (31.68)	3.50 39.77	7.82 133.78	7.54 130.06	7.40 132.28	6.82 128.54	---	---
d ^e	Cl	95	253	1207	9.89	3.33 (32.78)	3.69 40.50	7.42 134.93	7.28 129.71	7.82 131.56	7.82 123.64	---	---
e	Br	140	256	14410	8.56	3.29 (32.68)	3.65 40.42	7.85 132.68	7.55 128.47	7.39 130.08	7.47 131.06	---	---
f	NO ₂	211	258	16160	9.56	3.68 (33.80)	3.84 41.02	7.92 134.91	7.54 128.26	7.58 131.28	8.16 132.56	---	---
g ^f	COOH	298	---	---	9.62	3.28 (32.26)	3.52 39.87	8.04 133.26	7.56 128.15	7.64 131.61	7.34 125.99	---	169.65

^a In methanol. ^b C_{3,5'} (t, 2H); ¹³C NMR in DMSO, 4'-CH₃ appear at δ_{H} 2.23 (s, 3H); δ_{C} 20.56 ppm. ^c 4'-OCH₃ appear at δ_{H} 3.78 (s, 3H); δ_{C} 55.51 ppm. ^d C_{2,3,5,6'} (s, 4H). ^e 2'-COOH; in acetone-d₆; OH appear at δ_{H} 11.12 (s, 1H); C_{4,5'} (t, 2H).

TABLE II Properties, UV, ^1H NMR and ^{13}C NMR of N-[3-(4'-substituted phenylthio) propyl] phthalimide 2a-g

Cpd.	R	MP °C	λ^a nm	$\epsilon/\text{dm}^3 \text{ mol}^{-1} \text{ cm}^{-1}$	S % Found	^1H NMR (^{13}C NMR) δ ppm in CDCl_3									
						2 CH_2 (q, 2H)	S- CH_2 (t, 2H)	N- CH_2 (t, 2H)	$\text{C}_3, 6''$ (d, 2H)	$\text{C}_4, 5''$ (t, 2H)	$\text{C}_2, 6'$ (d, 2H)	$\text{C}_3, 5'$ (d, 2H)	C_4' (t, 1H)	$\text{C}=\text{O}$	
a	H	88	225	18170	10.26	1.92	2.85	3.68	7.75	7.45	7.00-7.23 (m)		126.5	169.26	
			253	8660		(27.62)	30.67	40.68	134.58	129.29	131.24	128.32	126.5	169.26	
b ^b	CH_3	81	225.5	18870	9.89	1.91	2.75	3.64	7.77	7.53	7.20	6.95	--	--	
			256	9520		(27.26)	30.18	40.61	133.92	128.96	131.18	127.12	125.23	169.12	
c ^c	OCH_3	60	228.5	21570	9.58	1.91	2.70	3.61	7.72	7.46	7.24	6.71	--	--	
			254	12170		(26.68)	30.09	40.48	134.78	129.82	131.28	128.23	117.12	167.92	
d ^d	Cl	90	227	19330	9.49	1.93	2.98	3.77	7.75	7.55	7.10	7.10	--	--	
			261	14420		(27.28)	31.20	41.02	133.62	129.62	132.54	125.92	127.31	167.12	
e	Br	103	228.5	19860	8.11	1.95	2.93	3.73	7.75	7.52	7.10	7.30	--	--	
			262	16610		(27.65)	31.08	40.96	133.98	128.56	129.86	131.32	127.85	169.64	
f	NO_2	212	229	19260	9.12	1.96	3.18	3.92	7.98	7.52	7.62	8.10	--	--	
			295	15660		(28.12)	32.52	44.50	134.98	129.10	131.28	132.78	128.53	169.58	
g ^e	COOH	170	--	--	9.18	1.92	2.91	3.81	7.98	7.54	7.76	7.32	--	--	
			--	--		(27.21)	33.12	38.78	134.98	128.38	131.68	127.12	129.52	169.12	

^a In methanol. ^b 4'- CH_3 appear at δ_{H} 2.39 (s, 3H); δ_{C} 20.45 ppm. ^c 4'- OCH_3 appear at δ_{H} 3.66 (s, 3H); δ_{C} 55.45 ppm.^d $\text{C}_{2,3,5,6}$ (s, 4H). ^e 2'- COOH ; in acetone - d_6 ; OH appear at δ_{H} 11.21 (s, 1H); $\text{C}_{4,5'}$ (t, 2H).


TABLE III Properties, UV, ^1H NMR and ^{13}C NMR of N-[4-(4'-substituted phenylthio) butyl] phthalimide 3a-g

Cpd.	R	MP °C	λ^a nm	ϵ/dm^2 mol^{-1} cm^{-1}	S % Found	^1H NMR (^{13}C NMR) δ ppm in CDCl_3									
						2 $\text{C}_{12}(m,$ 4H)	S - $\text{CH}_2(t,$ 2H)	N - $\text{CH}_2(t,$ 2H)	$\text{C}_{3,6}(d,$ 2H)	$\text{C}_{4,5}(t,$ 2H)	$\text{C}_{2,6}(d,$ 2H)	$\text{C}_{3,5}(d,$ 2H)	$\text{C}_4(t,$ 1H)	C = O	
^b _a	H	50	226.5	17770	10.0	1.72	2.90	3.65	7.75	7.50	7.21	7.12	6.95	---	
			255	9830		(21.6	31.22	40.50	134.68	129.98	132.12	128.26	124.5	167.25)	
^b _c	CH_3	75	227.5	19910	9.58	1.68	2.82	3.62	7.69	7.48	7.12	6.89	--	---	
			256.5	10940		(22.12	30.99	40.41	133.89	129.24	131.21	128.12	123.12	169.10)	
^c _d	OCH_3	67	229	20670	9.21	1.70	2.78	3.59	7.75	7.52	7.20	7.68	--	---	
			257	11220		(21.89	30.51	40.25	134.98	129.32	132.68	127.15	118.2	169.12)	
^d _e	Cl	80	226	14990	9.10	1.74	3.07	3.74	7.78	7.56	7.10	7.10	--	---	
			261.5	7270		(22.12	31.92	40.78	133.68	129.72	132.12	128.15	125.48	167.12)	
^e	Br	95	227.5	19040	8.00	1.75	3.03	3.72	7.76	7.58	7.06	7.27	--	---	
			262.5	16000		(21.26	31.74	40.70	134.68	128.68	130.70	131.02	125.69	169.12)	
^f	NO_2	158	229.5	19800	8.81	1.75	3.30	3.88	7.85	7.40	7.58	8.07	--	---	
			293.5	16820		(22.21	33.12	41.28	134.51	128.62	131.32	132.68	127.11	169.52)	
^g _f	COOH	136	--	--	9.00	1.69	2.88	3.68	7.95	7.48	7.72	7.32	--	---	
			--	--		(21.62	32.68	38.78	134.12	128.52	131.26	126.26	127.10	170.10)	

^a In methanol. ^b $\text{C}_{3,5}$ (t, 2H). ^c 4-CH_3 appear at δ_{H} 2.22 (s, 3H); δ_{C} 20.12 ppm. ^d 4-OCH_3 appear at δ_{H} 3.70 (s, 3H); δ_{C} 55.41 ppm. ^e $\text{C}_{2,3,5,6}$ (s, 4H). ^f 2-COOH ; in acetone- d_6 ; OH appear at δ_{H} 10.98 (s, 1H); $\text{C}_{4,5}$ (t, 2H).

TABLE IV Relative intensity of prominent peaks in the mass spectra of N-[3-(phenylthio) propyl] phthalimide 2a, N-[3-(4'-methylphenylthio) propyl] phthalimide 2b, N-[3-(4'-bromophenylthio) propyl] phthalimide 2c

Major peaks	2a		2b		2c	
	<i>m/z</i>	rel.int. %	<i>m/z</i>	rel.int. %	<i>m/z</i>	rel.int %
M ⁺	297	37	311	25	375	1
M-220	77	46	91	53	155	< 1
220	220	< 1	220	< 1	220	< 1
M-188	109	42	123	85	187	8
188	188	95	188	58	188	13
M-174	123	9	137	< 1	201	3
174	174	4	174	< 1	174	< 1
M-160	137	4	151	< 1	215	< 1
160	160	100	160	100	160	100
M-146	151	<1	165	< 1	229	< 1
146	146	5	146	< 1	146	3



2a; R=H ($M^+=297$), 2b; R=CH₃ ($M^+=311$), 2c; R=Br ($M^+=375$).


SCHEME 1

TABLE V Gradients of the correlated chemical shifts against σ^o constants

Cpd.	¹ H NMR		¹³ CMR	
	N-CH ₂	S-CH ₂	N-CH ₂	S-CH ₂
1a-f	0.321 (r=0.997)	0.632 (r=0.993)	1.158 (r=0.996)	2.030 (r=0.995)
2a-f	0.287 (r=0.989)	0.447 (r=0.992)	0.953 (r=0.996)	2.440 (r=0.977)
3a-f	0.271 (r=0.997)	0.497 (r=0.999)	0.932 (r=0.991)	2.341 (r=0.991)

The alkaline hydrolysis of compounds 1a-f, 2a-f and 3a-f with 5% sodium hydroxide solution in dioxane gave a mixture of phthalic acid, diaryl disulfides¹⁰ and alkylamines, Scheme 2. The gaseous ethylamine which has been generated during the alkaline hydrolysis reflux of compounds 1a-f was trapped with hydrochloric acid and identified as ethylamine hydrochloride²⁶. n-Propylamine hydrochloride and n-butylamine hydrochloride obtained from the acidified filtrate of 2a-f and 3a-f respectively were identified by benzooylation process^{27,28}. The acid hydrolysis of compounds 1a, 2a and 3a gave a mixture of phthalic acid, diphenyl disulfides¹⁰ and hydroxylamines²⁹, Scheme 2. In the alkaline hydrolysis the C-S bond fission could be rationalized by the attack of the OH ion on the sulfur atom to give sodium aryl sulfene¹⁰, rather than the α -carbanion formation²⁰, while the remaining moiety gives phthalate anion and alkyl amine, Scheme 2.

In the acid hydrolysis, the compounds are first protonated followed by a) hydrolysis to give phthalic acid. b) attack of H₂O on the α -carbon to give hydroxylamine and thiophenol. The latter undergoes air oxidation to give diphenyl disulfide, Scheme 2.



SCHEME 2

EXPERIMENTAL

Melting points are uncorrected, ¹H NMR and ¹³CMR in acetone-d₆ or CDCl₃ were recorded using varian XL 300 spectrometer with Me₄Si as internal standard, IR spectra (KBr pellets) were measured on a Perkin-Elmer 1430 ratio

recording infrared, UV spectra were recorded on a Shimadzu 160-A spectrophotometer and the mass spectra on MS 5988 spectrometer. Elemental analysis and mass spectra were carried out in the Faculty of Science, Cairo University, Egypt. The found analysis for carbon, hydrogen, nitrogen and sulfur were in excellent agreement with those calculated. Analytical Thin Layer Chromatography (TLC) was carried out on silica gel using (9.5:0.5) petroleum ether: ethylacetate, respectively.

General procedure for the synthesis of compounds 1, 2, 3a-g

N-(2-Bromoethyl)phthalimide, N-(3-bromopropyl)phthalimide and N-(4-bromobutyl)phthalimide (0.1 mole) in ethanol and the sodium salt of thiol compounds (Aldrich chemical Co. Inc) (0.1 mole) in ethanol were refluxed for 1–2h. Cooling, dilution and acidification with diluted hydrochloric acid gave a solid which was purified by recrystallization from aqueous ethanol.

Action of 5% sodium hydroxide solution

N-[Alkyl(4'-substituted phenylthio)]phthalimide 1a-g, 2a-f and 3a-f (1g) dissolved in dioxane and (10ml) of 5% sodium hydroxide solution were refluxed for 1h. On cooling and acidification a charactersitic thiol odor was detected. The reaction mixture was filtered. TLC of the precipitate indicated the presence of two compounds in all series which were separated by column chromatography. The first separated compound was identified as 4,4'-disubstituted diphenyl disulfide¹⁰ and the second compound in the mixture as phthalic acid (m.p. 210–11°C). The filtrate was vigorously shaken for 30 minutes with a mixture of 2 ml benzoyl chloride and 5 ml of 10% sodium hydroxide solution, then diluted with cold water. Series 2a-f and 3a-f gave a precipitate which was identified as N-(n-propyl)benzamide (m.p. 84°C, lit.²⁷ 83°C) and N-(n-butyl)benzamide (m.p.42°C, lit²⁸. 42°C). The gaseous ethylamine generated during the reflux of series 1a-f was trapped by hydrochloric acid and identified as ethylamine hydrochloride (m.p.78°C, lit²⁶.76–78°C). All of these compounds were confirmed by comparing their melting points and ¹H NMR spectra with a commercial specimen.

Action of 10% aqueous hydrochloric acid

N-[Alkyl(phenylthio)]phthalimide 1a, 2a and 3a (1g) dissolved in dioxane and (10 ml) of 10% aqueous hydrochloric acid were refluxed for 1h. On cooling and

dilution a charactersitic thiophenol odor was detected. The reaction mixture was filtered and the precipitate was chromatographically separated to phthalic acid (m.p. 210°C) and diphenyl disulfide (m.p. 61°C, lit¹⁰. 60°C). The filtrate was treated with aqueous NaHCO₃ and extracted with ether. Evaporation of the ether layer gave an oil in all series (n_D^{20} =1.46) which was identified by ¹H NMR as ethanolamine²⁹, 3-amino-1-propanol²⁹ and 4-amino-1-butanol²⁹, respectively.

References

- [1] A. A. Kassem and A. A. El-Bardan, *J. Chem. Eng. Data*, **31**, 496 (1986).
- [2] A. A. Kassem and A. A. El-Bardan and S. M. Mansour, *J. Chem. Eng. Data*, **32**, 483 (1987).
- [3] A. A. El-Bardan, E. A. Hamed and E. F. Saad, *J. Chem. Eng. Data*, **34**, 133 (1989).
- [4] A. A. El-Bardan, E. F. Saad, E. A. Hamed and A. A. Kassem, *Alex. J. Pharm. Sci.*, **4**, 16 (1990).
- [5] E. A. Hamed, A. A. El-Bardan and A. M. Moussa, *Phosphorus, Sulfur and Silicon*, **62**, 269 (1991).
- [6] A. A. El-Bardan, G. A. Gohar, F. M. El-Hegazy and E. A. Hamed, *Phosphorus, Sulfur and Silicon*, **69**, 148 (1992).
- [7] E. A. Hamed, A. El-Faham, E. F. Saad and A. A. El-Bardan, *Phosphorus, Sulfur and Silicon*, **11**, 19 (1996).
- [8] A. A. El-Bardan, E. A. Hamed, S. Abo El-Ella and A. M. Moussa, *Ind. J. Chem. Soc.* **74** (7), 575 (1997).
- [9] A. A. El-Bardan, *Phosphorus, Sulfur and Silicon*, **66**, 153 (1992).
- [10] F. M. El-Hegazy, A. A. El-Bardan and E. A. Hamed, *Phosphorus, Sulfur and Silicon*, **113**, 88 (1994).
- [11] Y. Riad, H. M. El-Nahas, E. El-Kady and A. A. El-Bardan, *J. Econ. Entomol.*, **85**, 2096 (1992).
- [12] E. F. Saad, A. A. El-Bardan and E. A. Hamed, *Spect. Lett.* **23**, 1347 (1990).
- [13] E. F. Saad, E. A. Hamed and A. A. El-Bardan, *Spect. Lett.* **24**, 413 (1991).
- [14] E. A. Hamed, A. A. El-Bardan, G. A. Gohar and A. N. Assad, *Spect. Lett.* **29**, (4) 549 (1996).
- [15] F. M. El-Hegazy, M. E. Mahmoud, E. F. Saad and E. A. Hamed, *Rapid Commun. Mass Spectrometry*, **11**, 316 (1997).
- [16] A. A. El-Bardan, N. M. El-Mallah and E. A. Hamed, *J. Phys. Org. Chem.*, **5**, 239 (1992).
- [17] N. M. El-Mallah, E. A. Hamed and A. A. El-Bardan, *Egypt. J. Chem.*, **3**, 319 (1995).
- [18] E. A. Hamed, A. A. El-Bardan and N. M. El-Mallah, *Int. J. Chem. Kinet.*, **28**, 283 (1996).
- [19] E. A. Hamed, A. A. El-Bardan, E. F. Saad, G. A. Gohar and G. M. Hassen, *J. Chem. Soc. Perkin Trans. 2*, 2415 (1997).
- [20] S. Oae, "Organic Sulfur Chemistry: Structure and Mechanism", CRC Press, New York (1991).
- [21] L. J. Ballemly, "The Infra Red Spectra of Complex Molecules", Wiley, New York (1978).
- [22] V. M. Parikh, "Absorption of Organic Molecules", Wiley, New York (1974).
- [23] R. M. Silverstein, G. C. Bassler and T. C. Morill, "Spectrometric Identification of Organic Compounds", 4th ed., Wiley, New York (1981); Ibid, 5th ed. (1991).
- [24] F. M. McLafferty, "Interpretation of Mass Spectra", Benjamin, New York (1966).
- [25] O. Exner, "In Correlation Analysis in Chemistry" (N.B. Chapman and J. Shorter, Eds) Plenum, London, Ch. 10 (1978).
- [26] G. W. Watt, J. B. Otto, *J. Am. Chem. Soc.*, **69**, 836 (1947).
- [27] A. W. Titherley, *J. Chem. Soc.*, **79**, 405 (1901).
- [28] G. H. Coleman, H. P. Howells, *J. Am. Chem. Soc.*, **45**, 3084 (1923).
- [29] L. Henry, *Ber.*, **33**, 3169 (1900).