

Synthesis of 3-Substituted-4*H*-1,4-Benzoxazines *via* Palladium-Catalysed Coupling Reactions

C. Buon, P. Bouyssou, G. Coudert*

Institut de Chimie Organique et Analytique associé au CNRS, Université d'Orléans, BP 6759
45067 Orléans Cedex 02, France

Received 23 October 1998; accepted 13 November 1998

Abstract : Here is described a high yield synthesis of 3-substituted 4*H*-1,4-benzoxazines *via* palladium-catalysed coupling reactions between organostannanes and a vinylphosphate obtained from a benzoxazin-3-one derivative.


© 1999 Published by Elsevier Science Ltd. All rights reserved.

Keywords: benzoxazines; coupling reactions; palladium and compounds; tin and compounds

In the course of a research program directed toward the design of biologically active derivatives containing a 1,4-benzoxazine subunit, we needed to have easy access to substituted 4*H*-1,4-benzoxazines. We recently reported a quite general procedure allowing the synthesis of 3-substituted-4*H*-1,4-benzoxazine *via* the regioselective lithiation-electrophilic substitution of the 4-Boc-4*H*-1,4-benzoxazine.¹ However the key step in the preparation of this unsaturated carbamic derivative from the corresponding 2,3-dihydrobenzoxazine involves a bromation-debromation sequence that is not easily achievable on a large scale.

We decided to search for a new alternative and complementary method allowing an efficient and rapid synthesis of 1,4-benzoxazines, particularly derivatives bearing aryl, heteroaryl and vinyl groups at C₃. A few years ago Kocienski described the synthesis in good yields of 6-(trimethylstannyl)3,4-dihydro-2*H*-pyrans from δ -valerolactones *via* the Pd(0) catalysed coupling of enol triflate derivatives with hexadimethylstannane.² Recently Hiemstra described the synthesis of lactam-derived enol triflates and their functionalization *via* catalytic coupling reaction with Pd(0).³

We postulated that 3-substituted-4*H*-1,4-benzoxazines could be obtained in a similar way from the benzoxazin-3-one **2** *via* the enol triflate **3** (Scheme 1).



The lactam **1** was first prepared from 2-aminophenol⁴ in 80 % yield; the N-Boc derivative **2** was then obtained in 96 % yield by treatment of **1** with di-*tert*-butyldicarbonate in dry tetrahydrofuran at room temperature in the presence of DMAP. The N-Boc benzoxazinone was treated with LDA (1.2 eq.) and TMEDA (1.2eq.) at -78°C in dry tetrahydrofuran. The corresponding lithium enolate was reacted then with

N-phenyltrifluoromethanesulfonimide which probably afforded the required enol triflate; unfortunately this compound decomposed when the temperature exceeded -50°C and could not be used further. The lack of stability of cyclic ketene acetal triflates was previously mentioned by Nicolaou who prepared and advantageously used cyclic ketene phosphates.⁵ Likewise, we chose to prepare the diphenylphosphate **4** from **2** via its lithium enolate (LDA, 1.2 eq.; TMEDA, 1.2 eq.; $(\text{PhO})_2\text{P}(\text{O})\text{Cl}$, 3eq.; THF; -78°C); the phosphate obtained in 86 % yield proved to be stable enough at room temperature and could be purified by flash chromatography on silicagel.

Coupling the intermediate **4** with various vinyl, aryl and heteroaryl stannanes (2 eq.), in the presence of LiCl (3 eq.) and a catalytic amount of $\text{Pd}(\text{PPh}_3)_4$ (0.05 eq) in THF at reflux resulted in the formation of 3-substituted derivatives **5** (Table 1).




Table 1. Palladium-Catalysed Coupling of vinyl phosphate **4** with organostannanes $\text{R}'\text{SnBu}_3$

5	$\text{R}'\text{SnBu}_3$	Product	Yield
5a	$\text{Bu}_3\text{Sn}-\text{CH}_2\text{CH}=\text{CH}_2$		87 %
5b	$\text{Bu}_3\text{Sn}-\text{CH}(\text{OEt})\text{CH}=\text{CH}_2$		82 %
5c	$\text{Bu}_3\text{Sn}-\text{C}_4\text{H}_3\text{S}$		91 %
5d	$\text{Bu}_3\text{Sn}-\text{C}_4\text{H}_3\text{O}$		96 %
5e	$\text{Bu}_3\text{Sn}-\text{C}_6\text{H}_5$		84 %
5f	$\text{Bu}_3\text{Sn}-\text{C}_{12}\text{H}_8\text{O}_2$		93 %

In summary we have developed a new versatile method for the preparation of the little known 3-substituted-4H-1,4-benzoxazines.

References

1. L. Chacun-Lefevre, C. Buon, P. Bouyssou, G. Coudert. *Tetrahedron Lett.* **1998**, 39, 5763-5764
2. C. Barber, K. Jarowicki, P. Kocienski. *Synlett.* **1991**, 197-198
3. T. Luker, H. Hiemstra, W. N. Speckamp. *J. Org. Chem.* **1997**, 62, 8131-8140
4. D. R. Shridar, M. Jogibhukta, V. S. H. Krishnan. *Org. Prep. Proced. Int.* **1982**, 14, 195-197
5. K. C. Nicolaou, G.-Q. Shi, J. L. Gunzner, P. Gärtner, Z. Yang. *J. Am. Chem. Soc.* **1997**, 119, 5467-5468