


Cycloaddition Reaction of 2,3-Disubstituted Oxiranes with Isocyanates  
by Highly Activated Catalyst;  $\text{Ph}_4\text{SbI}-\text{Bu}_3\text{SnI}$

Masahiro FUJIWARA, Akio BABA,\* Yuri TOMOHISA, and Haruo MATSUDA  
Department of Applied Chemistry, Faculty of Engineering,  
Osaka University, 2-1 Yamadaoka, Suita, Osaka 565

The combination of  $\text{Ph}_4\text{SbI}$  and  $\text{Bu}_3\text{SnI}$  proved to be a new excellent catalyst for effecting cycloaddition reaction of inactive 2,3-disubstituted oxiranes with isocyanates under neutral conditions, yielding oxazolidinones in high yields, while either of both metal iodides had no activity individually.

The cycloaddition of oxiranes toward isocyanates has been ardently studied for a convenient preparation of oxazolidinones, and a variety of catalysts have been reported.<sup>1)</sup> Quite recently we also reported that organotin halide-base complexes and organostibonium halides are very effective catalysts for this reaction.<sup>2)</sup> These compounds show high and unique catalytic activity in the reaction of monosubstituted oxiranes under mild conditions. Unfortunately, even these active catalysts have lower effect for the cycloaddition using 2,3-disubstituted oxiranes. The development of a versatile catalyst for reactions of these inactive oxiranes is a significant problem, and if realized, a wide range of oxazolidinones could be readily obtained from the direct cycloaddition with isocyanates.



Although showing not high activity toward this type of reaction,  $\text{Ph}_4\text{SbI}$ , as

Table 1. Reaction of 2,3-Disubstituted Oxiranes with Isocyanates<sup>a)</sup>

Entry	Oxirane 1		Isocyanate 2	Solvent	Cat. system	Yield of 3 / %
	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>			
1	-(CH <sub>2</sub> ) <sub>4</sub> -		Ph	PhH	Ph <sub>4</sub> SbI - Bu <sub>3</sub> SnI	99
2	"		"	"	Ph <sub>4</sub> SbI	15
3	"		"	"	Bu <sub>3</sub> SnI	0
4	"		"	"	Bu <sub>4</sub> NI	0 <sup>b)</sup>
5	"		"	"	Ph <sub>4</sub> SbI - Bu <sub>2</sub> SnI <sub>2</sub>	87
6	"		"	"	Ph <sub>4</sub> SbI - SnI <sub>2</sub>	48
7	"		"	"	Ph <sub>4</sub> SbI - ZnI <sub>2</sub>	35
8	"		"	"	Ph <sub>4</sub> SbI - AlI <sub>3</sub>	13
9	"		"	"	Ph <sub>4</sub> SbI - CuI	0
10	"		"	CH <sub>3</sub> CN	Ph <sub>4</sub> SbI - Bu <sub>3</sub> SnI	48
11	"		"	HMPA	Ph <sub>4</sub> SbI - Bu <sub>3</sub> SnI	47
12	"		"	HMPA	Ph <sub>4</sub> SbI - ZnI <sub>2</sub>	23 + 31 <sup>c)</sup>
13	"		p-ClC <sub>6</sub> H <sub>4</sub>	PhH	Ph <sub>4</sub> SbI - Bu <sub>3</sub> SnI	100
14	"		p-MeC <sub>6</sub> H <sub>4</sub>	"	Ph <sub>4</sub> SbI - Bu <sub>3</sub> SnI	90
15 <sup>d)</sup>	"		iso-C <sub>3</sub> H <sub>7</sub>	"	Ph <sub>4</sub> SbI - Bu <sub>3</sub> SnI	78
16 <sup>e)</sup>	-(CH <sub>2</sub> ) <sub>3</sub> -		Ph	"	Ph <sub>4</sub> SbI - Bu <sub>3</sub> SnI	94
17 <sup>f)</sup>	Ph	Ph (trans)	"	"	Ph <sub>4</sub> SbI - Bu <sub>3</sub> SnI	65 (trans)

a) Oxirane 10 mmol, R-NCO 10 mmol, catalyst 1 mmol, solvent 5 ml, 80 °C, 2 h, isocyanate was added slowly over 60 min. b) All phenyl isocyanate turned into the trimer within 10 h. c) Iminodioxolane derivative(see reference 6), determined by GC. d) 70 °C, 1 h. e) 60 °C, 1 h. f) 80 °C, 28 h.

previously reported, can cleave monosubstituted oxiranes at their hindered site to produce 3,4-disubstituted oxazolidinones predominantly.<sup>2a)</sup> This result suggests that Ph<sub>4</sub>SbI may possess an inherent catalytic ability toward the reaction with 2,3-disubstituted oxiranes. We wish to describe herein that the reaction of inactive oxiranes such as cyclohexene oxide and cyclopentene oxide with isocyanates can be promoted by a catalyst, Ph<sub>4</sub>SbI activated by Bu<sub>3</sub>SnI. Several oxazolidinones were obtained in high yields as shown in Table 1.

In a typical procedure, to a mixture of cyclohexene oxide (10 mmol),  $\text{Ph}_4\text{SbI}$  (1 mmol),  $\text{Bu}_3\text{SnI}$  (1 mmol) and benzene (3 ml), phenyl isocyanate (10 mmol) in benzene (2 ml) was added dropwise during a course of 60 min at 80 °C. After additional 2 h the resultant mixture was subjected to separation by column chromatography on silica gel, and 4,5-tetramethylene-3-phenyl-1,3-oxazolidin-2-one (**3a**) was eluted by  $\text{CHCl}_3$  (99% yield).

As summarized in Table 1, the combination catalyst of  $\text{Ph}_4\text{SbI}$  and  $\text{Bu}_3\text{SnI}$  showed the highest activity, yielding **3a** quantitatively, while either of them exhibited little activity individually. Representative conventional catalysts,  $\text{Bu}_4\text{NI}$  or  $\text{LiBr}$ , gave only the trimer of phenyl isocyanate. It is notable that this reaction media is considered to be almost neutral because of lower acidity of the both of  $\text{Ph}_4\text{SbI}$  and  $\text{Bu}_3\text{SnI}$ .<sup>3)</sup> This cycloaddition appeared to be highly dependent on the nature of the additives to  $\text{Ph}_4\text{SbI}$  and solvents as revealed in Table 1. The efficiency of additives decreases in the following order;  $\text{Bu}_3\text{SnI} > \text{Bu}_2\text{SnI}_2 \gg \text{SnI}_2$ ,  $\text{ZnI}_2 > \text{AlI}_3$ . Aluminum triiodide<sup>4)</sup> caused predominantly polymerization of cyclohexene oxide due to its strong acidity. In particular, it is noteworthy that the addition of  $\text{CuI}$  completely depressed the catalytic ability of  $\text{Ph}_4\text{SbI}$ , and isocyanate was recovered nearly quantitatively. Consequently,  $\text{Bu}_3\text{SnI}$  of lowest acidity was the most suitable additive. Among various solvents examined, benzene was found to be a solvent of choice, while polar solvents such as acetonitrile and hexamethylphosphoramide (HMPA) were unsatisfactory. These polar solvents have coordinate ability to organotin halides,<sup>5)</sup> which would weaken an interaction of  $\text{Ph}_4\text{SbI}$  with  $\text{Bu}_3\text{SnI}$ , leading to a lower yield of **3a**. Of interest is the observation that 4,5-tetramethylene-2-phenylimino-1,3-dioxolane (**4**) was produced by  $\text{Ph}_4\text{SbI}$ - $\text{ZnI}_2$  catalyst in HMPA (entry 12).<sup>6)</sup> To our knowledge, this is the first example of isolation of **4** in the direct reaction of phenyl isocyanate with oxiranes, although it has been assumed to be an intermediate of the formation of oxazolidinones.<sup>7)</sup>

The reaction of trans-stilbene oxide with phenyl isocyanate gave trans-4,5-diphenyloxazolidinone<sup>8)</sup> in 65% yield without the formation of cis isomer. Stereospecificity of this reaction promoted by this type of catalysts is now under investigation.

This study was partially supported by the Grant-in-Aids for Scientific Research No.60430022 from the Ministry of Education, Science and Culture.

## References

- 1) For example, M. E. Dyen and D. Swern, *Chem. Rev.*, 67, 197 (1967), and references cited therein: J. E. Herweh and W. J. Kauffman, *Tetrahedron Lett.*, 1971, 809.
- 2) a) A. Baba, M. Fujiwara, and H. Matsuda, *Tetrahedron Lett.*, 27, 77 (1986);  
b) I. Shibata, A. Baba, H. Iwasaki, and H. Matsuda, *J. Org. Chem.*, 51, 2177 (1986).
- 3) R. Nomura, A. Ninagawa, and H. Matsuda, *J. Org. Chem.*, 45, 3735 (1980);  
N. Iwamoto, A. Ninagawa, H. Matsuda, and S. Matsuda, *Kogyo Kagaku Zasshi*, 70, 1400 (1971).
- 4)  $AlI_3$  was prepared in situ; S. Andersson, *Synthesis*, 1985, 437.
- 5) A. G. Davies and P. J. Smith, "Comprehensive Organometallic Chemistry," ed by G. Wilkinson, F. G. A. Stone, and W. Abel, Pergamon Press, New York (1982), Vol. 2, p. 519.
- 6) 4,5-Tetramethylene-2-phenylimino-1,3-dioxolane; mp 84-85 °C; IR (KBr)  $1700\text{ cm}^{-1}$  (C=O); MS (m/e) 217 ( $M^+$ );  $^1H$  NMR ( $CDCl_3$ )  $\delta$  = 1.1-2.2 (8H, m), 4.45-4.7 (2H, m), 6.8-7.4 (5H, m);  $^{13}C$  NMR ( $CDCl_3$ )  $\delta$  = 19.6 (t), 19.9 (t), 26.7 (t), 27.0 (t), 75.1 (d), 77.0 (d), 123.11 (d), 123.17 (d), 128.5 (d), 145.5 (s), 153.2 (s). Found: C, 71.78; H, 6.93; N, 6.37%. Calcd for  $C_{13}H_{15}NO_2$ : C, 71.87; H, 6.96; N, 6.45%.
- 7) K. Gulbins and K. Hamann, *Angew. Chem.*, 73, 434 (1961); T. Mukaiyama, T. Fujisawa, H. Nohira, and T. Hyugaji, *J. Org. Chem.*, 27, 3337 (1962).
- 8) trans-4.5-Diphenyl-3-phenyloxazolin-2-one; mp 128 °C; IR (KBr)  $1740\text{ cm}^{-1}$  (C=O); MS (m/e) 315 ( $M^+$ );  $^1H$  NMR ( $CDCl_3$ )  $\delta$  = 5.14 (1H, d,  $J$  = 6.3 Hz), 5.27 (1H, d,  $J$  = 6.3 Hz), 6.9-7.6 (15H, m).

(Received September 1, 1986)