## A concise approach towards the synthesis of steganone analogues

## Adrian Bradley,<sup>a</sup> William B. Motherwell<sup>\*a</sup> and Feroze Ujjainwalla<sup>b</sup>

<sup>a</sup> Department of Chemistry, Christopher Ingold Laboratories, University College London, 20 Gordon Street, London, UK WC1H 0AJ. E-mail: w.b.motherwell@ucl.ac.uk

<sup>b</sup> Department of Chemistry, Imperial College of Science, Technology and Medicine, South Kensington, London, UK SW7 2AY

Received (in Cambridge, UK) 27th January 1999, Accepted 12th April 1999

The cobalt-mediated [2+2+2] cycloaddition of a tethered deca-1,9-diyne is used as a key step in a highly convergent route to steganone analogues.

Steganone **1** is one member of the family of four bisbenzocyclooctadiene lignan lactones isolated from the Ethiopean shrub *Steganotaenia araliciae* Hochst and characterised by Kupchan<sup>1</sup>



in 1973. These compounds have displayed significant activity in vivo against P-388 leukemia in mice and in vitro against cells derived from human carcinoma of the nasopharynx (KB). They act as spindle poisons and function by stabilising dimeric tubulin and thereby inhibiting formation of microtubules.<sup>2</sup> These intriguing structures, comprising an eight-membered ring fused to a biaryl moiety, are composed of three stereochemical elements, two deriving from the trans-fused y-lactone, whilst the third is the stereogenic axis of the atropisomeric biaryl unit. Since its discovery, steganone 1 has proven to be a favoured target for sustained synthetic effort, and for more than two decades a wide variety of synthetic strategies have culminated in a number of total/formal syntheses.3 Careful scrutiny of these approaches reveals striking parallels in retrosynthetic analysis, with no less than ten of the syntheses requiring a late stage installation of the trans-fused y-lactone. Moreover, in every single instance to date some variant of a biaryl coupling reaction is used to forge the crucial carbon-carbon bond between the two aromatic rings.

Since the properties of steganone 1 itself are well known, our own objective in this area was to develop a highly convergent route which would be potentially amenable to the syntheses of a wide variety of differentially substituted aromatic analogues for use in structure-activity studies. To the best of our knowledge, only a single analogue of this type has thus far been reported.<sup>4</sup> As shown in Scheme 1, the key element of the present strategy involves the use of the tethered divne 2 in a cobalt mediated [2+2+2] cycloaddition,<sup>5</sup> thereby enabling closure of the eight-membered carbocyclic ring with concomitant construction of a usefully functionalised northern aryl unit for further elaboration. We also envisaged that the required trans stereochemistry round the  $\gamma$ -lactone 2 would be derived by alkylation of the lactone enolate from 3 with the benzylic bromide 4 or congeners thereof which would allow yet further variation in aromatic functionality to be introduced at an antepenultimate stage.

In the event, as outlined in Scheme 2, it was possible to develop a concise and highly efficient route to multigram

quantities of the required divne 2. Thus, the protected lactone 3 was readily prepared from racemic paraconic acid 5 via palladium-catalysed coupling of the derived acid chloride with Me<sub>3</sub>SiC=CSnMe<sub>3</sub> followed by ketalisation of the resultant ynone 6, whilst the known benzylic alcohol<sup>7</sup> 7 was available from 3,4,5-trimethoxybenzaldehyde via a straightforward sequence involving iodination, borohydride reduction, and a second palladium catalysed coupling reaction of the resultant iodoarene with Me<sub>3</sub>SiC=CH. The transformation to the corresponding bromide 4 was then acheived using the PPh<sub>3</sub>-CBr<sub>4</sub> protocol. The coupling reaction of the two fragments proceeded smoothly through prior generation of the lithium enolate of lactone 3 with LDA at -78 °C followed by alkylation with the benzylic bromide 4 at -35 °C to give 8 as a single diastereoisomer. Deprotection of the acetylenic groups then furnished the parent divne 2.

Our attention then focused on the crucial CpCo(CO)<sub>2</sub>mediated [2+2+2] cycloaddition step using diyne 2, Me<sub>3-</sub> SiC=CSiMe<sub>3</sub> as the third acetylenic component, and the standard irradiation conditions developed by Vollhardt.5 To our consternation, initial experiments failed to yield any significant amounts of the desired biaryl 9, and the beautifully crystalline orange material which could be isolated as a single diastereoisomer in up to 57% yield was shown by X-ray crystallography to be the cobaltacyclobutadiene complex 10 (Scheme 3). Although the formation of similar complexes has been previously observed<sup>9</sup> and is generally considered to arise when either co-ordination or insertion of the third alkyne component is problematic, it was nevertheless encouraging to note that the structure 10 confirmed both the viability of the eight-membered ring closure and also the trans stereochemistry around the ylactone. Gratifyingly however, whether via the intermediacy of the putative metallocyclopentadiene precursor to 10 or one of the other two possible candidates from reaction with Me<sub>3</sub>-

[2+2+2] /cloadditior Сус MeC MeO MeC MeO ÓМе ÓΜε 2 SiMe<sub>3</sub> MeO R Me<sub>3</sub>Si MeO ÓMe 3 4 Scheme 1



Scheme 2 Reagents and conditions: i,  $(COCl)_2$ ,  $CH_2Cl_2$ , DMF (cat), room temp., 1 h; ii,  $Me_3SiC\equiv CSnMe_3$ ,  $Pd(PPh_3)_2Cl_2$ ,  $CICH_2CH_2Cl$ , 50 °C, 16 h, 67% over 2 steps; iii, HOCH\_2CH\_2OH, PhH, PPTS (cat), reflux, 12 h, 87%; iv,  $I_2$ ,  $AgO_2CCF_3$ ,  $CH_2Cl_2$ , 20 °C, 12 h, quant.; v, NaBH<sub>4</sub>, MeOH, -5 °C, 30 min, quant.; vi,  $Me_3SiC\equiv CH$ ,  $Pd(PPh_3)_2Cl_2$ , CuI,  $Et_2NH$ , 50 °C, 2 h, 92%; viii,  $At_1DA$ ,  $THF_7 - 78 °C$ , 1 h, then add  $3t_7 - 35 °C$ , 1 h, 80%; ix,  $K_2CO_3$ , MeOH, 20 °C, 15 h, 95%.

 $SiC \equiv CSiMe_3$ , further variation in experimental conditions gave the desired biaryl **9** possessing the carbocyclic steganone core in 20% yield.

The structure of the indicated atropisomer of the biaryl **9**, which corresponds to the natural series, was established both by NOE experiments and also by deprotection of the ketal using wet formic acid which furnished the parent ketone **11** ( $v_{max}$  1674 cm<sup>-1</sup>) in 54% yield (Scheme 3). Deprotection accompanied by selective mono- or di-protodesilylation could also be achieved in a single step by treatment of **9** with TFA either at room temperature or at reflux to afford the further steganone analogues **12** and **13** in 73 and 51% yield, respectively.

In summary, these preliminary results demonstrate the viability of using a tethered deca-1,9-diyne in a cobalt-mediated [2+2+2] cyclobenzannulation strategy for the construction of steganone analogues in a highly convergent manner. Within this framework, current studies are directed towards the use of further  $2\pi$  addends and additional metal-based cyclotrimerisation reactions both for aromatic and heteroaromatic rings in order to further enhance the flexibility of this approach.

We thank the EPSRC and Zeneca Agrochemicals for the award of a CASE studentship to F. U., and University College London for the provision of a Provosts' studentship to A. B. We are also indebted to Professor D. Williams and Dr A. M. Z. Slawin for X-ray structural determination and the University of London Intercollegiate Research School (ULIRS) for mass spectral measurements. Finally, one of us (W. B. M.) wishes to



Scheme 3 Reagents and conditions: i, Me<sub>3</sub>SiC=CSiMe<sub>3</sub>, CpCo(CO)<sub>2</sub>, 50–65 °C, reflux, hv, addition of 2 in THF over 9 h; ii, Me<sub>3</sub>SiC=CSiMe<sub>3</sub>, MeCN, reflux, hv, addition of 2 in THF over 9 h.

acknowledge his personal debt of gratitude to the late Professor R. A. Raphael FRS, whose mastery of the Art of Organic Synthesis was matched only by his ability to encourage and to instil his love and enthusiasm for the subject in others.

## Notes and references

- 1 S. M. Kupchan, R. W. Britton, M. F. Ziegler, C. J. Gilmore, R. J. Restivo and R. Bryan, J. Am. Chem. Soc., 1973, 95, 1335
- 2 R. W. J. Wang, L. I. Rebbun and M. S. Kupchan, *Cancer Res.*, 1977, 37, 3071.
- 3 N. S. Narasimhan and I. R. Aiden, Tetrahedron Lett., 1988, 29, 2987; N. S. Narasimhan and I. R. Aiden, Indian J. Chem., Sect. B., 1993, 32B, 211; A. I. Meyers, J. R. Flisak and R. A. Aitken, J. Am. Chem. Soc., 1987, 109, 5446; P. Magnus, J. Schultz and T. Gallagher, J. Am. Chem. Soc., 1985, 107, 4984; R. Dhal, J. P. Robin and E. Brown, Tetrahedron 1983, 39, 2787; M. Mervic, Y. Ben-David and E. Ghera, Tetrahedron Lett., 1981, 22, 5091; F. E. Ziegler, I. C. Chliwner, K. W. Fowler, S. J. Kanfer, S. J. Kuo and N. D. Sinha, J. Am. Chem. Soc., 1980, 102, 790; E. Brown, R. Dhal and J. P. Robin, Tetrahedron Lett., 1979, 733; E. R. Larson and R. A. Raphael, Tetrahedron Lett., 1979, 5041; G. R. Krow, K. M. Damadoran, E. Michener, R. Wolf and J. Gaure, J. Org. Chem., 1978, 43, 3950; D. Becker, L. R. Hughes and R. A. Raphael, J. Chem. Soc., Perkin Trans. 1, 1977, 1674; L. R. Hughes and R. A. Raphael, Tetrahedron Lett., 1976, 1543; A. S. Kende and L. S. Liebskind, J. Am. Chem. Soc., 1976, 98, 267; M. Uemura, A. Daimon and Y. Hayashi, J. Chem. Soc., Chem. Commun., 1995, 1943.
- 4 D. Becker, L. R. Hughes and R. A. Raphael, J. Chem. Soc., Chem. Commun., 1974, 430.
- 5 K. P. C.Vollhardt, Angew. Chem., Int. Ed. Engl., 1984, 28, 539.
- 6 J. F. Tocanne and C. Asselineau, Bull. Chem. Soc. Fr., 1965, 3346.
- 7 M. E. Garst and B. J. McBride, J. Org. Chem., 1989, 54, 249.
- 8 F. Ujjainwalla, PhD thesis, University of London, 1993. We are grateful to Drs D. Williams and A. M. Z. Slawin for the crystal structure determination, which will be discussed in the full paper.
- 9 R. Gleiter, D. Kratz, M. L. Ziegler and B. Nuber, *Tetrahedron Lett.*, 1990, **31**, 6175; C.-A. Chang, J. A. King Jr. and K. P. C. Vollhardt, *J. Chem. Soc., Chem. Commun.*, 1981, 53; C.-A. Chang, C. G. Francisco, T. R. Gadek, J. A. King Jr., E. D. Sternberg and K. P. C. Vollhardt, in *Organic Synthesis Today and Tomorrow*, ed. B. M. Trost and C. R. Hutchinson, Pergamon, New York, 1981, 71; P. Phanasavath, C. Aubert and M. Malacria, *Tetrahedron Lett.*, 1998, **39**, 1561.

Communication 9/00743A