


Stefania De Montis, Claudia Fattuoni,* Enzo Cadoni, Maria G. Cabiddu,
Michele Usai, Salvatore Cabiddu

Dipartimento di Scienze Chimiche, University of Cagliari, Cittadella Universitaria di Monserrato,
SS. 554, Bivio per Sestu. I-09042 Monserrato (CA), Italy
Tel.: +39-070-6754387; fax: +39-070-6754388; E-mail: cfattuon@unica.it

Received March 28, 2008



4*H*-1,4-Benzothiazine-1,1-dioxide derivatives were synthesized through a sequence of almost quantitative reactions. The commercial starting material 2-(methylsulfonyl)aniline was Boc-protected, *N*-acylated and oxidized at the sulfur atom to obtain a sulfonyl derivative. An anionic transposition of the acyl group followed by a simultaneous deprotection-cyclization gave the title products in excellent yields. All products and intermediates were fully characterized.

J. Heterocyclic Chem., **45**, 1445 (2008).

INTRODUCTION

1,4-Benzothiazine derivatives have drawn much interest in the pharmacologic field due to their structural similarity with the phenothiazine system [1]. In particular the synthesis of the *S*-oxidized system, 4*H*-1,4-benzothiazine-1,1-dioxide, has been the object of several studies and patents for industrial and pharmacological applications [2-5]. The synthesis of such systems usually started from 2-aminobenzenethiols which were treated with β -diketones to give the 2,3-disubstituted heterocyclic ring through the formation of an intermediate enaminoketone [4-8]. The further step, *i.e.* the oxidation at the sulfur, was usually performed with 30% hydrogen peroxide in glacial acetic acid [6-8]. An alternative method reported by Schou started from pyridinium sulfonate salts which were converted into 2-cyanomethylsulfonyl acetanilides: base catalyzed ring closure gave the *S*-oxidized benzothiazinic system [3]. Another recent method involved the microwave assisted cyclization of α -phenylsulfonyl enaminoacrylates [9].

In this paper we report a new high yield synthesis of 3-substituted 4*H*-1,4-benzothiazine-1,1-dioxides: this procedure is characterized by few almost quantitative steps allowing the preparation of the target compounds bearing a free, unsubstituted NH suitable for further functionalization. The starting compound 2-(methylsulfonyl)aniline (**1**) is commercial and all reagents are easily available and inexpensive.

RESULTS AND DISCUSSION

The first step of the synthetic pathway consisted in the protection of the amino group in **1** with Boc (*tert*-

butoxycarbonyl), an easily removable protecting group, to give **2** (Scheme I). This compound was then lithiated with lithium diisopropylamide (LDA) at -50 °C and treated with an acyl chloride RCOCl where R was an aryl, heteroaryl or alkyl group to give the *N*-acylated products **3a-e** (yield 60-77%, Table 1). Compounds **3a-e** were oxidized at the sulfur atom with *m*-chloroperbenzoic acid (MCPBA) to give the sulfones **4a-e** (yield 80-100%, Table 1), which after treatment with butyllithium (BuLi) at -50 °C underwent an acyl migration from the nitrogen atom to the methylsulfonyl group affording products **5a-e** (yield 71-100%, Table 1). The acyl group transfer is likely to occur by an intramolecular mechanism: in fact, from the reaction mixture no product of type **A** or **B** could be isolated, even performing the reaction with a higher substrate concentration (Figure 1).




Figure 1

The treatment of compounds of type **5** with trifluoroacetic acid (TFA) allowed the simultaneous Boc-deprotection and cyclization giving products **6a-e** (yield 70-100%, Table 1). It is noteworthy that products of type **5** did not cyclize in the presence of BuLi, through an attack of the anionic nitrogen to the carbonyl group: this is due to the low reactivity as nucleophile of the Boc-protected N, even as an anion (Figure 2).

Scheme I




Figure 2

The cyclization took place only when the N was deprotected in an acidic medium: the attack of the free NH_2 to the carbonyl carbon did not lead to the expected imine, but to an enaminic product because of the strong acidity of the $\text{SO}_2\text{-CH}_2$ hydrogens.

All products were fully characterized by elemental analysis, IR, $^1\text{H-NMR}$, $^{13}\text{C-NMR}$ and mass spectroscopy (see Experimental).

The ^{13}C NMR spectra of products **3e** and **4e**, bearing two *tert*-butyl groups, showed a remarkable feature: the signals due to the methyl groups of the *tert*-butyl moieties are observed as an unresolved multiplet. This phenomenon clearly depended upon the closeness of such bulky groups, whose free rotation around the C-O or C-C bond was forbidden. When the same spectra were recorded at 55 °C, the signals relative to the CH_3 appeared as two clean singlets.

Table 1. Yields of pure isolated products **3a-e**, **4a-e**, **5a-e** and **6a-e**.

R		3	4	5	6
Ph	a	77	90	96	100
4-MeOC ₆ H ₄	b	70	92	71	81
4-FC ₆ H ₄	c	65	92	78	75
	d	65	100	80	81
C(CH ₃) ₃	e	60	80	100	70

In conclusion, we have reported an easy method to prepare monosubstituted 4*H*-1,4-benzothiazine-1,1-dioxides, using inexpensive reagents and allowing to introduce in the 3-position different groups (aryl, heteroaryl or alkyl).

EXPERIMENTAL

NMR spectra were recorded on a Varian VXR-300 spectrometer with tetramethylsilane as internal reference. IR spectra were recorded on a Perkin-Elmer 1310 grating spectrophotometer. The GC-MS analyses were performed at 70 eV with a Hewlett-Packard 5989A GC-MS system with HP 5890 GC fitted with a capillary column (50 m×0.2 mm) packed with DH 50.2 Petrocol (0.50 μm film thickness) using a direct-inlet system for solid products. Chromatography was performed on silica gel 60, 0.04-0.063 mm (Fluka). Melting points were determined with a Kofler hot stage microscope and are uncorrected. Microanalyses were carried out on a Carlo Erba 1106 element analyzer. Reagent-grade commercially available reagents and solvents were used. Solutions of BuLi in hexane were purchased from Aldrich Chemical Company and were analyzed before use [10]. 2-(Methylsulfonyl)aniline (**1**) (Fluka) was protected by literature method to give **2** [11].

General procedure for the synthesis of compounds **3a-e**.

To a stirred solution of LDA (24.0 mL, 31.6 mmol) in anhydrous THF (60 mL) at -50 °C, a solution of **2** (3.00 g, 12.6 mmol) in anhydrous THF (60 mL) was added under argon, and stirring was continued for 1 h. Then the appropriate acyl chloride was added at the same temperature, and after 30 minutes the mixture was poured into a saturated solution of aqueous NH_4Cl . The organic layer was separated, the aqueous layer extracted with CH_2Cl_2 , the extracts combined, dried (Na_2SO_4) and concentrated to give products **3a-e**.

tert-Butyl benzoyl[2-(methylthio)phenyl]carbamate (**3a**).

The crude product was purified by chromatography using diethyl ether/light petroleum (1:2) as eluent to give **3a** as colourless crystals, 3.33 g (77%), mp 122-123 °C (EtOH); ir (Nujol): 1725, 1670 cm^{-1} ; ^1H nmr (CDCl_3): δ 1.21 (s, 9H, CH_3C), 2.44 (s, 3H, SCH_3), 7.23 (m, 2H, ArH), 7.35 (m, 2H, ArH), 7.43 (m, 2H, ArH), 7.49 (m, 1H, ArH), 7.81 (m, 2H, ArH); ^{13}C nmr (CDCl_3): δ 15.7, 27.3, 83.4, 125.9, 127.3, 127.9, 128.1, 128.9, 129.3, 131.3, 136.9, 137.1, 137.7, 152.5, 172.0; ms: m/z 343 (M^+ , 5), 243 (6), 196 (48), 165 (56), 105 (100), 77 (72), 57 (77), 41 (36). *Anal.* calcd. for $\text{C}_{19}\text{H}_{21}\text{NO}_3\text{S}$: C, 66.45; H, 6.16; N, 4.08; S, 9.34. Found: C, 66.42; H, 6.18; N, 4.02; S, 9.30.

tert-Butyl 4-methoxybenzoyl[2-(methylthio)phenyl]-carbamate (**3b**).

The crude product was purified by chromatography using first light petroleum and then diethyl ether/light petroleum (1:2) as eluents to give **3b** as white crystals, 3.29 g (70%), mp 114-115 °C (EtOH/ H_2O); ir (nujol):

1720, 1665 cm^{-1} ; ^1H nmr (CDCl_3): δ 1.27 (s, 9H, CH_3C), 2.43 (s, 3H, SCH_3), 3.84 (s, 3H, OCH_3), 6.92 (d, 2H, $J=9.0$ Hz, ArH), 7.20 (m, 2H, ArH), 7.31 (m, 2H, ArH), 7.81 (d, 2H, $J=9.0$ Hz, ArH); ^{13}C nmr (CDCl_3): δ 15.6, 27.5, 55.3, 83.0, 113.2, 125.8, 127.1, 128.7, 128.7, 129.3, 130.7, 137.5, 137.7, 152.8, 162.4, 171.3; ms: m/z 373 (M^+ , 0.1), 273 (2), 226 (14), 165 (18), 135 (100), 107 (8), 77 (18), 57 (35), 41 (32). *Anal.* calcd. for $\text{C}_{20}\text{H}_{23}\text{NO}_4\text{S}$: C, 64.32; H, 6.21; N, 3.75; S, 8.59. Found: C, 64.28; H, 6.25; N, 3.78; S, 8.61.

tert-Butyl 4-fluorobenzoyl[2-(methylthio)phenyl]-carbamate (3c). The crude product was washed with light petroleum first and then with diethyl ether to give **3c** as white crystals, 2.96 g (65%), mp 109–111 $^\circ\text{C}$ (EtOH); ir (nujol): 1725, 1670 cm^{-1} ; ^1H nmr (CDCl_3): 1.35 (s, 9H, CH_3C), 2.51 (s, 3H, SCH_3), 7.20 (m, 2H, ArH), 7.30 (br s, 2H, ArH), 7.42 (br s, 2H, ArH), 7.93 (m, 2H, ArH); ^{13}C nmr (CDCl_3): δ 15.4, 27.3, 83.3, 114.9 ($J_{\text{CF}} = 22.0$ Hz), 125.7, 127.0, 128.9, 129.1, 130.6 ($J_{\text{CF}} = 8.6$ Hz), 132.7 ($J_{\text{CF}} = 3.6$ Hz), 136.9, 137.5, 152.3, 164.4 ($J_{\text{CF}} = 252.8$ Hz), 170.6; ms: m/z 361 (M^+ , 3), 261 (17), 214 (100), 165 (80), 123 (93), 95 (32), 57 (43), 41 (26). *Anal.* calcd. for $\text{C}_{19}\text{H}_{20}\text{FNO}_3\text{S}$: C, 63.14; H, 5.58; N, 3.88; S, 8.87. Found: C, 63.11; H, 5.61; N, 3.92; S, 8.92.

tert-Butyl 2-(methylthio)phenyl(2-thienylcarbonyl)-carbamate (3d). The crude product was purified by flash-chromatography using diethyl ether/light petroleum (1:2) as eluent to give **3d** as yellow crystals, 2.86 g (65%), mp 123–125 $^\circ\text{C}$ (EtOH); ir (nujol): 1720, 1665 cm^{-1} ; ^1H nmr (CDCl_3): δ 1.28 (s, 9H, CH_3C), 2.29 (s, 3H, SCH_3), 6.88 (br s, 1H, ArH), 7.08 (br s, 2H, ArH), 7.19 (br s, 2H, ArH), 7.39 (br s, 1H, ArH), 7.45 (br s, 1H, ArH); ^{13}C nmr (CDCl_3): δ 15.3, 27.4, 83.1, 125.6, 126.8, 126.8, 128.9, 129.2, 131.9, 132.6, 137.0, 138.1, 138.4, 152.3, 164.2; ms: m/z 349 (M^+ , 15), 294 (52), 250 (44), 249 (38), 232 (29), 202 (100), 165 (66), 111 (68), 57 (79), 41 (42). *Anal.* calcd. for $\text{C}_{17}\text{H}_{16}\text{NO}_3\text{S}_2$: C, 58.43; H, 5.48; N, 4.01; S, 18.35. Found: C, 58.39; H, 5.50; N, 4.03; S, 18.37.

tert-Butyl 2,2-dimethylpropanoyl[2-(methylthio)phenyl]-carbamate (3e). The crude product was purified by chromatography using diethyl ether/light petroleum (1:8) as eluent to give **3e** as colourless crystals, 2.44 g (60%), mp 69–71 $^\circ\text{C}$ (H_2O); ir (nujol): 1720 and 1665 cm^{-1} ; ^1H nmr (CDCl_3): δ 1.38 (s, 9H, CH_3C), 1.40 (s, 9H, CH_3C), 2.42 (s, 3H, SCH_3), 7.04 (d, 1H, $J = 7.6$ Hz, ArH), 7.16 (m, 1H, ArH), 7.27 (m, 2H, ArH); ^{13}C nmr (CDCl_3): δ 15.5, 27.9, 43.5, 82.4, 125.5, 126.9, 128.3, 128.5, 137.7, 138.3, 153.1, 183.7; ms: m/z 323 (M^+ , 6), 224 (13), 223 (17), 232 (29), 176 (47), 165 (98), 57 (100), 41 (60). *Anal.* calcd. for $\text{C}_{17}\text{H}_{25}\text{NO}_3\text{S}$: C, 63.13; H, 7.79; N, 4.33; S, 9.91. Found: C, 63.11; H, 7.82; N, 4.30; S, 9.94.

General procedure for the synthesis of compounds 4a-e. A solution of one compound of type **3** (5.6 mmol) in CH_2Cl_2 (150 mL) was treated with MCPBA (3.03 g, 12.3 mmol) at 0 $^\circ\text{C}$. After the addition was complete the temperature was allowed to warm to room temperature and the reaction was monitored by tlc till completion. The mixture was treated with 10% aqueous NaOH, the organic layer was dried (Na_2SO_4) and concentrated to give products **4a-e**.

tert-Butyl benzoyl[2-(methylsulfonyl)phenyl]carbamate (4a). The crude product was washed with diethyl ether to give **4a** as colourless crystals, 1.89 g (90%), mp 178–179 $^\circ\text{C}$ (EtOH); ir (nujol): 1715, 1680, 1310, 1155 cm^{-1} ; ^1H nmr (CDCl_3): δ 1.20 (s, 9H, CH_3C), 3.14 (s, 3H, SO_2CH_3), 7.40 (m, 4H, ArH), 7.62 (t, 1H, $J = 7.5$ Hz, ArH), 7.76 (m, 3H, ArH), 8.15 (d, 1H, $J = 7.5$ Hz, ArH); ^{13}C nmr (CDCl_3): δ 27.3, 44.4, 84.3, 128.0, 128.0,

129.5, 130.8, 131.4, 131.8, 134.7, 136.5, 137.6, 137.6, 152.1, 172.6; ms: m/z 276 (3), 197 (23), 196 (29), 105 (100), 77 (45), 57 (47), 41 (28). *Anal.* calcd. for $\text{C}_{19}\text{H}_{21}\text{NO}_5\text{S}$: C, 60.78; H, 5.64; N, 3.73; S, 8.54. Found: C, 60.72; H, 5.66; N, 3.76; S, 8.57.

tert-Butyl 4-methoxybenzoyl[2-(methylsulfonyl)phenyl]-carbamate (4b). The crude product was washed with diethyl ether to give **4b** as colourless crystals, 2.09 g (92%), mp 166–168 $^\circ\text{C}$ (EtOH); ir (nujol): 1715, 1670, 1310, 1150 cm^{-1} ; ^1H nmr (CDCl_3): δ 1.26 (s, 9H, CH_3C), 3.12 (s, 3H, SO_2CH_3), 3.87 (s, 3H, OCH_3), 6.96 (d, 2H, $J = 8.8$ Hz, ArH), 7.38 (d, 1H, $J = 7.8$ Hz, ArH), 7.60 (t, 1H, $J = 7.8$ Hz, ArH), 7.72 (t, 1H, $J = 7.8$ Hz, ArH), 7.79 (d, 2H, $J = 8.8$ Hz, ArH), 8.14 (d, 1H, $J = 7.8$ Hz, ArH); ^{13}C nmr (CDCl_3): δ 27.4, 44.3, 55.4, 83.9, 113.2, 128.1, 129.3, 130.7, 130.8, 131.8, 134.6, 137.5, 137.9, 152.4, 162.6, 171.8; ms: m/z 305 (2), 226 (13), 135 (100), 77 (7), 57 (14), 41 (10). *Anal.* calcd. for $\text{C}_{20}\text{H}_{23}\text{NO}_6\text{S}$: C, 59.24; H, 5.72; N, 3.45; S, 7.91. Found: C, 59.22; H, 5.74; N, 3.43; S, 7.90.

tert-Butyl 4-fluorobenzoyl[2-(methylsulfonyl)phenyl]-carbamate (4c). The crude product was washed with diethyl ether to give **4c** as white crystals, 2.02 g (92%), mp 178–179 $^\circ\text{C}$ (EtOH); ir (nujol): 1710, 1670, 1310, 1150 cm^{-1} ; ^1H nmr (CDCl_3): δ 1.25 (s, 9H, CH_3C), 3.12 (s, 3H, SO_2CH_3), 7.14 (t, 2H, $J = 8.2$ Hz, ArH), 7.38 (d, 1H, $J = 7.8$ Hz, ArH), 7.62 (t, 1H, $J = 7.8$ Hz, ArH), 7.74 (t, 1H, $J = 7.8$ Hz, ArH), 7.81 (m, 2H, ArH), 8.14 (d, 1H, $J = 7.8$ Hz, ArH); ^{13}C nmr (CDCl_3): δ 27.4, 44.4, 83.5, 115.2 ($J_{\text{CF}} = 22.0$ Hz), 129.6, 130.7 ($J_{\text{CF}} = 8.6$ Hz), 130.8, 131.8, 132.4 ($J_{\text{CF}} = 3.6$ Hz), 134.7, 137.4, 137.5, 152.1, 164.7 ($J_{\text{CF}} = 252.8$ Hz), 171.4; ms: m/z 293 (8), 214 (70), 197 (22), 123 (100), 77 (25), 57 (9), 41 (7). *Anal.* calcd. for $\text{C}_{19}\text{H}_{20}\text{FNO}_5\text{S}$: C, 58.00; H, 5.12; N, 3.56; S, 8.15. Found: C, 57.99; H, 5.14; N, 3.57; S, 8.18.

tert-Butyl 2-(methylsulfonyl)phenyl(2-thienylcarbonyl)-carbamate (4d). The crude product was washed with diethyl ether to give **4d** as colourless crystals, 2.13 g (100%), mp 138–139 $^\circ\text{C}$ (EtOH); ir (nujol): 1720, 1680, 1320, 1165 cm^{-1} ; ^1H nmr (CDCl_3): δ 1.24 (s, 9H, CH_3C), 3.04 (s, 3H, SO_2CH_3), 7.00 (t, 1H, $J = 4.3$ Hz, ArH), 7.27 (d, 1H, $J = 7.7$ Hz, ArH), 7.48 (m, 2H, ArH), 7.60 (m, 2H, ArH), 8.01 (d, 1H, $J = 7.7$ Hz, ArH); ^{13}C nmr (CDCl_3): δ 27.3, 44.1, 84.1, 127.0, 129.4, 130.5, 131.6, 132.0, 132.9, 134.5, 137.3, 137.4, 137.9, 152.1, 165.1; ms: m/z 282 (19), 281 (12), 264 (20), 202 (46), 197 (16), 111 (100), 57 (47), 41 (31). *Anal.* calcd. for $\text{C}_{17}\text{H}_{16}\text{NO}_5\text{S}_2$: C, 53.53; H, 5.02; N, 3.67; S, 16.81. Found: C, 53.50; H, 5.05; N, 3.65; S, 16.84.

tert-Butyl 2,2-dimethylpropanoyl[2-(methylsulfonyl)phenyl]carbamate (4e). The crude product was purified by flash-chromatography using diethyl ether/light petroleum (1:1) as eluent to give **4e** as colourless crystals, 1.59 g (80%), mp 119–120 $^\circ\text{C}$ (EtOH); ir (nujol): 1725, 1700, 1310, 1150 cm^{-1} ; ^1H nmr (CDCl_3): δ 1.37 (s, 9H, CH_3C), 1.45 (s, 9H, CH_3C), 3.10 (s, 3H, SO_2CH_3), 7.17 (d, 1H, $J = 7.6$ Hz, ArH), 7.53 (t, 1H, $J = 7.6$ Hz, ArH), 7.66 (t, 1H, $J = 7.6$ Hz, ArH), 8.07 (d, 1H, $J = 7.6$ Hz, ArH); ^{13}C nmr (CDCl_3): δ 27.6, 43.2, 69.0, 83.5, 128.7, 130.3, 130.9, 134.4, 136.9, 139.4, 152.1, 183.6; ms: m/z 197 (15), 176 (14), 171 (29), 149 (18), 85 (18), 57 (100), 41 (47). *Anal.* calcd. for $\text{C}_{17}\text{H}_{25}\text{NO}_5\text{S}$: C, 57.44; H, 7.09; N, 3.94; S, 9.02. Found: C, 57.41; H, 7.11; N, 3.92; S, 9.03.

General procedure for the synthesis of compounds 5a-e. A solution of one compound of type **4** (1.3 mmol) in anhydrous THF (50 mL) was treated dropwise with a 1.6 *M* solution of BuLi in hexane (2.0 mL, 3.2 mmol) at -50 $^\circ\text{C}$ under argon, and stirring was continued for 10 minutes. The mixture was poured

into a saturated solution of aqueous NH_4Cl , the organic layer was separated, the aqueous layer extracted with CH_2Cl_2 , the extracts combined, dried (Na_2SO_4) and concentrated to give products **5a-e**.

tert-Butyl 2-[(2-oxo-2-phenylethyl)sulfonyl]phenyl-carbamate (5a). The crude product was washed with diethyl ether to give **5a** as colourless crystals, 0.47 g (96%); mp 118–120 °C (EtOH); ir (nujol): 3350, 1730, 1675, 1315, 1155 cm^{-1} ; ^1H nmr (CDCl_3): δ 1.51 (s, 9H, CH_3C), 4.76 (s, 2H, SO_2CH_2), 7.07 (t, 1H, $J = 8.2$ Hz, ArH), 7.41 (t, 2H, $J = 7.5$ Hz, ArH), 7.56 (m, 2H, ArH), 7.78 (d, 1H, $J = 8.2$ Hz, ArH), 7.85 (d, 2H, $J = 8.2$ Hz, ArH), 8.30 (d, 1H, $J = 8.2$ Hz, ArH), 8.75 (s, 1H, NH); ^{13}C nmr (CDCl_3): δ 27.9, 62.4, 81.1, 120.5, 122.2, 124.2, 128.6, 128.8, 130.1, 134.0, 135.4, 135.4, 138.4, 151.8, 187.1; ms: m/z 375 (M^+ , 7), 319 (12), 275 (40), 152 (15), 106 (45), 105 (100), 77 (19), 57 (90), 41 (37). *Anal.* calcd. for $\text{C}_{19}\text{H}_{21}\text{NO}_5\text{S}$: C, 60.78; H, 5.64; N, 3.73; S, 8.54. Found: C, 60.75; H, 5.66; N, 3.75; S, 8.57.

tert-Butyl 2-[(2-(4-methoxyphenyl)-2-oxoethyl)sulfonyl]phenylcarbamate (5b). The crude product was washed with light petroleum to give **5b** as colourless crystals, 0.37 g (71%), mp 114–117 °C (EtOH/ H_2O); ir (nujol): 3340, 1720, 1665, 1310, 1155 cm^{-1} ; ^1H nmr (CDCl_3): δ 1.49 (s, 9H, CH_3C), 3.83 (s, 3H, OCH_3), 4.67 (s, 2H, SO_2CH_2), 6.87 (d, 2H, $J = 8.1$ Hz, ArH), 7.07 (t, 1H, $J = 7.9$ Hz, ArH), 7.54 (t, 1H, $J = 7.9$ Hz, ArH), 7.76 (d, 1H, $J = 7.9$ Hz, ArH), 7.83 (d, 2H, $J = 8.1$ Hz, ArH), 8.30 (d, 1H, $J = 7.9$ Hz, ArH), 8.72 (s, 1H, NH); ^{13}C nmr (CDCl_3): δ 28.1, 55.5, 62.5, 81.2, 114.0, 120.7, 122.4, 124.4, 128.6, 130.2, 131.5, 135.5, 138.5, 152.0, 164.4, 185.3; ms: m/z 405 (M^+) (4), 305 (32), 135 (100), 77 (13), 57 (47), 41 (78). *Anal.* calcd. for $\text{C}_{20}\text{H}_{23}\text{NO}_6\text{S}$: C, 59.24; H, 5.72; N, 3.45; S, 7.91. Found: C, 59.22; H, 5.75; N, 3.48; S, 7.90.

tert-Butyl 2-[(2-(4-fluorophenyl)-2-oxoethyl)sulfonyl]phenylcarbamate (5c). The crude product was purified by chromatography using diethyl ether/light petroleum (1:1) as eluent to give **5c** as colourless crystals, 0.40 g (78%), mp 137–139 °C (EtOH); ir (nujol): 3350, 1730, 1670, 1320, 1155 cm^{-1} ; ^1H nmr (CDCl_3): δ 1.44 (s, 9H, CH_3C), 4.63 (s, 2H, SO_2CH_2), 7.06 (m, 3H, ArH), 7.51 (t, 1H, $J = 8.0$ Hz, ArH), 7.70 (d, 1H, $J = 8.0$ Hz, ArH), 7.85 (m, 2H, ArH), 8.25 (d, 1H, $J = 8.0$ Hz, ArH), 8.60 (s, 1H, NH); ^{13}C nmr (CDCl_3): 28.2, 62.8, 81.5, 116.1 ($J_{\text{CF}} = 22.0$ Hz), 120.9, 122.6, 124.3, 130.3, 132.0 ($J_{\text{CF}} = 9.7$ Hz), 135.8, 138.6, 138.6, 152.0, 166.4 ($J_{\text{CF}} = 257.6$ Hz), 185.6; ms: m/z 393 (M^+ , 2), 293 (32), 211 (10), 171 (17), 152 (15), 123 (65), 106 (20), 95 (12), 57 (100), 41 (48). *Anal.* calcd. for $\text{C}_{19}\text{H}_{19}\text{FNO}_5\text{S}$: C, 58.00; H, 5.12; N, 3.56; S, 8.15. Found: C, 57.99; H, 5.14; N, 3.58; S, 8.18.

tert-Butyl 2-[(2-oxo-2-(2-thienyl)ethyl)sulfonyl]phenyl-carbamate (5d). The crude product was purified by chromatography using diethyl ether/light petroleum (1:1) as eluent to give **5d** as colourless crystals, 0.40 g (80%), mp 104–106 °C (EtOH/ H_2O); ir (nujol): 3360, 1740, 1670, 1330, 1170 cm^{-1} ; ^1H nmr (CDCl_3): δ 1.53 (s, 9H, CH_3C), 4.64 (s, 2H, SO_2CH_2), 7.09–7.14 (m, 2H, ArH), 7.58 (t, 1H, $J = 8.0$ Hz, ArH), 7.68 (d, 1H, $J = 4.5$ Hz, ArH), 7.74 (d, 1H, $J = 4.5$ Hz, ArH), 7.81 (d, 1H, $J = 8.0$ Hz, ArH), 8.32 (d, 1H, $J = 8.0$ Hz, ArH), 8.71 (s, 1H, NH); ^{13}C nmr (CDCl_3): δ 28.2, 63.7, 81.4, 120.8, 122.5, 124.1, 128.6, 130.3, 134.8, 135.8, 136.5, 138.6, 143.0, 152.0, 179.1; ms: m/z 381 (M^+ , 5), 282 (21), 281 (37), 152 (14), 111 (35), 106 (20), 57 (100), 41 (48). *Anal.* calcd. for $\text{C}_{17}\text{H}_{19}\text{NO}_5\text{S}_2$: C, 53.53; H, 5.02; N, 3.67; S, 16.81. Found: C, 53.50; H, 5.05; N, 3.65; S, 16.82.

tert-Butyl 2-[(3,3-dimethyl-2-oxobutyl)sulfonyl]phenyl-carbamate (5e). The crude product was washed with diethyl ether to give **5e** as colourless crystals, 0.46 g (100%), mp 56–58 °C (EtOH/ H_2O); ir (nujol): 3350, 1730, 1710, 1320, 1150 cm^{-1} ; ^1H nmr (CDCl_3): δ 1.04 (s, 9H, CH_3C), 1.46 (s, 9H, CH_3C), 4.24 (s, 2H, SO_2CH_2), 7.10 (t, 1H, $J = 7.9$ Hz, ArH), 7.52 (t, 1H, $J = 7.9$ Hz, ArH), 7.78 (d, 1H, $J = 7.9$ Hz, ArH), 8.23 (d, 1H, $J = 7.9$ Hz, ArH), 8.64 (s, 1H, NH); ^{13}C nmr (CDCl_3): δ 25.4, 28.2, 45.3, 59.9, 81.4, 121.9, 122.6, 125.3, 130.4, 135.4, 138.3, 152.2, 202.3; ms: m/z 355 (M^+ , 14), 299 (17), 256 (18), 198 (67), 171 (15), 156 (19), 57 (100), 41 (66). *Anal.* calcd. for $\text{C}_{17}\text{H}_{25}\text{NO}_5\text{S}$: C, 57.44; H, 7.09; N, 3.94; S, 9.02. Found: C, 57.42; H, 7.10; N, 3.97; S, 9.05.

General procedure for the synthesis of compounds 6a-e. A solution of one compound of type **5** (1.3 mmol), CH_2Cl_2 (4.5 mL) and trifluoroacetic acid (1.9 mL, 24.3 mmol) was stirred at room temperature for 10 minutes, then cooled at 0 °C and treated with 20% aqueous NaOH. The solid thus separated was filtered and ir lamp dried.

3-Phenyl-4H-1,4-benzothiazine 1,1-dioxide (6a). White crystals, 0.33 g (100%), mp 250–252 °C (EtOH) (lit. [12] 271–274 °C); ir (nujol): 3300, 1310, 1110 cm^{-1} ; ^1H nmr (CDCl_3): δ 6.39 (s, 1H), 7.45 (t, 1H, $J = 7.5$ Hz), 7.66–7.77 (m, 5H), 7.83–7.86 (m, 2H), 8.00 (d, 1H, $J = 7.5$ Hz), 11.02 (s, 1H, D_2O exchangeable, NH); ^{13}C nmr (CDCl_3): δ 96.4, 118.5, 122.1, 123.6, 123.7, 127.7, 129.1, 130.9, 132.4, 133.9, 137.2, 145.3; ms: m/z 257 (M^+ , 30), 193 (100), 165 (17). *Anal.* calcd. for $\text{C}_{14}\text{H}_{11}\text{NO}_2\text{S}$: C, 65.35; H, 4.31; N, 5.44; S, 12.46. Found: C, 65.32; H, 4.34; N, 5.47; S, 12.49.

3-(4-Methoxyphenyl)-4H-1,4-benzothiazine 1,1-dioxide (6b). White crystals, 0.30 g (81%), mp 278–280 °C (EtOH) (lit. [12] 256–260 °C); ir (nujol): 3280, 1260, 1115 cm^{-1} ; ^1H nmr (DMSO): δ 3.95 (s, 3H, OCH_3), 6.22 (s, 1H), 7.20 (d, 2H, $J = 8.6$ Hz), 7.39 (t, 1H, $J = 7.4$ Hz), 7.63 (d, 1H, $J = 7.4$ Hz), 7.70 (t, 1H, $J = 7.4$ Hz), 7.82 (d, 2H, $J = 8.6$ Hz), 7.93 (d, 1H, $J = 7.4$ Hz); ^{13}C nmr (DMSO): δ 55.53, 95.17, 114.30, 118.46, 121.90, 123.35, 123.58, 125.96, 129.15, 132.11, 137.24, 144.96, 161.29; ms: m/z 287 (M^+ , 40), 223 (86), 208 (100), 180 (38), 152 (14). *Anal.* calcd. for $\text{C}_{15}\text{H}_{13}\text{NO}_3\text{S}$: C, 62.70; H, 4.56; N, 4.87; S, 11.16. Found: C, 62.67; H, 4.58; N, 4.85; S, 11.19.

3-(4-Fluorophenyl)-4H-1,4-benzothiazine 1,1-dioxide (6c). Colourless crystals, 0.27 g (75%), mp 315–317 °C (EtOH); ir (nujol): 3300, 1310, 1115 cm^{-1} ; ^1H nmr (CDCl_3): δ 6.37 (s, 1H), 7.43–7.98 (m, 9H); ^{13}C nmr (CDCl_3): δ 96.3, 116.1 ($J_{\text{CF}} = 22.0$ Hz), 118.8, 122.1, 123.7, 123.7, 130.3 ($J_{\text{CF}} = 8.6$ Hz), 130.7, 132.4, 137.5, 144.6, 163.7 ($J_{\text{CF}} = 247.8$ Hz); ms: m/z 275 (M^+ , 40), 211 (100), 165 (19). *Anal.* calcd. for $\text{C}_{14}\text{H}_9\text{FNO}_2\text{S}$: C, 61.08; H, 3.66; N, 5.09; S, 11.65. Found: C, 61.05; H, 3.68; N, 5.11; S, 11.67.

3-(2-Thienyl)-4H-1,4-benzothiazine 1,1-dioxide (6d). White crystals, 0.30 g (81%), mp 276–278 °C (EtOH); ir (nujol): 3300, 1290, 1100 cm^{-1} ; ^1H nmr (CDCl_3): δ 6.44 (s, 1H), 7.38 (t, 1H, $J = 4.2$ Hz), 7.46 (t, 1H, $J = 7.2$ Hz), 7.69–7.78 (m, 2H), 7.89 (d, 1H, $J = 4.2$ Hz), 7.88–7.99 (m, 2H), 10.94 (s, 1H, D_2O exchangeable, NH); ^{13}C nmr (CDCl_3): δ 95.5, 118.4, 121.9, 123.8, 123.8, 128.4, 129.0, 129.8, 132.4, 135.4, 136.8, 139.0; ms: m/z 263 (M^+ , 62), 199 (100), 171 (17), 167 (15). *Anal.* calcd. for $\text{C}_{12}\text{H}_9\text{NO}_2\text{S}_2$: C, 54.73; H, 3.44; N, 5.32; S, 24.35. Found: C, 54.70; H, 3.48; N, 5.30; S, 24.30.

3-tert-Butyl-4H-1,4-benzothiazine 1,1-dioxide (6e). Colourless crystals, 0.22 g (70%), mp 260–262 °C (EtOH); ir (nujol): 3310, 1240, 1100 cm^{-1} ; ^1H nmr (DMSO): δ 1.39 (s, 9H, CH_3C), 5.91 (s, 1H), 7.37 (t, 1H, $J = 7.2$ Hz), 7.70 (m, 2H), 7.90 (d, 1H, $J =$

7.2 Hz); ^{13}C nmr (DMSO): δ 28.1, 35.9, 94.3, 118.0, 121.9, 122.8, 123.0, 132.1, 137.4, 154.3; ms: m/z 237 (M^+ , 37), 222 (30), 173 (14), 158 (100). *Anal.* calcd. for $\text{C}_{12}\text{H}_{15}\text{NO}_2\text{S}$: C, 60.73; H, 6.37; N, 5.90; S, 13.51. Found: C, 60.70; H, 6.39; N, 5.92; S, 13.54.

Acknowledgement. Financial support from the Ministero dell'Università e della Ricerca Scientifica e Tecnologica, (Rome) and from the University of Cagliari is gratefully acknowledged. One of us (Michele Usai) acknowledges financial support from Consorzio Interuniversitario Nazionale Metodologie e Processi Innovativi di Sintesi (C.I.N.M.P.I.S.) Bari.

REFERENCES

- [1] Thomas, L.; Gupta, A.; Gupta, V. *J. Fluor. Chem.* **2003**, *122*, 207-213, and references cited therein.
- [2] Armenise, D.; Trapani, G.; Arrivo, V.; Laraspata, E.; Morlacchi, F. *J. Heterocyclic Chem.* **2000**, *37*, 1611-1616, and references cited therein.
- [3] Schou, S. C.; Hansen, H. C.; Tagmose, T. M.; Boonen, H. C. M.; Worsaae, A.; Drabowski, M.; Wahl, P.; Arkhammar, P. O. G.; Bodvarsdottir, T.; Antoine, M. H.; Lebrun, P.; Hansen, J. B. *Bioorg. Med. Chem.* **2005**, *13*, 141-155.
- [4] Armenise, D.; Trapani, G.; Stasi, F.; Morlacchi, F. *Arch. Pharm. Pharm. Med. Chem.* **1998**, *331*, 54-58.
- [5] Rathore, B. S.; Kumar, M. *Bioorg. Med. Chem.* **2006**, *14*, 5678-5682.
- [6] Hamadi, M. Y.; Gupta, R.; Gupta, R. R. *J. Fluor. Chem.* **1999**, *94*, 169-174.
- [7] Sharma, P. R.; Gupta, V.; Gautam, D. C.; Gupta, R. R. *Phosphorus, Sulfur, Silicon Relat. Elem.* **2003**, *178*, 1483-1488.
- [8] Kumar, G.; Gupta, V.; Gautam, D. C.; Gupta, R. R. *Phosphorus, Sulfur, Silicon Relat. Elem.* **2004**, *179*, 1941-1948.
- [9] Lopez, S.E.; Salazar, J.; Rebollo, O.; Restrepo, J. *J. Heterocyclic Chem.* **2005**, *42*, 1007-1010.
- [10] Duhamel, L.; Plaquevent, J. C. *J. Org. Chem.* **1979**, *44*, 3404-3405.
- [11] Cabiddu, M.G.; Cabiddu, S.; Cadoni, E.; De Montis, S.; Fattuoni, C.; Melis, S. *Tetrahedron* **2003**, *59*, 2893-2897.
- [12] MacKenzie, N.E.; Thomson, R.H.; Greenhalgh, C.W. *J. Chem. Soc. Perkin I* **1980**, 2923-2932.