This article was downloaded by: [North West University] On: 21 December 2014, At: 05:04 Publisher: Taylor & Francis Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



# Organic Preparations and Procedures International: The New Journal for Organic Synthesis

Publication details, including instructions for authors and subscription information: http://www.tandfonline.com/loi/uopp20

# A FACILE SYNTHESIS OF NEW 2-PHOSPHONOBENZYLOXY-1,3,2-BENZOXAZA(DIOXA OR DIAZA)PHOSPHOLES

Jia Zhou<sup>a</sup> & Ruyu Chen<sup>a</sup>

<sup>a</sup> Institute of Elemento-Organic Chemistry, The State Key Laboratory of Elemento-Organic Chemistry, Nankai University , Tianjin, 300071, PR CHINA

Published online: 09 Feb 2009.

To cite this article: Jia Zhou & Ruyu Chen (1998) A FACILE SYNTHESIS OF NEW 2-PHOSPHONOBENZYLOXY-1,3,2-BENZOXAZA(DIOXA OR DIAZA)PHOSPHOLES, Organic Preparations and Procedures International: The New Journal for Organic Synthesis, 30:6, 716-719, DOI: <u>10.1080/00304949809355332</u>

To link to this article: http://dx.doi.org/10.1080/00304949809355332

### PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms &

Conditions of access and use can be found at <u>http://www.tandfonline.com/page/terms-and-conditions</u>

#### **OPPI BRIEFS**

- a) A. Mckillop, J. D. Hunt, R. D. Naylor and E. C. Taylor, J. Am. Chem. Soc., 93, 4918 (1971);
  b) S. B. Shim, K. Kim and Y. H. Kim, Tetrahedron Lett., 28, 645 (1987).
- 6. G. A. Olah, T. L. Ho, Synthesis, 610 (1976).
- 7. Y. Yukawa, M. Sakai, S. Suzuki, Bull. Chem. Soc. Jpn, 39, 2266 (1966).
- 8. P. Vankar, R. Rathore and S. Chandrasekaran, J. Org. Chem., 51, 3063 (1986).
- 9. J. G. Lee, K. H. Kwak and J. P. Hwang, Synth. Commun., 22, 2435 (1992).
- 10. H. Firouzabadi, N. Iranpoor, F. Kiaeezadeh and J. Toofan, Tetrahedron, 42, 719 (1986).
- 11. A. Wall, P. A. Ganeshpure and S. Satish, Bull. Chem. Soc. Jpn, 66, 1847 (1993).
- 12. B. Tamami and N. Goudarzian, Eur. Polym. J., 28, 1035 (1992).
- 13. H. Firouzabadi, E. Mottaghinejad and M. Seddighi, Synth. Commun., 19, 3469 (1989).
- 14. L. Singh and R. N. Kam, *ibid.*, 23, 3139 (1993).
- 15. H. Firouzabadi and I. Mohammadpoor-Baltork, ibid., 24, 489 (1994).
- J. Buckingham and S. M. Donaghy, Heilbron's "Dictionary of Organic Compounds", 5th ed., Chapman and Hall, New York, NY (1982).

\*\*\*\*\*\*

### A FACILE SYNTHESIS OF NEW 2-PHOSPHONOBENZYLOXY-1,3,2-BENZOXAZA(DIOXA OR DIAZA)PHOSPHOLES

Submitted by (04/06/98)

Jia Zhou\* and Ruyu Chen

Institute of Elemento-Organic Chemistry The State Key Laboratory of Elemento-Organic Chemistry Nankai University, Tianjin 300071, P. R. CHINA

Phosphorus-containing heterocyclic compounds have received much attention due to their biological importance.<sup>1-3</sup> Previously, we reported the synthesis of some thio (seleno) phosphate-phosphonate derivatives and their significant herbicidal, antiviral and fungicidal activities.<sup>4</sup> In continuation of our work of elaborating phosphate-phosphonates into a variety of heterocyclic systems of biological

**OPPI BRIEFS** 

interest, we herein describe an efficient, one-pot synthesis of 2-phosphonobenzyloxy-1,3,2benzoxaza(dioxa or diaza)phosphole 2-sulfides starting from the readily available synthon 1 (*Scheme*).



The phosphorylation of  $\alpha$ -hydroxyalkylphosphonates **1** was carried out with *tris*(diethylamino)phosphine activated by iodine<sup>5</sup> to give the resultant phosphite-phosphonates **3** in nearly quantitative yields under mild conditions. Intermediates **3** underwent a cyclization with pyrocatechol, *o*phenylenediamine or *o*-aminophenol to produce the cyclic phosphite-phosphonates **5**, which were then treated with sulfur to afford the expected title compounds **6a-d** in overall yields of 79.1-87.3%.

In summary, a convenient and efficient one-pot procedure has been developed for the synthesis of benzophospholes containing thio phosphate-phosphonate linkages in good yields under mild conditions. This method has considerable utility and can be extended to the syntheses of some new heterocyclic systems with potential biological interest.

#### **EXPERIMENTAL SECTION**

NMR spectra were taken on a Bruker AC-P200 spectrometer. Tetramethylsilane (TMS) was used as an internal standard for <sup>1</sup>H NMR, and 85%  $H_3PO_4$  was used as an external standard for <sup>31</sup>P NMR spectroscopy. The nuclei that are deshielded relative to their respective standards are assigned a positive chemical shift. IR spectra were recorded on a Shimadzu-435 spectrometer. Mass spectra were recorded on a Hewlett-Packard 5988 instrument. Elemental analyses were carried out on a Yana MT-3 instrument.  $\alpha$ -Hydroxyalkylphosphonates 1<sup>6</sup> and tris(diethylamino)phosphine<sup>7</sup> were prepared according to published methods.

Representative Experimental Procedure. 2-[(P,P-Diethoxy)phosphono(*p*-methyl)benzyloxy]-1,3,2-benzodioxaphosphole 2-Sulfide (6a).- A mixture of *tris*(diethylamino)phosphine (0.78 g, 3.15 mmol) and iodine (0.038 g, 0.15 mmol) in 50 mL of anhydrous benzene was heated at 70° in a stream of nitrogen for approximately 20 min until the precipitate dissolved. The solution was cooled to ambient temperature, and diethyl 1-hydroxy-1-(*p*-methyl)benzyl phosphonate (0.77 g, 3.0 mmol) was added. After having been maintained at 75° for 1.5 h, the reaction mixture was cooled to room temperature, and pyrocatechol (0.33 g, 3.0 mmol) was added. Then the reaction mixture was heated at 75° for 3 h. Transformation to the desired compound **6a** was accomplished by adding sulfur (0.15 g, 4.69 mmol) and keeping the mixture at 70° for 2.5 h. The solvent was removed *in vacuo*, and the compound was extracted from the residue with 10 mL of acetone. After filtration, the solvent was distilled off, and purification of the product was achieved with silica gel flash column chromatography using ethyl acetate/petroleum ether (bp. 60-90°) (1:4, v/v) as the eluent. This afforded **6a** as a colorless syrup (1.02 g, 79%). IR (film): 3190, 3031, 2909, 1602, 1499,1465, 1253 (s, P=O), 1053, 1016, 944, 881 (s, P-O-C), 830 (m), 754 (s), 714 (m, P=S) cm<sup>-1</sup>; <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  1.23 (t, 6 H, 2 CH<sub>3</sub>), 2.28 (ds, 3 H, CH<sub>3</sub>), 4.07 (m, 4 H, 2 CH<sub>2</sub>), 5.67 (dd, 1 H, CH, J = 14.0 Hz), 6.71-7.33 (m, 8 H, 2 C<sub>6</sub>H<sub>4</sub>); <sup>31</sup>P NMR (CDCl<sub>3</sub>);  $\delta$  15.77, 16.17, 69.69, 69.29 (two signals overlapped), 68.89, <sup>3</sup>J<sub>P1P2</sub> = 32.34 Hz; MS (EI) *m*/*z*: 428 (M<sup>+</sup>, 4.96%), 213 (15.1), 171 (13.09), 139 (12.82), 119 (24.49), 111 (17.24), 105 (100), 93 (22.26), 91 (35.67), 77 (14.9), 65(26.16).

Anal. Calcd for C<sub>18</sub>H<sub>22</sub>O<sub>6</sub>P<sub>2</sub>S: C, 50.46; H, 5.19. Found: C, 50.61; H, 5.42

**2-[(P,P-Diethoxy)phosphono**(*p*-methyl)benzyloxy]-1,3,2-benzodiazaphosphole 2-Sulfide (6b).- A colorless syrup; yield 87%. IR (film): 3388, 3127, 2971, 1607, 1511, 1490 (s), 1251 (s, P=O), 1091, 1049, 1015, 918 (s, P-O-C, P-N, C-N), 831 (s), 734 (s), 698 (s, P=S) cm<sup>-1</sup>; <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  1.22 (m, 6 H, 2 CH<sub>3</sub>), 2.34 (ds, 3 H, CH<sub>3</sub>), 3.92-4.18 (m, 4 H, 2 CH<sub>2</sub>), 5.76 (dd, 1 H, CH, J = 12.6 Hz), 5.78 (d, 1 H, NH, J = 18 Hz), 6.14 (d, 1 H, NH, J = 18 Hz), 6.68-7.36 (m, 8 H, 2 C<sub>6</sub>H<sub>4</sub>). <sup>31</sup>P NMR (CDCl<sub>3</sub>):  $\delta$  17.36, 16.69, 70.94, 70.27, <sup>3</sup>J<sub>P1P2</sub> = 24.25 Hz; MS (EI) *m/z*: 426 (M<sup>+</sup>, 19.12%), 290 (4.79), 242 (10.93), 213 (11.99), 185 (16.07), 169 (10.37), 154 (21.10), 137 (34.3), 119 (11.77), 109 (5.09), 105 (100), 93 (16.75), 91 (22.07), 65 (21.55).

Anal. Calcd for C<sub>18</sub>H<sub>24</sub>N<sub>2</sub>O<sub>4</sub>P<sub>2</sub>S: C, 50.70; H, 5.68; N, 6.57. Found: C, 50.98; H, 5.61; N, 6.86

**2-[(P,P-Diethoxy)phosphono**(*p*-methyl)benzyloxy]-1,3,2-benzoxazaphosphole 2-Sulfide (6c).- A colorless syrup; yield 82%. IR (film): 3407 (m, N-H), 2969, 1617, 1588, 1502, 1458 (s), 1252 (s, P=O), 1182, 1054, 1021, 953, 923 (s, P-O-C, P-N, C-N), 848 (s), 745 (s), 710 (s, P=S) cm<sup>-1</sup>; <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  1.16 (m, 6 H, 2 CH<sub>3</sub>), 2.32 (ds, 3 H, CH<sub>3</sub>), 2.83-4.52 (m, 4 H, 2 CH<sub>2</sub>), 5.58-5.92 (dq, 1 H, CH, J = 14.0 Hz), 6.54-6.83 (dd, 1 H, NH, J = 18.3 Hz), 6.63-7.42 (m, 8 H, 2 C<sub>6</sub>H<sub>4</sub>); <sup>31</sup>P NMR (CDCl<sub>3</sub>):  $\delta$  17.54 (d), 17.47 (d), 74.02 (d), 73.34 (d), <sup>3</sup>J<sub>P1P2</sub> = 42.22 Hz; MS (EI) *m/z*: 427 (M<sup>+</sup>, 8.42%), 392 (25.87), 364 (39.42), 272 (14.49), 241 (11.69), 213 (42.16), 180 (26.78), 138 (13.0), 121 (30.97), 108 (56.04), 105 (100), 93 (29.78), 91 (21.71), 80 (41.97), 74 (24.40), 65 (29.24).

Anal. Calcd for C18H23NO5P2S: C, 50.58; H, 5.44; N, 3.28. Found: C, 50.39; H, 5.52; N, 3.41

**2-[(P,P-Diethoxy)phosphono**(*p*-chloro)benzyloxy]-1,3,2-benzodiazaphosphole 2-Sulfide (6d).- A colorless syrup; yield 87%. IR (film): 3380 (s, NH), 3123, 2964, 1628, 1602, 1488 (s), 1247 (s, P=O), 1083, 1011, 971, 918 (s, P-O-C, P-N, C-N), 827 (s), 733 (s), 709 (s, P=S) cm<sup>-1</sup>; <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  1.17-1.34 (m, 6 H, 2 CH<sub>3</sub>), 3.96-4.17 (m, 4 H, 2 CH<sub>2</sub>), 5.79 (dd, 1 H, CH, J = 12.52 Hz), 5.82 (d, 1 H, NH, J = 20.84 Hz), 6.21 (d, 1 H, NH, J = 18.78 Hz), 6.68-7.42 (m, 8 H, 2 C<sub>6</sub>H<sub>4</sub>); <sup>31</sup>P NMR (CDCl<sub>3</sub>):  $\delta$  16.90, 16.61, 71.10, 70.81, <sup>3</sup>J<sub>P1P2</sub> = 23.50 Hz; MS (EI) *m*/z: 446 (M<sup>+</sup>, 29.24%), 290 (5.84), 262 (24.11), 233 (10.85), 206 (16.09), 185 (13.34), 169 (23.05), 154 (90.20), 137 (54.73), 127 (31.86), 125 (100), 121 (10.50), 105 (18.89), 93 (13.45), 77 (16.47), 65 (24.00).

Anal. Calcd for C<sub>17</sub>H<sub>21</sub>ClN<sub>2</sub>O<sub>4</sub>P<sub>2</sub>S: C, 45.69; H, 4.75; N, 6.27. Found: C, 45.53; H, 4.62; N, 6.53

Acknowledgement.- The authors thank the National Natural Science Foundation of China for financial support.

#### REFERENCES

- S. Y. Wu and J. E. Casida, *Phosphorus, Sulfur and Silicon*, **102**, 177 (1995); C. D. Reddy, M. S. Reddy and K. D. Berlin, *ibid.*, **102**, 103 (1995); A. Hirashima, *Agric. Bio. Chem.*, **53**, 175 (1989).
- H. L. Wang, J. Zhou, Y. G. Qiu, K. S. Feng and R. Y. Chen, *Phosphorus, Sulfur and Silicon*, 104, 135 (1995); R. Y. Chen and R. Bao, *Synthesis*, 618 (1989); R. Y. Chen and J. S. Tang, *Phosphorus, Sulfur and Silicon*, 48, 259 (1990); R. Y. Chen and L. F. Cheng, *ibid.*, 44, 193 (1989); R. Y. Chen and M. Zhang, *Science in China (Series B)*, 38, 152 (1995); *Chem. Abstr.*, 123, 144000f (1995).
- Y. Nishizuka, *Nature*, **308**, 693 (1984); V. L. Boyd, M. F. Summers, S. M. Ludeman, W. Egan, G. Zon and J. B. Regan, *J. Med. Chem.*, **30**, 366 (1987).
- 4. J. Zhou, R. Y. Chen and X. F. Yang, Heteroatom Chem., 9, 369 (1998).
- 5. H. Nöth and H. J. Vetter, Chem. Ber., 94, 1505 (1961).
- 6. F. Texier-Boullet and A. Foucaud, Synthesis, 916 (1982).
- 7. C. Stuebe and H. P. Lankelma, J. Am. Chem. Soc., 78, 976 (1956).

\*\*\*\*\*\*