

Tetrahedron Letters 39 (1998) 4667-4670

TETRAHEDRON LETTERS

Synthesis of Spirolactams from Tyrosine Amides and Related Substances

Norbert A. Braun,^a Marco A. Ciufolini,^{*,a} Karl Peters,^b Eva-Maria Peters^b

^aUniversité Claude Bernard Lyon 1 et Ecole Supérieure de Chimie, Physique, Electronique de Lyon 43, Boulevard du 11 Novembre 1918, 69622 Villeurbanne cedex, France

and

Department of Chemistry, MS 60, Rice University, 6100 Main Street, Houston, Texas 77005-1892, U.S.A. bMax-Planck-Institut für Festkörperforschung, Heisenbergstraße 1, 70506 Stuttgart, Germany

Received 16 March 1998; accepted 22 April 1998

Abstract

Oxidation of oxazolines derived from phenolic ω -arylalkanoic acids such as tyrosine with iodobenzene diacetate leads to spirocyclic amides in moderate yields. This reaction was heretofore unknown due to the propensity of free amide analogs of the oxazolines to furnish lactones upon oxidation. © 1998 Elsevier Science Ltd. All rights reserved.

Keywords: oxazolines, spiro compounds, cyclization, oxidation

Spirolactams of type 1 are useful intermediates in the synthesis of natural azaspirocyclic compounds. An appealing route to these heterocycles may be visualized in terms of oxidative spirocyclization of phenolic amides 3, followed by hydrogenation of the resulting dienone 2 (Scheme 1) [1-3]. Unfortunately, this transformation fails due to the propensity of the nucleophilic oxygen atom of the amide to intercept the electrophilic intermediate arising through activation of the phenol. Thus, oxidation of 3 with PhI(OCOCF₃)₂ (= PIFA) leads to lactone 5, probably by way of iminoether 4, which is readily hydrolyzed during isolation [4,5]. Work by Knapp [6] suggested that the effects responsible for the nucleophilicity of the oxygen atom of an amide should operate in favor of the nitrogen atom in imino derivatives



email: ciufi@cpe.fr; fax: (int) 33 (0)4 72 43 29 63

0040-4039/98/\$19.00 © 1998 Elsevier Science Ltd. All rights reserved. *PII:* S0040-4039(98)00874-0



(a) Ac₂O, 15% aq. NaOH, 93-95%; (b) aminoalcohol (see Table 1), BOP-Cl, CH₂Cl₂, Et₃N, 72-82%; (c) Burgess reagent, THF, 70°C, sealed tube, 68-74%; (d) K₂CO₃, MeOH, 73-81%; (e) PPh₃, CCl₄, Et₃N, 1:1 pyridine - MeCN, aminoalcohol (see Table 1), 44-68%.

thereof. We now report that the heretofore unknown transformation $3\rightarrow 2$ may indeed be achieved by oxidative cyclization of 2-oxazolines [7] derived from amides 3.

Oxazolines 10 were made either by dehydration [8] of the preformed amides 8 [9] with the Burgess reagent [10] (method A), or, more conveniently, by condensation of phenolic acids 6 with suitable ethanolamines as described by Vorbrüggen [11] (method B, Scheme 2 and Table 1). It is necessary to protect the phenol as an acetate during oxazoline formation by method A. Seemingly, condensation of the Burgess reagent with the phenolic OH results in polar byproducts that lower the yield of desired 10. No need for protection exists when method B is used.

Reaction of oxazolines 10 with iodobenzene diacetate under Kita-like conditions [4] provided the desired spirolactams 11. These were prone to undergo Michael-type cyclization to heterocycles 13a and/or 13b. Consequently, the crude 11 were converted immediately to the corresponding acetates 12 (Scheme 3). Interestingly, oxidation of 10 with PIFA produced mixtures containing only small amounts of desired 11. Probably the CF₃COOH liberated during these reactions damaged the acid-sensitive dienones. It is conceivable that the use of suitable acid traps (e.g. propylene oxide) in conjunction with PIFA might correct the



(a) PhI(OAc)₂, CF₃CH₂OH; (b) Ac₂O, pyridine, DMAP, (Table 1); (c) standing.

Entry	a	b	С	d	e
Acid	n = 1	n = 1	n = 1	n = 1	n = 2
6	R ¹ ⇒ H	R ¹ = H	R ¹ = NHTs	R ¹ = NHBOC	R ¹ = H
Amino-	Ph OH	∩OH	Phr H OH	Phr H OH	Ph H OH
alcohol	H ^W NH ₂	NH₂	NH2	NH2	NH2
Method	Α	в	В	Α	Α
Cpds.	n = 1	n = 1	n = 1	n = 1	n = 2
10 / 12	R ¹ = H; R ² = Bn	R ¹ = R ² = H	R ¹ = NHTs; R ² = Bn	R ¹ = NHBOC; R ² = Bn	R ¹ = H; R ² = Bn
Yield % of 12^a	47	42	41	22	17

^aRepresentative procedure for the cyclization of oxazolines **10**. A solution of $Phl(OAc)_2$ (1.1 mmol) in trifluoroethanol (5 mL) is added to a solution of oxazoline (1.0 mmol) in trifluoroethanol (5 mL) at rt. The mixture was stirred for 30 min, then solid NaHCO₃ was added to the solution and the mixture was filtered and concentrated. Standard acylation conditions (Ac₂O, pyridine, DMAP, rt) provided **12**.

problem; however, experiments in this direction have not yet been conducted. The structure of **12a** was confirmed by single crystal X-ray diffractometry.¹ A summary of representative experiments appears in Table 1, wherein the reported yields refer to purified acetates **12**.² In general, cyclization of oxazolines leading to a spiro[4.5] system proceeds in fair yields (40-50%), whereas the reaction of those giving a spiro[5.5] product is much less efficient (entry **e**). Tyrosine derivatives are good substrates for the new process, provided that the α -amino functionality is blocked as a sulfonamide (cf. entry **c**). In tyrosine *carbamates* (e.g., *N*-BOC compound **10d**), by contrast, the carbamate carbonyl competes effectively with the oxazoline nitrogen for capture of the electrophilic intermediate obtained by activation of the phenol. This results in several byproducts and in significantly lowered yields of **11** / **12**. Scrutiny of NMR spectra of crude **11** shows no evidence of byproducts resulting from aromatization of the oxazoline to an oxazole. The moderate yields of **11** / **12** seem to be rather attributable to

Table 1: representative formation and cyclization reactions of oxazolines 10

¹Crystal data for **12a** (C₂₀H₂₁NO₄; M = 339.39). Crystal dimension 0.25 x 0.25 x 1.4 mm; colorless needle; orthorhombic, space group $P2_{12}1_{21}$, a = 1008.8(1), b = 1102.5(1), c = 1610.3(3) pm, $V = 1791.0(4) \times 10^6$ pm³, Z = 4, d (calcd) = 1.295 g x cm⁻³, μ (Mo $K\alpha$) = 0.09 mm⁻¹, $\lambda = 0.71073$, $2\theta_{max} = 55^\circ$, 5564 reflections measured, 4124 unique and 2858 with $F > 3\sigma(F)$. Full matrix LSQ. Empirical absorption correction by Ψ scans. Hydrogen positions of riding model with fixed isotopic U. R(F) and R_W(F) = 0.066 and 0.061, respectively. Further details of the crystal structure may be obtained from the Director of the Cambridge Crystallographic Data Centre, 12 Union Road, Cambridge CB2 1EW (UK), by quoting the full journal citation.

²All compounds were fully characterized [¹H, ¹³C NMR (CDCl₃), IR, MS, $[\alpha]_D^{20}$ (CHCl₃), HRMS]. Data for representative compounds: **8a**: colorless crystals, mp 96 - 98 °C. ¹H: 2.29 (s, 3H), 2.28 - 2.47 (m, 2H), 2.77 (d, 7.6 Hz, 2H), 2.88 - 2.95 (m, 2H), 3.39 (dd, 3.5, 10.8 Hz, 1H), 3.45 (dd, 4.0, 10.8 Hz, 1H), 4.10 (m, 1H), 5.62 (d, 7.6 Hz, 1H), 6.97 & 7.18 (d, 8.5 Hz, each 2H), 7.14 - 7.30 (m, 5H). ¹³C: 21.1 (q), 31.3 (t), 36.8 (t), 38.8 (t), 52.2 (d), 63.5 (t), 121.7, 126.6, 128.5, 129.2 & 129.5 (d) 137.7, 138.3 & 149.0 (s), 170.2 (s), 172.1 (s). $[\alpha] = -14.7^{\circ}$ (c = 1.63). **10a**: colorless crystals, mp 141 - 143 °C. - ¹H (DMSO): 2.36 - 2.41 (m, 2H), 2.59 (dd, 7.1, 13.6 Hz, 1H), 2.67 - 2.72 (m, 2H), 2.80 (dd, 6.1, 13.6 Hz, 1H), 3.83 (dd, 7.3, 8.1 Hz, 1H), 4.14 (dd, 8.1, 9.4 Hz, 1H), 4.25 (m, 1H), 6.64 & 6.98 (d, 8.5 Hz, each 2H), 7.18 - 7.29 (m, 5H). ¹³C (DMSO): 29.7 (t), 30.8 (t), 41.4 (t), 66.8 (d), 71.1 (t), 115.3, 126.3, 128.3, 129.3 & 129.4 (d), 130.6, 138.5 & 156.1 (s), 165.9 (s). $[\alpha] = -26.8^{\circ}$ (c = 1.16, MeOH). **12a**: colorless needles, mp 163 - 165 °C. ¹H: 1.85 - 2.07 (m, 2H), 2.05 (s, 3H), 2.48 - 2.56 (m, 2H), 2.87 (dd, 4.9, 13.2 Hz, 1H), 3.12 (m, 1H), 3.47 (dd, 10.6, 13.2 Hz, 1H), 4.38 (dd, 5.4, 11.3 Hz, 1H), 4.53 (dd, 8.1, 11.3 Hz, 1H), 5.14 (dd, 3.1, 10.1 Hz, 1H), 5.85 & 6.21 (dd, each 1H, 2.1, 10.1 Hz, 1H), 6.78 (dd, 3.1, 10.1 Hz, 1H), 7.12 - 7.32 (m, 5H). ¹³C: 21.0 (q), 30.0 (t), 30.5 (t), 34.5 (t), 57.0 (d), 62.8 (s), 63.5 (t), 127.1 & 128.6 (d), 129.2 & 129.6 (d, C=C), 129.9 (d) 137.9 (s), 148.6 & 149.2 (d, C=C), 170.2 (s), 175.0 (s), 184.2 (s). [α] = + 31.4° (c = 0.44).



polymer formation. Finally, attempts to induce oxidative cyclization in solvents other than trifluoroethanol (e.g., MeCN) have so far been unsuccessful.

As exemplified in Scheme 4 with compound 12a, the dienone system may be hydrogenated at atmospheric pressure to give spirocycles of the type 1; e.g., 14. This reduction occurred cleanly in the presence of platinum on carbon or platinum oxide (Adams catalyst), whereas other hydrogenation catalysts provided also variable quantities of rearomatized product 15. The ratio of desired lactam 14 to amide 15 ranged from 100:0 with Pt(C) to 60:40 with Pd(C) to 0:100 with Rh(C). Aromatization presumably involves formal oxidative addition of zerovalent metal to the dienone C–N σ bond, followed by hydrogenolysis of the intermediate complex. Indeed, the extent of aromatization seems to correlate with the reduction potential of Rh (0.600 V), Pd (0.951 V) and Pt (1.118 V) [12]. A side reaction observed with Pt catalyst is hydrogenation, and it may be generally controlled by careful monitoring of the course of the reduction (TLC) and by avoiding prolonged reaction times.

In summary, the preparation of spirolactams 12 by oxidative cyclization of phenolic ω arylalkanoic acids, heretofore an elusive process, may be achieved *via* oxazoline intermediates. This chemistry is likely to facilitate the synthesis of various heterocyclic natural products. Additional developments in these areas will be described in due course.

Acknowledgment: We thank the NIH (CA-55268), the NSF (CHE 95-26183), the Robert A. Welch Foundation (C-1007), the CNRS and the MESR for support of our research program. MAC is a Fellow of the Alfred P. Sloan Foundation (1994-1998).

References

- [1] Alternative methods: Bryce MR, Gardiner JM, Horton PJ, Smith SA. J. Chem. Res. S 1989:1.
- [2] Other spirolactam syntheses: Boivin J, Yousfi M, Zard SZ. Tetrahedron Lett. 1997;38:5985-5988.
- [3] See also: Cossy J, Bouzide A. Tetrahedron 1997;53:5775-5784.
- [4] Kita Y, Tohma H, Kikuchi K, Inagaki M, Yakura T. J. Org. Chem. 1991;56:435-438.
- [5] Tamaru Y, Yakura T, Haruta J, Kita Y. J. Org. Chem. 1987;52:3927-3930.
- [6] Knapp S. Iodolactamization: Aspects and Applications. In: Pearson WH, editor. Advances in Heterocyclic Natural Product Synthesis, vol. 3. Greenwich, CT: JAI Press, 1996:57-98.
- [7] Review on 2-oxazoline chemistry: Gant TG, Meyers AI. Tetrahedron 1994;50:2297-2360.
- [8] Diago-Meseguer J, Palomo-Coll AL, Fernandez-Lizarbe JR, Zugaza-Bilbao A. Synthesis 1980:547-551.
- [9] Wipf P, Miller CP. Tetrahedron Lett. 1992;33:907-910.
- [10] Atkins GM, Burgess EM. J. Am. Chem. Soc. 1968;90:4744-4745.
- [11] Vorbrüggen H, Krolikiewicz, K. Tetrahedron Lett. 1981;22:4471-4474.
- [12] Vanysek P. Electrochemical Series. In: Lide DR, editor. Handbook of Chemistry and Physics, 78th edition. Boca Raton, FL: CRC Press, 1997:8/20-8/30.